

QUARKS

The u -, d -, and s -quark masses are estimates of so-called “current-quark masses,” in a mass-independent subtraction scheme such as \overline{MS} at a scale $\mu \approx 2$ GeV. The c - and b -quark masses are the “running” masses in the \overline{MS} scheme. For the b -quark we also quote the 1S mass. These can be different from the heavy quark masses obtained in potential models.

u

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

Mass $m = 1.5$ to 3.3 MeV ^[a] Charge = $\frac{2}{3} e$ $I_z = +\frac{1}{2}$
 $m_u/m_d = 0.35$ to 0.60

d

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

Mass $m = 3.5$ to 6.0 MeV ^[a] Charge = $-\frac{1}{3} e$ $I_z = -\frac{1}{2}$
 $m_s/m_d = 17$ to 22
 $\bar{m} = (m_u + m_d)/2 = 2.5$ to 5.0 MeV

s

$$I(J^P) = 0(\frac{1}{2}^+)$$

Mass $m = 105^{+25}_{-35}$ MeV ^[a] Charge = $-\frac{1}{3} e$ Strangeness = -1
 $m_s / ((m_u + m_d)/2) = 25$ to 30

c

$$I(J^P) = 0(\frac{1}{2}^+)$$

Mass $m = 1.27^{+0.07}_{-0.11}$ GeV Charge = $\frac{2}{3} e$ Charm = $+1$

b

$$I(J^P) = 0(\frac{1}{2}^+)$$

Charge = $-\frac{1}{3} e$ Bottom = -1

Mass $m = 4.20^{+0.17}_{-0.07}$ GeV (\overline{MS} mass)

t

$$I(J^P) = 0(\frac{1}{2}^+)$$

Charge = $\frac{2}{3} e$ Top = $+1$

Mass $m = 171.3 \pm 1.1 \pm 1.2$ GeV ^[b] (direct observation of top events)
 $\Gamma(Wb)/\Gamma(Wq(q = b, s, d)) = 0.99^{+0.09}_{-0.08}$

t DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	ρ (MeV/c)
$Wq (q = b, s, d)$			—
Wb			—
$\ell\nu_\ell$ anything	[c,d] $(9.4 \pm 2.4) \%$		—
$\gamma q (q=u,c)$	[e] $< 5.9 \times 10^{-3}$	95%	—
$\Delta T = 1$ weak neutral current ($T1$) modes			
$Zq (q=u,c)$	$T1$ [f] $< 3.7 \%$	95%	—

b' (4th Generation) Quark, Searches for

- Mass $m > 190$ GeV, CL = 95% ($p\bar{p}$, quasi-stable b')
- Mass $m > 199$ GeV, CL = 95% ($p\bar{p}$, neutral-current decays)
- Mass $m > 128$ GeV, CL = 95% ($p\bar{p}$, charged-current decays)
- Mass $m > 46.0$ GeV, CL = 95% (e^+e^- , all decays)

t' (4th Generation) Quark, Searches for

- Mass $m > 256$ GeV, CL = 95% ($p\bar{p}$, $t'\bar{t}'$ prod., $t' \rightarrow Wq$)

Free Quark Searches

All searches since 1977 have had negative results.

NOTES

- [a] The ratios m_u/m_d and m_s/m_d are extracted from pion and kaon masses using chiral symmetry. The estimates of u and d masses are not without controversy and remain under active investigation. Within the literature there are even suggestions that the u quark could be essentially massless. The s -quark mass is estimated from SU(3) splittings in hadron masses.
- [b] Based on published top mass measurements using data from Tevatron Run-I and Run-II. Including also the most recent unpublished results from Run-II, the Tevatron Electroweak Working Group reports a top mass of $173.1 \pm 0.6 \pm 1.1$ GeV. See the note "The Top Quark" in the Quark Particle Listings of this *Review*.
- [c] ℓ means e or μ decay mode, not the sum over them.
- [d] Assumes lepton universality and W -decay acceptance.
- [e] This limit is for $\Gamma(t \rightarrow \gamma q)/\Gamma(t \rightarrow Wb)$.
- [f] This limit is for $\Gamma(t \rightarrow Zq)/\Gamma(t \rightarrow Wb)$.