



$$I(J^P) = 0(0^-)$$

I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

B_c^+ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
6274.9 ± 0.8 OUR AVERAGE			
6274.28 ± 1.40 ± 0.32	¹ AAIJ	17L LHCB	$p\bar{p}$ at 7, 8 TeV
6274.0 ± 1.8 ± 0.4	² AAIJ	14AQ LHCB	$p\bar{p}$ at 7, 8 TeV
6276.28 ± 1.44 ± 0.36	³ AAIJ	13AS LHCB	$p\bar{p}$ at 7, 8 TeV
6273.7 ± 1.3 ± 1.6	⁴ AAIJ	12AV LHCB	$p\bar{p}$ at 7 TeV
6275.6 ± 2.9 ± 2.5	⁵ AALTONEN	08M CDF	$\rho\bar{p}$ at 1.96 TeV
6300 ± 14 ± 5	⁵ ABAZOV	08T D0	$\rho\bar{p}$ at 1.96 TeV
6400 ± 390 ± 130	⁶ ABE	98M CDF	$\rho\bar{p}$ at 1.8 TeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
6285.7 ± 5.3 ± 1.2	⁵ ABULENCIA	06C CDF	Repl. by AALTONEN 08M
6320 ± 60	⁷ ACKERSTAFF	98O OPAL	$e^+e^- \rightarrow Z$

¹ Measured using $B_c^+ \rightarrow J/\psi D^0 K^+$ decays.

² Uses $B_c^+ \rightarrow J/\psi p\bar{p}\pi^+$ decays.

³ AAIJ 13AS uses the $B_c^+ \rightarrow J/\psi D_s^+$.

⁴ AAIJ 12AV uses the $B_c^+ \rightarrow J/\psi\pi^+$ mode and also measures the mass difference $M(B_c^+) - M(B^+) = 994.6 \pm 1.3 \pm 0.6 \text{ MeV}/c^2$.

⁵ Measured using a fully reconstructed decay mode of $B_c \rightarrow J/\psi\pi$.

⁶ ABE 98M observed $20.4^{+6.2}_{-5.5}$ events in the $B_c^+ \rightarrow J/\psi(1S)\ell\nu\ell$ with a significance of > 4.8 standard deviations. The mass value is estimated from $m(J/\psi(1S)\ell)$.

⁷ ACKERSTAFF 98O observed 2 candidate events in the $B_c^+ \rightarrow J/\psi(1S)\pi^+$ channel with an estimated background of 0.63 ± 0.20 events.

B_c^+ MEAN LIFE

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFLAV) and are described at <http://www.slac.stanford.edu/xorg/hflav/>. The averaging/rescaling procedure takes into account correlations between the measurements.

VALUE (10^{-12} s)	DOCUMENT ID	TECN	COMMENT
0.507 ± 0.009 OUR EVALUATION			
0.507 ± 0.009 OUR AVERAGE			
0.5134 ± 0.0110 ± 0.0057	^{1,2} AAIJ	15G LHCB	$p\bar{p}$ at 7, 8 TeV
0.509 ± 0.008 ± 0.012	³ AAIJ	14G LHCB	$p\bar{p}$ at 8 TeV
0.452 ± 0.048 ± 0.027	² AALTONEN	13 CDF	$\rho\bar{p}$ at 1.96 TeV
0.448 $^{+0.038}_{-0.036}$ ± 0.032	⁴ ABAZOV	09H D0	$\rho\bar{p}$ at 1.96 TeV
0.463 $^{+0.073}_{-0.065}$ ± 0.036	⁴ ABULENCIA	06O CDF	$\rho\bar{p}$ at 1.96 TeV
0.46 $^{+0.18}_{-0.16}$ ± 0.03	⁴ ABE	98M CDF	$\rho\bar{p}$ 1.8 TeV

¹ Also measures the width difference $\Delta\Gamma = \Gamma_{B_c^+} - \Gamma_{B^+} = 4.46 \pm 0.14 \pm 0.07 \text{ mm}^{-1} c$.

² Uses fully reconstructed $B_c^+ \rightarrow J/\psi \pi^+$ decays.

³ Measured using $B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu X$ decays.

⁴ The lifetime is measured from the $J/\psi e$ decay vertices.

B_c^+ DECAY MODES $\times B(\bar{b} \rightarrow B_c)$

B_c^- modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)	Confidence level
The following quantities are not pure branching ratios; rather the fraction $\Gamma_i/\Gamma \times B(\bar{b} \rightarrow B_c)$.		
Γ_1 $J/\psi(1S)\ell^+\nu_\ell$ anything	$(5.2^{+2.4}_{-2.1}) \times 10^{-5}$	
Γ_2 $J/\psi(1S)\mu^+\nu_\mu$		
Γ_3 $J/\psi(1S)\pi^+$	seen	
Γ_4 $J/\psi(1S)K^+$	seen	
Γ_5 $J/\psi(1S)\pi^+\pi^+\pi^-$	seen	
Γ_6 $J/\psi(1S)a_1(1260)$	$< 1.2 \times 10^{-3}$	90%
Γ_7 $J/\psi(1S)K^+K^-\pi^+$	seen	
Γ_8 $J/\psi(1S)\pi^+\pi^+\pi^+\pi^-\pi^-$	seen	
Γ_9 $\psi(2S)\pi^+$	seen	
Γ_{10} $J/\psi(1S)D^0K^+$	seen	
Γ_{11} $J/\psi(1S)D^*(2007)^0K^+$	seen	
Γ_{12} $J/\psi(1S)D^*(2010)^+K^{*0}$	seen	
Γ_{13} $J/\psi(1S)D^+K^{*0}$	seen	
Γ_{14} $J/\psi(1S)D_s^+$	seen	
Γ_{15} $J/\psi(1S)D_s^{*+}$	seen	
Γ_{16} $J/\psi(1S)p\bar{p}\pi^+$	seen	
Γ_{17} $\chi_c^0\pi^+$	$(2.4^{+0.9}_{-0.8}) \times 10^{-5}$	
Γ_{18} $p\bar{p}\pi^+$	not seen	
Γ_{19} D^0K^+		
Γ_{20} $D^0\pi^+$		
Γ_{21} $D^{*0}\pi^+$		
Γ_{22} $D^{*0}K^+$		
Γ_{23} $D^*(2010)^+\bar{D}^0$	$< 6.2 \times 10^{-3}$	90%
Γ_{24} D^+K^{*0}	$< 0.20 \times 10^{-6}$	90%
Γ_{25} $D^+\bar{K}^{*0}$	$< 0.16 \times 10^{-6}$	90%
Γ_{26} $D_s^+K^{*0}$	$< 0.28 \times 10^{-6}$	90%
Γ_{27} $D_s^+\bar{K}^{*0}$	$< 0.4 \times 10^{-6}$	90%

Γ_{28}	$D_s^+ \phi$	< 0.32	$\times 10^{-6}$	90%
Γ_{29}	$K^+ K^0$	< 4.6	$\times 10^{-7}$	90%
Γ_{30}	$B_s^0 \pi^+ / B(\bar{b} \rightarrow B_s)$	$(2.37^{+0.37}_{-0.35}) \times 10^{-3}$		

B_c^+ BRANCHING RATIOS

$\Gamma(J/\psi(1S)\ell^+ \nu_\ell \text{ anything}) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_1 / \Gamma \times B$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
8.1±1.2 OUR AVERAGE Error includes scale factor of 1.3.				
$8.6 \pm 1.0 \pm 0.3$		^{1,2} AALTONEN	16A CDF	$p\bar{p}$ at 1.96 TeV
$5.2^{+2.4}_{-2.1}$		³ ABE	98M CDF	$p\bar{p}$ 1.8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<16	90	⁴ ACKERSTAFF	98O OPAL	$e^+e^- \rightarrow Z$
<19	90	⁵ ABREU	97E DLPH	$e^+e^- \rightarrow Z$
<12	90	⁶ BARATE	97H ALEP	$e^+e^- \rightarrow Z$

¹ AALTONEN 16A reports $[\Gamma(B_c^+ \rightarrow J/\psi(1S)\ell^+ \nu_\ell \text{ anything}) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow J/\psi(1S)K^+)] = 0.211 \pm 0.012^{+0.021}_{-0.020}$ which we multiply by our best values $B(\bar{b} \rightarrow B^+) = (40.5 \pm 0.6) \times 10^{-2}$, $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.010 \pm 0.029) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² AALTONEN 16A also measures the cross-section $\sigma(B_c) \times B(B_c \rightarrow J/\psi \mu \nu_\mu) = 0.60 \pm 0.09$ nb and estimates the total cross-section $\sigma(B_c)$ to be in the range 25 ± 4 to 52 ± 8 nb for $p_T(B_c) > 6$ GeV/c and $|y(B_c)| < 1$.

³ ABE 98M result is derived from the measurement of $[\sigma(B_c) \times B(B_c \rightarrow J/\psi(1S)\ell \nu_\ell)] / [\sigma(B^+) \times B(B^+ \rightarrow J/\psi(1S)K^+)] = 0.132^{+0.041}_{-0.037}(\text{stat}) \pm 0.031(\text{sys})^{+0.032}_{-0.020}(\text{lifetime})$ by using PDG 98 values of $B(b \rightarrow B^+)$ and $B(B^+ \rightarrow J/\psi(1S)K^+)$.

⁴ ACKERSTAFF 98O reports $B(Z \rightarrow B_c X) / B(Z \rightarrow qq) \times B(B_c \rightarrow J/\psi(1S)\ell \nu_\ell) < 6.95 \times 10^{-5}$ at 90%CL. We rescale to our PDG 98 values of $B(Z \rightarrow b\bar{b})$.

⁵ ABREU 97E value listed is for an assumed $\tau_{B_c} = 0.4$ ps and improves to 1.6×10^{-4} for $\tau_{B_c} = 1.4$ ps.

⁶ BARATE 97H reports $B(Z \rightarrow B_c X) / B(Z \rightarrow qq) \cdot B(B_c \rightarrow J/\psi(1S)\ell \nu_\ell) < 5.2 \times 10^{-5}$ at 90%CL. We rescale to our PDG 96 values of $B(Z \rightarrow b\bar{b})$. A $B_c^+ \rightarrow J/\psi(1S)\mu^+ \nu_\mu$ candidate event is found, compared to all the known background sources 2×10^{-3} , which gives $m_{B_c} = 5.96^{+0.25}_{-0.19}$ GeV and $\tau_{B_c} = 1.77 \pm 0.17$ ps.

$\Gamma(J/\psi(1S)\pi^+) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_3 / \Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
seen		¹ AAIJ	15M LHCB	pp at 8 TeV
seen		² KHACHATRY...15AA	CMS	pp at 7 TeV
seen		AALTONEN	13 CDF	$p\bar{p}$ at 1.96 TeV
seen		³ AAIJ	12AV LHCB	pp at 7 TeV
seen		AALTONEN	08M CDF	$p\bar{p}$ at 1.96 TeV
seen		ABAZOV	08T D0	$p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2.4 \times 10^{-4}$	90	⁴ ACKERSTAFF 98O	OPAL	$e^+e^- \rightarrow Z$
$<3.4 \times 10^{-4}$	90	⁵ ABREU	97E	DLPH $e^+e^- \rightarrow Z$
$<8.2 \times 10^{-5}$	90	⁶ BARATE	97H	ALEP $e^+e^- \rightarrow Z$
$<2.0 \times 10^{-5}$	95	⁷ ABE	96R	CDF $p\bar{p}$ 1.8 TeV

¹ AAIJ 15M reports a measurement of $B(B_c^+ \rightarrow J/\psi\pi^+) / B(B^+ \rightarrow J/\psi K^+) \cdot f_c/f_u = (0.683 \pm 0.018 \pm 0.009)\%$ at $p_T(B) < 20$ GeV and $2.0 < y(B) < 4.5$.

² KHACHATRYAN 15AA reports a measurement of $B(B_c^+ \rightarrow J/\psi\pi^+) / B(B^+ \rightarrow J/\psi K^+) \cdot f_c/f_u = (0.48 \pm 0.05 \pm 0.03 \pm 0.05)\%$, at $p_T > 15$ GeV and $|\eta(B)| < 1.6$.

³ AAIJ 12AV reports a measurement of $B(B_c^+ \rightarrow J/\psi\pi^+)/B(B^+ \rightarrow J/\psi K^+) f_c/f_u = (0.68 \pm 0.10 \pm 0.03 \pm 0.05)\%$ at $p_T(B) > 4$ GeV and $2.5 < \eta(B) < 4.5$.

⁴ ACKERSTAFF 98O reports $B(Z \rightarrow B_c X)/B(Z \rightarrow qq) \times B(B_c \rightarrow J/\psi(1S)\pi^+) < 1.06 \times 10^{-4}$ at 90%CL. We rescale to our PDG 98 values of $B(Z \rightarrow b\bar{b})$.

⁵ ABREU 97E value listed is for an assumed $\tau_{B_c} = 0.4$ ps and improves to 2.7×10^{-4} for $\tau_{B_c} = 1.4$ ps.

⁶ BARATE 97H reports $B(Z \rightarrow B_c X)/B(Z \rightarrow qq) \cdot B(B_c \rightarrow J/\psi(1S)\pi) < 3.6 \times 10^{-5}$ at 90%CL. We rescale to our PDG 96 values of $B(Z \rightarrow b\bar{b})$.

⁷ ABE 96R reports $B(b \rightarrow B_c X)/B(b \rightarrow B^+ X) \cdot B(B_c^+ \rightarrow J/\psi(1S)\pi^+)/B(B^+ \rightarrow J/\psi(1S)K^+) < 0.053$ at 95%CL for $\tau_{B_c} = 0.8$ ps. It changes from 0.15 to 0.04 for $0.17 \text{ ps} < \tau_{B_c} < 1.6$ ps. We rescale to our PDG 96 values of $B(b \rightarrow B^+) = 0.378 \pm 0.022$ and $B(B^+ \rightarrow J/\psi(1S)K^+) = 0.00101 \pm 0.00014$.

$\Gamma(J/\psi(1S)\pi^+)/\Gamma(J/\psi(1S)\mu^+\nu_\mu)$ Γ_3/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
$(4.69 \pm 0.28 \pm 0.46) \times 10^{-2}$	¹ AAIJ	14W	LHCB pp at 7 TeV

¹ AAIJ 14W reports also a measurement $B(B_c^+ \rightarrow J/\psi\pi^+) / B(B_c^+ \rightarrow J/\psi\mu^+\nu_\mu) = 0.271 \pm 0.016 \pm 0.016$ in the region $m_{J/\psi\mu^+} > 5.3$ GeV.

$\Gamma(J/\psi(1S)K^+)/\Gamma(J/\psi(1S)\pi^+)$ Γ_4/Γ_3

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.079 \pm 0.007 \pm 0.003$		AAIJ	16AF	LHCB pp at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.069 \pm 0.019 \pm 0.005$	50	AAIJ	13BY	LHCB Repl. by AAIJ 16AF
-----------------------------	----	------	------	-------------------------

$\Gamma(J/\psi(1S)\pi^+\pi^+\pi^-)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_5/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
seen		AAIJ	12Y	LHCB pp at 7 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<5.7 \times 10^{-4}$	90	¹ ABREU	97E	DLPH $e^+e^- \rightarrow Z$
-----------------------	----	--------------------	-----	-----------------------------

¹ ABREU 97E value listed is independent of $0.4 \text{ ps} < \tau_{B_c} < 1.4$ ps.

$\Gamma(J/\psi(1S)\pi^+\pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+)$ Γ_5/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
2.4 ± 0.4 OUR AVERAGE			
$2.55 \pm 0.80 \pm 0.33^{+0.04}_{-0.01}$	KHACHATRY...15AA	CMS	pp at 7 TeV
$2.41 \pm 0.30 \pm 0.33$	AAIJ	12Y	LHCB pp at 7 TeV

$\Gamma(J/\psi(1S) a_1(1260))/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_6/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2 \times 10^{-3}$	90	¹ ACKERSTAFF 980	OPAL	$e^+ e^- \rightarrow Z$

¹ ACKERSTAFF 980 reports $B(Z \rightarrow B_c X)/B(Z \rightarrow qq) \times B(B_c \rightarrow J/\psi(1S) a_1(1260)) < 5.29 \times 10^{-4}$ at 90%CL. We rescale to our PDG 98 values of $B(Z \rightarrow b\bar{b})$.

 $\Gamma(J/\psi(1S) K^+ K^- \pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_7/\Gamma \times B$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	13CA LHCB	pp at 7, 8 TeV

¹ A signal yield of 78 ± 14 decays is reported with a significance of 6.2 standard deviations using an integrated luminosity of 3 fb^{-1} data.

 $\Gamma(J/\psi(1S) K^+ K^- \pi^+)/\Gamma(J/\psi(1S) \pi^+)$ Γ_7/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
$0.53 \pm 0.10 \pm 0.05$	¹ AAIJ	13CA LHCB	pp at 7, 8 TeV

¹ A signal yield of 78 ± 14 decays is reported with a significance of 6.2 standard deviations using an integrated luminosity of 3 fb^{-1} data.

 $\Gamma(J/\psi(1S) \pi^+ \pi^+ \pi^+ \pi^- \pi^-)/\Gamma(J/\psi(1S) \pi^+)$ Γ_8/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
$1.74 \pm 0.44 \pm 0.24$	¹ AAIJ	14P LHCB	pp at 7, 8 TeV

¹ A signal yield of 32 ± 8 decays is reported with a significance of 4.5 standard deviations.

 $\Gamma(\psi(2S) \pi^+)/\Gamma(J/\psi(1S) \pi^+)$ Γ_9/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
$0.268 \pm 0.032 \pm 0.007 \pm 0.006$	¹ AAIJ	15AY LHCB	pp at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.250 \pm 0.068 \pm 0.014 \pm 0.006$	¹ AAIJ	13AMLHCB	Repl. by AAIJ 15AY
---------------------------------------	-------------------	----------	--------------------

¹ The last uncertainty is due to the uncertainty of the $B(\psi(2S) \rightarrow \mu^+ \mu^-)/B(J/\psi \rightarrow \mu^+ \mu^-)$ ratio measurement.

 $\Gamma(J/\psi(1S) D^0 K^+)/\Gamma(J/\psi(1S) \pi^+)$ Γ_{10}/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
$0.432 \pm 0.136 \pm 0.028$	AAIJ	17L LHCB	pp at 7, 8 TeV

 $\Gamma(J/\psi(1S) D^*(2007)^0 K^+)/\Gamma(J/\psi(1S) D^0 K^+)$ Γ_{11}/Γ_{10}

VALUE	DOCUMENT ID	TECN	COMMENT
$5.1 \pm 1.8 \pm 0.4$	AAIJ	17L LHCB	pp at 7, 8 TeV

 $\Gamma(J/\psi(1S) D^*(2010)^+ K^{*0})/\Gamma(J/\psi(1S) D^0 K^+)$ Γ_{12}/Γ_{10}

VALUE	DOCUMENT ID	TECN	COMMENT
$2.10 \pm 1.08 \pm 0.34$	AAIJ	17L LHCB	pp at 7, 8 TeV

 $\Gamma(J/\psi(1S) D^+ K^{*0})/\Gamma(J/\psi(1S) D^0 K^+)$ Γ_{13}/Γ_{10}

VALUE	DOCUMENT ID	TECN	COMMENT
$0.63 \pm 0.39 \pm 0.08$	AAIJ	17L LHCB	pp at 7, 8 TeV

$\Gamma(J/\psi(1S)D_s^+)/\Gamma(J/\psi(1S)\pi^+)$ Γ_{14}/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
3.1 ± 0.5 OUR AVERAGE			
3.8 ± 1.1 ± 0.4	AAD	16H ATLS	pp at 7, 8 TeV
2.90 ± 0.57 ± 0.24	AAIJ	13AS LHCB	pp at 7, 8 TeV

 $\Gamma(J/\psi(1S)D_s^{*+})/\Gamma(J/\psi(1S)\pi^+)$ Γ_{15}/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
10.4 ± 3.1 ± 1.6	AAD	16H ATLS	pp at 7, 8 TeV

 $\Gamma(J/\psi(1S)D_s^{*+})/\Gamma(J/\psi(1S)D_s^+)$ Γ_{15}/Γ_{14}

VALUE	DOCUMENT ID	TECN	COMMENT
2.5 ± 0.5 OUR AVERAGE			
2.8 $^{+1.2}_{-0.8}$ ± 0.3	AAD	16H ATLS	pp at 7, 8 TeV
2.37 ± 0.56 ± 0.10	AAIJ	13AS LHCB	pp at 7, 8 TeV

 $\Gamma(J/\psi(1S)p\bar{p}\pi^+)/\Gamma(J/\psi(1S)\pi^+)$ Γ_{16}/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
0.143 $^{+0.041}_{-0.036}$	AAIJ	14AQ LHCB	pp at 7, 8 TeV

 $\Gamma(\chi_c^0\pi^+)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
24.2 $^{+8.6}_{-7.7}$ ± 0.4	^{1,2} AAIJ	16AT LHCB	pp at 7 and 8 TeV

¹AAIJ 16AT reports $[\Gamma(B_c^+ \rightarrow \chi_c^0\pi^+)/\Gamma_{\text{total}}] \times [\Gamma(\bar{b} \rightarrow B^+)/\Gamma_{\text{total}}] = (9.8^{+3.4}_{-3.0} \pm 0.8) \times 10^{-6}$ which we divide by our best value $\Gamma(\bar{b} \rightarrow B^+)/\Gamma_{\text{total}} = 0.405 \pm 0.006$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²The significance of the observed signal is 4.0 standard deviations.

 $\Gamma(p\bar{p}\pi^+)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	¹ AAIJ	16K LHCB	pp at 7, 8 TeV

¹Measures the ratio $(f_c/f_u) \times B(B_c^+ \rightarrow p\bar{p}\pi^+) < 3.6 \times 10^{-8}$ at 95% CL, in the region $m(p\bar{p}) < 2.85 \text{ GeV}/c^2$, where f_c (f_u) represents the fragmentation fraction of the b -quark into the B_c^+ (B_u^+) meson.

 $\Gamma(D^0K^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{19}/\Gamma \times B$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
3.8 $^{+1.2}_{-1.0}$ ± 0.1	¹ AAIJ	17AG LHCB	pp at 7, 8 TeV

¹AAIJ 17AG reports $[\Gamma(B_c^+ \rightarrow D^0K^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] = (9.3^{+2.8}_{-2.5} \pm 0.6) \times 10^{-7}$ which we multiply by our best value $B(\bar{b} \rightarrow B^+) = (40.5 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D^0\pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{20}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.6 \times 10^{-7}$	95	¹ AAIJ	17AG LHCB	pp at 7, 8 TeV

¹AAIJ 17AG reports $[\Gamma(B_c^+ \rightarrow D^0\pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 3.9 \times 10^{-7}$ which we multiply by our best value $B(\bar{b} \rightarrow B^+) = 40.5 \times 10^{-2}$.

$\Gamma(D^{*0}\pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{21}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4 \times 10^{-7}$	95	¹ AAIJ	17AG LHCB	pp at 7, 8 TeV

¹AAIJ 17AG reports $[\Gamma(B_c^+ \rightarrow D^{*0}\pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 1.1 \times 10^{-6}$ which we multiply by our best value $B(\bar{b} \rightarrow B^+) = 40.5 \times 10^{-2}$.

$\Gamma(D^{*0}K^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{22}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4 \times 10^{-7}$	95	¹ AAIJ	17AG LHCB	pp at 7, 8 TeV

¹AAIJ 17AG reports $[\Gamma(B_c^+ \rightarrow D^{*0}K^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 1.1 \times 10^{-6}$ which we multiply by our best value $B(\bar{b} \rightarrow B^+) = 40.5 \times 10^{-2}$.

$\Gamma(D^*(2010)^+\bar{D}^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{23}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.2 \times 10^{-3}$	90	¹ BARATE	98Q ALEP	$e^+e^- \rightarrow Z$

¹BARATE 98Q reports $B(Z \rightarrow B_c X) \times B(B_c \rightarrow D^*(2010)^+\bar{D}^0) < 1.9 \times 10^{-3}$ at 90%CL. We rescale to our PDG 98 values of $B(Z \rightarrow b\bar{b})$.

$\Gamma(D^+K^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{24}/\Gamma \times B$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<0.20	90	¹ AAIJ	13R LHCB	pp at 7 TeV

¹AAIJ 13R reports $[\Gamma(B_c^+ \rightarrow D^+K^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 0.5 \times 10^{-6}$ which we multiply by our best value $B(\bar{b} \rightarrow B^+) = 40.5 \times 10^{-2}$.

$\Gamma(D^+\bar{K}^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{25}/\Gamma \times B$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<0.16	90	¹ AAIJ	13R LHCB	pp at 7 TeV

¹AAIJ 13R reports $[\Gamma(B_c^+ \rightarrow D^+\bar{K}^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 0.4 \times 10^{-6}$ which we multiply by our best value $B(\bar{b} \rightarrow B^+) = 40.5 \times 10^{-2}$.

$\Gamma(D_s^+K^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{26}/\Gamma \times B$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<0.28	90	¹ AAIJ	13R LHCB	pp at 7 TeV

¹AAIJ 13R reports $[\Gamma(B_c^+ \rightarrow D_s^+K^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 0.7 \times 10^{-6}$ which we multiply by our best value $B(\bar{b} \rightarrow B^+) = 40.5 \times 10^{-2}$.

$\Gamma(D_s^+ \bar{K}^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{27}/\Gamma \times B$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<0.4	90	¹ AAIJ	13R LHCB	pp at 7 TeV

¹ AAIJ 13R reports $[\Gamma(B_c^+ \rightarrow D_s^+ \bar{K}^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 1.1 \times 10^{-6}$ which we multiply by our best value $B(\bar{b} \rightarrow B^+) = 40.5 \times 10^{-2}$.

$\Gamma(D_s^+ \phi)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{28}/\Gamma \times B$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<0.32	90	¹ AAIJ	13R LHCB	pp at 7 TeV

¹ AAIJ 13R reports $[\Gamma(B_c^+ \rightarrow D_s^+ \phi)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 0.8 \times 10^{-6}$ which we multiply by our best value $B(\bar{b} \rightarrow B^+) = 40.5 \times 10^{-2}$.

$\Gamma(K^+ K^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{29}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.6 $\times 10^{-7}$	90	¹ AAIJ	13BS LHCB	pp at 7 TeV

¹ Derived from $\Gamma(K^+ K^0)/\Gamma \times B(\bar{b} \rightarrow B_c) / (B(B^+ \rightarrow K^0 \pi^+) B(\bar{b} \rightarrow B^+)) < 5.8\%$ at 90% CL using normalization mode $B(B^+ \rightarrow K^0 \pi^+) = (23.97 \pm 0.53 \pm 0.71) \times 10^{-6}$ and assuming a B production ratio $f(\bar{b} \rightarrow B_u^+) = 0.33$.

$\Gamma(B_s^0 \pi^+ / B(\bar{b} \rightarrow B_s)) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_{30}/\Gamma \times B$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$2.37 \pm 0.31 \pm 0.11^{+0.17}_{-0.13}$	¹ AAIJ	13BU LHCB	pp at 7, 8 TeV

¹ The last uncertainty is due to the uncertainty of the B_c^+ lifetime measurement.

POLARIZATION IN B_c^+ DECAY

In decays involving two vector mesons, one can distinguish among the states in which meson polarizations are both longitudinal (L) or both are transverse and parallel (\parallel) or perpendicular (\perp) to each other with the parameters Γ_L/Γ , Γ_{\perp}/Γ , and the relative phases ϕ_{\parallel} and ϕ_{\perp} . See the definitions in the note on “Polarization in B Decays” review in the B^0 Particle Listings.

Γ_L/Γ in $B_c^+ \rightarrow J/\psi D_s^{*+}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.54 ± 0.15 OUR AVERAGE			
0.62 ± 0.24	¹ AAD	16H ATLS	pp at 7, 8 TeV
0.48 ± 0.20	² AAIJ	13AS LHCB	pp at 7, 8 TeV

¹ AAD 16H measures $1 - \Gamma_L/\Gamma = 0.38 \pm 0.24$.

² AAIJ 13AS measures $1 - \Gamma_L/\Gamma = 0.52 \pm 0.20$.

B_c^+ REFERENCES

AAIJ	17AG	PRL 118 111803	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17L	PR D95 032005	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAD	16H	EPJ C76 4	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAIJ	16AF	JHEP 1609 153	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	16AT	PR D94 091102	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	16K	PL B759 313	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	16A	PR D93 052001	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AAIJ	15AY	PR D92 072007	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15G	PL B742 29	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15M	PRL 114 132001	R. Aaij <i>et al.</i>	(LHCb Collab.)
KHACHATRY...	15AA	JHEP 1501 063	V. Khachatryan <i>et al.</i>	(CMS Collab.)
AAIJ	14AQ	PRL 113 152003	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14G	EPJ C74 2839	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14P	JHEP 1405 148	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14W	PR D90 032009	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13AM	PR D87 071103	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13AS	PR D87 112012	R. Aaij <i>et al.</i>	(LHCb Collab.)
Also		PR D89 019901 (errat.)	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13BS	PL B726 646	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13BU	PRL 111 181801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13BY	JHEP 1309 075	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13CA	JHEP 1311 094	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13R	JHEP 1302 043	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	13	PR D87 011101	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AAIJ	12AV	PRL 109 232001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12Y	PRL 108 251802	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABAZOV	09H	PRL 102 092001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
AALTONEN	08M	PRL 100 182002	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	08T	PRL 101 012001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABULENCIA	06C	PRL 96 082002	A. Abulencia <i>et al.</i>	(CDF Collab.)
ABULENCIA	06O	PRL 97 012002	A. Abulencia <i>et al.</i>	(CDF Collab.)
ABE	98M	PRL 81 2432	F. Abe <i>et al.</i>	(CDF Collab.)
Also		PR D58 112004	F. Abe <i>et al.</i>	(CDF Collab.)
ACKERSTAFF	98O	PL B420 157	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
BARATE	98Q	EPJ C4 387	R. Barate <i>et al.</i>	(ALEPH Collab.)
PDG	98	EPJ C3 1	C. Caso <i>et al.</i>	(PDG Collab.)
ABREU	97E	PL B398 207	P. Abreu <i>et al.</i>	(DELPHI Collab.)
BARATE	97H	PL B402 213	R. Barate <i>et al.</i>	(ALEPH Collab.)
ABE	96R	PRL 77 5176	F. Abe <i>et al.</i>	(CDF Collab.)
PDG	96	PR D54 1	R. M. Barnett <i>et al.</i>	(PDG Collab.)