

$$I^G(J^{PC}) = 0^-(1^{--})$$

See the Review on “ $\psi(2S)$ and χ_c branching ratios” before the $\chi_{c0}(1P)$ Listings.

$\psi(2S)$ MASS

OUR FIT includes measurements of $m_{\psi(2S)}$, $m_{\psi(3770)}$, and $m_{\psi(3770)} - m_{\psi(2S)}$.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3686.097 ± 0.025 OUR FIT				Error includes scale factor of 2.6.
3686.097 ± 0.010 OUR AVERAGE				
3686.099 ± 0.004 ± 0.009		¹ ANASHIN 15	KEDR	$e^+e^- \rightarrow \text{hadrons}$
3686.12 ± 0.06 ± 0.10	4k	AAIJ 12H	LHCB	$pp \rightarrow J/\psi \pi^+ \pi^- X$
3685.95 ± 0.10	413	² ARTAMONOV 00	OLYA	$e^+e^- \rightarrow \text{hadrons}$
3685.98 ± 0.09 ± 0.04		³ ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3686.114 ± 0.007 ^{+0.011} _{-0.016}		⁴ ANASHIN 12	KEDR	$e^+e^- \rightarrow \text{hadrons}$
3686.111 ± 0.025 ± 0.009		AULCHENKO 03	KEDR	$e^+e^- \rightarrow \text{hadrons}$
3686.00 ± 0.10	413	⁵ ZHOLENTZ 80	OLYA	e^+e^-

¹ Supersedes AULCHENKO 03 and ANASHIN 12.

² Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).

³ Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $J/\psi(1S)$ mass from AULCHENKO 03.

⁴ From the scans in 2004 and 2006. ANASHIN 12 reports the value $3686.114 \pm 0.007 \pm 0.011^{+0.002}_{-0.012}$ MeV, where the third uncertainty is due to assumptions on the interference between the resonance and hadronic continuum. We combined the two systematic uncertainties.

⁵ Superseded by ARTAMONOV 00.

$m_{\psi(2S)} - m_{J/\psi(1S)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
589.188 ± 0.028 OUR AVERAGE			
589.194 ± 0.027 ± 0.011	¹ AULCHENKO 03	KEDR	$e^+e^- \rightarrow \text{hadrons}$
589.7 ± 1.2	LEMOIGNE 82	GOLI	$185 \pi^- \text{Be} \rightarrow \gamma \mu^+ \mu^- A$
589.07 ± 0.13	¹ ZHOLENTZ 80	OLYA	e^+e^-
588.7 ± 0.8	LUTH 75	MRK1	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
588 ± 1	² BAI 98E	BES	e^+e^-

¹ Redundant with data in mass above.

² Systematic errors not evaluated.

$\psi(2S)$ WIDTH

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
294 ± 8 OUR FIT				
286 ± 16 OUR AVERAGE				
358 ± 88 ± 4		ABLIKIM 08B	BES2	$e^+e^- \rightarrow \text{hadrons}$
290 ± 25 ± 4	2.7k	ANDREOTTI 07	E835	$\rho\bar{p} \rightarrow e^+e^-, J/\psi X$
331 ± 58 ± 2		ABLIKIM 06L	BES2	$e^+e^- \rightarrow \text{hadrons}$
264 ± 27		¹ BAI 02B	BES2	e^+e^-
287 ± 37 ± 16		² ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$

¹ From a simultaneous fit to the hadronic and $\mu^+\mu^-$ cross section, assuming $\Gamma = \Gamma_h + \Gamma_e + \Gamma_\mu + \Gamma_\tau$ and lepton universality. Does not include vacuum polarization correction.

² The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].

$\psi(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 hadrons	(97.85 ± 0.13) %	
Γ_2 virtual $\gamma \rightarrow$ hadrons	(1.73 ± 0.14) %	S=1.5
Γ_3 ggg	(10.6 ± 1.6) %	
Γ_4 γgg	(1.03 ± 0.29) %	
Γ_5 light hadrons	(15.4 ± 1.5) %	
Γ_6 e^+e^-	(7.93 ± 0.17) × 10 ⁻³	
Γ_7 $\mu^+\mu^-$	(8.0 ± 0.6) × 10 ⁻³	
Γ_8 $\tau^+\tau^-$	(3.1 ± 0.4) × 10 ⁻³	

Decays into $J/\psi(1S)$ and anything

Γ_9 $J/\psi(1S)$ anything	(61.4 ± 0.6) %
Γ_{10} $J/\psi(1S)$ neutrals	(25.37 ± 0.32) %
Γ_{11} $J/\psi(1S)\pi^+\pi^-$	(34.67 ± 0.30) %
Γ_{12} $J/\psi(1S)\pi^0\pi^0$	(18.23 ± 0.31) %
Γ_{13} $J/\psi(1S)\eta$	(3.37 ± 0.05) %
Γ_{14} $J/\psi(1S)\pi^0$	(1.268 ± 0.032) × 10 ⁻³

Hadronic decays

Γ_{15} $\pi^0 h_c(1P)$	(8.6 ± 1.3) × 10 ⁻⁴	
Γ_{16} $3(\pi^+\pi^-)\pi^0$	(3.5 ± 1.6) × 10 ⁻³	
Γ_{17} $2(\pi^+\pi^-)\pi^0$	(2.9 ± 1.0) × 10 ⁻³	S=4.7
Γ_{18} $\rho a_2(1320)$	(2.6 ± 0.9) × 10 ⁻⁴	
Γ_{19} $\rho\bar{p}$	(2.88 ± 0.10) × 10 ⁻⁴	
Γ_{20} $\Delta^{++}\bar{\Delta}^{--}$	(1.28 ± 0.35) × 10 ⁻⁴	
Γ_{21} $\Lambda\bar{\Lambda}\pi^0$	< 2.9 × 10 ⁻⁶	CL=90%
Γ_{22} $\Lambda\bar{\Lambda}\eta$	(2.5 ± 0.4) × 10 ⁻⁵	
Γ_{23} $\Lambda\bar{p}K^+$	(1.00 ± 0.14) × 10 ⁻⁴	
Γ_{24} $\Lambda\bar{p}K^+\pi^+\pi^-$	(1.8 ± 0.4) × 10 ⁻⁴	
Γ_{25} $\Lambda\bar{\Lambda}\pi^+\pi^-$	(2.8 ± 0.6) × 10 ⁻⁴	

Γ_{26}	$\Lambda\bar{\Lambda}$	$(3.81 \pm 0.13) \times 10^{-4}$	S=1.4
Γ_{27}	$\Lambda\bar{\Sigma}^+\pi^- + \text{c.c.}$	$(1.40 \pm 0.13) \times 10^{-4}$	
Γ_{28}	$\Lambda\bar{\Sigma}^-\pi^+ + \text{c.c.}$	$(1.54 \pm 0.14) \times 10^{-4}$	
Γ_{29}	$\Lambda\bar{\Sigma}^0$	$(1.23 \pm 0.24) \times 10^{-5}$	
Γ_{30}	$\Sigma^0\bar{p}K^+ + \text{c.c.}$	$(1.67 \pm 0.18) \times 10^{-5}$	
Γ_{31}	$\Sigma^+\bar{\Sigma}^-$	$(2.32 \pm 0.12) \times 10^{-4}$	
Γ_{32}	$\Sigma^0\bar{\Sigma}^0$	$(2.35 \pm 0.09) \times 10^{-4}$	S=1.1
Γ_{33}	$\Sigma(1385)^+\bar{\Sigma}(1385)^-$	$(8.5 \pm 0.7) \times 10^{-5}$	
Γ_{34}	$\Sigma(1385)^-\bar{\Sigma}(1385)^+$	$(8.5 \pm 0.8) \times 10^{-5}$	
Γ_{35}	$\Sigma(1385)^0\bar{\Sigma}(1385)^0$	$(6.9 \pm 0.7) \times 10^{-5}$	
Γ_{36}	$\Xi^-\bar{\Xi}^+$	$(2.87 \pm 0.11) \times 10^{-4}$	S=1.1
Γ_{37}	$\Xi^0\bar{\Xi}^0$	$(2.3 \pm 0.4) \times 10^{-4}$	S=4.2
Γ_{38}	$\Xi(1530)^0\bar{\Xi}(1530)^0$	$(5.2 \begin{smallmatrix} +3.2 \\ -1.2 \end{smallmatrix}) \times 10^{-5}$	
Γ_{39}	$K^-\Lambda\bar{\Xi}^+ + \text{c.c.}$	$(3.9 \pm 0.4) \times 10^{-5}$	
Γ_{40}	$\Xi(1690)^-\bar{\Xi}^+ \rightarrow K^-\Lambda\bar{\Xi}^+ +$	$(5.2 \pm 1.6) \times 10^{-6}$	
Γ_{41}	$\Xi(1820)^-\bar{\Xi}^+ \rightarrow K^-\Lambda\bar{\Xi}^+ +$	$(1.20 \pm 0.32) \times 10^{-5}$	
Γ_{42}	$K^-\Sigma^0\bar{\Xi}^+ + \text{c.c.}$	$(3.7 \pm 0.4) \times 10^{-5}$	
Γ_{43}	$\Omega^-\bar{\Omega}^+$	$(5.2 \pm 0.4) \times 10^{-5}$	
Γ_{44}	$\pi^0\rho\bar{p}$	$(1.53 \pm 0.07) \times 10^{-4}$	
Γ_{45}	$N(940)\bar{p} + \text{c.c.} \rightarrow \pi^0\rho\bar{p}$	$(6.4 \begin{smallmatrix} +1.8 \\ -1.3 \end{smallmatrix}) \times 10^{-5}$	
Γ_{46}	$N(1440)\bar{p} + \text{c.c.} \rightarrow \pi^0\rho\bar{p}$	$(7.3 \begin{smallmatrix} +1.7 \\ -1.5 \end{smallmatrix}) \times 10^{-5}$	S=2.5
Γ_{47}	$N(1520)\bar{p} + \text{c.c.} \rightarrow \pi^0\rho\bar{p}$	$(6.4 \begin{smallmatrix} +2.3 \\ -1.8 \end{smallmatrix}) \times 10^{-6}$	
Γ_{48}	$N(1535)\bar{p} + \text{c.c.} \rightarrow \pi^0\rho\bar{p}$	$(2.5 \pm 1.0) \times 10^{-5}$	
Γ_{49}	$N(1650)\bar{p} + \text{c.c.} \rightarrow \pi^0\rho\bar{p}$	$(3.8 \begin{smallmatrix} +1.4 \\ -1.7 \end{smallmatrix}) \times 10^{-5}$	
Γ_{50}	$N(1720)\bar{p} + \text{c.c.} \rightarrow \pi^0\rho\bar{p}$	$(1.79 \begin{smallmatrix} +0.26 \\ -0.70 \end{smallmatrix}) \times 10^{-5}$	
Γ_{51}	$N(2300)\bar{p} + \text{c.c.} \rightarrow \pi^0\rho\bar{p}$	$(2.6 \begin{smallmatrix} +1.2 \\ -0.7 \end{smallmatrix}) \times 10^{-5}$	
Γ_{52}	$N(2570)\bar{p} + \text{c.c.} \rightarrow \pi^0\rho\bar{p}$	$(2.13 \begin{smallmatrix} +0.40 \\ -0.31 \end{smallmatrix}) \times 10^{-5}$	
Γ_{53}	$\pi^0 f_0(2100) \rightarrow \pi^0\rho\bar{p}$	$(1.1 \pm 0.4) \times 10^{-5}$	
Γ_{54}	$\eta\rho\bar{p}$	$(6.0 \pm 0.4) \times 10^{-5}$	
Γ_{55}	$\eta f_0(2100) \rightarrow \eta\rho\bar{p}$	$(1.2 \pm 0.4) \times 10^{-5}$	
Γ_{56}	$N(1535)\bar{p} \rightarrow \eta\rho\bar{p}$	$(4.4 \pm 0.7) \times 10^{-5}$	
Γ_{57}	$\omega\rho\bar{p}$	$(6.9 \pm 2.1) \times 10^{-5}$	
Γ_{58}	$\phi\rho\bar{p}$	$< 2.4 \times 10^{-5}$	CL=90%
Γ_{59}	$\pi^+\pi^-\rho\bar{p}$	$(6.0 \pm 0.4) \times 10^{-4}$	
Γ_{60}	$\rho\bar{n}\pi^- \text{ or c.c.}$	$(2.48 \pm 0.17) \times 10^{-4}$	
Γ_{61}	$\rho\bar{n}\pi^-\pi^0$	$(3.2 \pm 0.7) \times 10^{-4}$	
Γ_{62}	$2(\pi^+\pi^-\pi^0)$	$(4.8 \pm 1.5) \times 10^{-3}$	
Γ_{63}	$\eta\pi^+\pi^-$	$< 1.6 \times 10^{-4}$	CL=90%

Γ_{64}	$\eta\pi^+\pi^-\pi^0$	$(9.5 \pm 1.7) \times 10^{-4}$	
Γ_{65}	$2(\pi^+\pi^-\eta)$	$(1.2 \pm 0.6) \times 10^{-3}$	
Γ_{66}	$\eta'\pi^+\pi^-\pi^0$	$(4.5 \pm 2.1) \times 10^{-4}$	
Γ_{67}	$\omega\pi^+\pi^-$	$(7.3 \pm 1.2) \times 10^{-4}$	S=2.1
Γ_{68}	$b_1^\pm\pi^\mp$	$(4.0 \pm 0.6) \times 10^{-4}$	S=1.1
Γ_{69}	$b_1^0\pi^0$	$(2.4 \pm 0.6) \times 10^{-4}$	
Γ_{70}	$\omega f_2(1270)$	$(2.2 \pm 0.4) \times 10^{-4}$	
Γ_{71}	$\pi^0\pi^0 K^+ K^-$	$(2.6 \pm 1.3) \times 10^{-4}$	
Γ_{72}	$\pi^+\pi^- K^+ K^-$	$(7.3 \pm 0.5) \times 10^{-4}$	
Γ_{73}	$\pi^0\pi^0 K_S^0 K_L^0$	$(1.3 \pm 0.5) \times 10^{-3}$	
Γ_{74}	$\rho^0 K^+ K^-$	$(2.2 \pm 0.4) \times 10^{-4}$	
Γ_{75}	$K^*(892)^0 \bar{K}_2^*(1430)^0$	$(1.9 \pm 0.5) \times 10^{-4}$	
Γ_{76}	$K^+ K^- \pi^+ \pi^- \eta$	$(1.3 \pm 0.7) \times 10^{-3}$	
Γ_{77}	$K^+ K^- 2(\pi^+ \pi^-) \pi^0$	$(1.00 \pm 0.31) \times 10^{-3}$	
Γ_{78}	$K^+ K^- 2(\pi^+ \pi^-)$	$(1.9 \pm 0.9) \times 10^{-3}$	
Γ_{79}	$K_1(1270)^\pm K^\mp$	$(1.00 \pm 0.28) \times 10^{-3}$	
Γ_{80}	$K_S^0 K_S^0 \pi^+ \pi^-$	$(2.2 \pm 0.4) \times 10^{-4}$	
Γ_{81}	$\rho^0 \rho \bar{p}$	$(5.0 \pm 2.2) \times 10^{-5}$	
Γ_{82}	$K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$	$(6.7 \pm 2.5) \times 10^{-4}$	
Γ_{83}	$2(\pi^+ \pi^-)$	$(2.4 \pm 0.6) \times 10^{-4}$	S=2.2
Γ_{84}	$\rho^0 \pi^+ \pi^-$	$(2.2 \pm 0.6) \times 10^{-4}$	S=1.4
Γ_{85}	$K^+ K^- \pi^+ \pi^- \pi^0$	$(1.26 \pm 0.09) \times 10^{-3}$	
Γ_{86}	$\omega f_0(1710) \rightarrow \omega K^+ K^-$	$(5.9 \pm 2.2) \times 10^{-5}$	
Γ_{87}	$K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.}$	$(8.6 \pm 2.2) \times 10^{-4}$	
Γ_{88}	$K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.}$	$(9.6 \pm 2.8) \times 10^{-4}$	
Γ_{89}	$K^*(892)^+ K^- \rho^0 + \text{c.c.}$	$(7.3 \pm 2.6) \times 10^{-4}$	
Γ_{90}	$K^*(892)^0 K^- \rho^+ + \text{c.c.}$	$(6.1 \pm 1.8) \times 10^{-4}$	
Γ_{91}	$\eta K^+ K^-$, no $\eta\phi$	$(3.1 \pm 0.4) \times 10^{-5}$	
Γ_{92}	$\omega K^+ K^-$	$(1.62 \pm 0.11) \times 10^{-4}$	S=1.1
Γ_{93}	$\omega K^*(892)^+ K^- + \text{c.c.}$	$(2.07 \pm 0.26) \times 10^{-4}$	
Γ_{94}	$\omega K_2^*(1430)^+ K^- + \text{c.c.}$	$(6.1 \pm 1.2) \times 10^{-5}$	
Γ_{95}	$\omega \bar{K}^*(892)^0 K^0$	$(1.68 \pm 0.30) \times 10^{-4}$	
Γ_{96}	$\omega \bar{K}_2^*(1430)^0 K^0$	$(5.8 \pm 2.2) \times 10^{-5}$	
Γ_{97}	$\omega X(1440) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.}$	$(1.6 \pm 0.4) \times 10^{-5}$	
Γ_{98}	$\omega X(1440) \rightarrow \omega K^+ K^- \pi^0$	$(1.09 \pm 0.26) \times 10^{-5}$	
Γ_{99}	$\omega f_1(1285) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.}$	$(3.0 \pm 1.0) \times 10^{-6}$	
Γ_{100}	$\omega f_1(1285) \rightarrow \omega K^+ K^- \pi^0$	$(1.2 \pm 0.7) \times 10^{-6}$	
Γ_{101}	$3(\pi^+ \pi^-)$	$(3.5 \pm 2.0) \times 10^{-4}$	S=2.8
Γ_{102}	$\rho \bar{p} \pi^+ \pi^- \pi^0$	$(7.3 \pm 0.7) \times 10^{-4}$	
Γ_{103}	$K^+ K^-$	$(7.5 \pm 0.5) \times 10^{-5}$	
Γ_{104}	$K_S^0 K_L^0$	$(5.34 \pm 0.33) \times 10^{-5}$	
Γ_{105}	$\pi^+ \pi^- \pi^0$	$(2.01 \pm 0.17) \times 10^{-4}$	S=1.7

Γ_{106}	$\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0$	$(1.9^{+1.2}_{-0.4}) \times 10^{-4}$	
Γ_{107}	$\rho(770)\pi \rightarrow \pi^+\pi^-\pi^0$	$(3.2 \pm 1.2) \times 10^{-5}$	S=1.8
Γ_{108}	$\pi^+\pi^-$	$(7.8 \pm 2.6) \times 10^{-6}$	
Γ_{109}	$K_1(1400)^\pm K^\mp$	$< 3.1 \times 10^{-4}$	CL=90%
Γ_{110}	$K_2^*(1430)^\pm K^\mp$	$(7.1^{+1.3}_{-0.9}) \times 10^{-5}$	
Γ_{111}	$K^+K^-\pi^0$	$(4.07 \pm 0.31) \times 10^{-5}$	
Γ_{112}	$K_S^0 K_L^0 \pi^0$	$< 3.0 \times 10^{-4}$	CL=90%
Γ_{113}	$K_S^0 K_L^0 \eta$	$(1.3 \pm 0.5) \times 10^{-3}$	
Γ_{114}	$K^+K^*(892)^- + \text{c.c.}$	$(2.9 \pm 0.4) \times 10^{-5}$	S=1.2
Γ_{115}	$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	$(1.09 \pm 0.20) \times 10^{-4}$	
Γ_{116}	$\phi\pi^+\pi^-$	$(1.18 \pm 0.26) \times 10^{-4}$	S=1.5
Γ_{117}	$\phi f_0(980) \rightarrow \pi^+\pi^-$	$(7.5 \pm 3.3) \times 10^{-5}$	S=1.6
Γ_{118}	$2(K^+K^-)$	$(6.3 \pm 1.3) \times 10^{-5}$	
Γ_{119}	ϕK^+K^-	$(7.0 \pm 1.6) \times 10^{-5}$	
Γ_{120}	$2(K^+K^-)\pi^0$	$(1.10 \pm 0.28) \times 10^{-4}$	
Γ_{121}	$\phi\eta$	$(3.10 \pm 0.31) \times 10^{-5}$	
Γ_{122}	$\phi\eta'$	$(3.1 \pm 1.6) \times 10^{-5}$	
Γ_{123}	$\omega\eta'$	$(3.2^{+2.5}_{-2.1}) \times 10^{-5}$	
Γ_{124}	$\omega\pi^0$	$(2.1 \pm 0.6) \times 10^{-5}$	
Γ_{125}	$\rho\eta'$	$(1.9^{+1.7}_{-1.2}) \times 10^{-5}$	
Γ_{126}	$\rho\eta$	$(2.2 \pm 0.6) \times 10^{-5}$	S=1.1
Γ_{127}	$\omega\eta$	$< 1.1 \times 10^{-5}$	CL=90%
Γ_{128}	$\phi\pi^0$	$< 4 \times 10^{-7}$	CL=90%
Γ_{129}	$\eta_c\pi^+\pi^-\pi^0$	$< 1.0 \times 10^{-3}$	CL=90%
Γ_{130}	$\bar{p}\bar{p}K^+K^-$	$(2.7 \pm 0.7) \times 10^{-5}$	
Γ_{131}	$\bar{\Lambda}nK_S^0 + \text{c.c.}$	$(8.1 \pm 1.8) \times 10^{-5}$	
Γ_{132}	$\phi f_2'(1525)$	$(4.4 \pm 1.6) \times 10^{-5}$	
Γ_{133}	$\Theta(1540)\bar{\Theta}(1540) \rightarrow$ $K_S^0 p K^- \bar{n} + \text{c.c.}$	$< 8.8 \times 10^{-6}$	CL=90%
Γ_{134}	$\Theta(1540)K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	$< 1.0 \times 10^{-5}$	CL=90%
Γ_{135}	$\Theta(1540)K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	$< 7.0 \times 10^{-6}$	CL=90%
Γ_{136}	$\bar{\Theta}(1540)K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	$< 2.6 \times 10^{-5}$	CL=90%
Γ_{137}	$\bar{\Theta}(1540)K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	$< 6.0 \times 10^{-6}$	CL=90%
Γ_{138}	$K_S^0 K_S^0$	$< 4.6 \times 10^{-6}$	

Radiative decays

Γ_{139}	$\gamma\chi_{c0}(1P)$	$(9.79 \pm 0.20) \%$	
Γ_{140}	$\gamma\chi_{c1}(1P)$	$(9.75 \pm 0.24) \%$	
Γ_{141}	$\gamma\chi_{c2}(1P)$	$(9.52 \pm 0.20) \%$	
Γ_{142}	$\gamma\eta_c(1S)$	$(3.4 \pm 0.5) \times 10^{-3}$	S=1.3
Γ_{143}	$\gamma\eta_c(2S)$	$(7 \pm 5) \times 10^{-4}$	

Γ_{144}	$\gamma\pi^0$	$(1.04 \pm 0.22) \times 10^{-6}$	S=1.4
Γ_{145}	$\gamma\eta'(958)$	$(1.24 \pm 0.04) \times 10^{-4}$	
Γ_{146}	$\gamma f_2(1270)$	$(2.73^{+0.29}_{-0.25}) \times 10^{-4}$	S=1.8
Γ_{147}	$\gamma f_0(1370) \rightarrow \gamma K \bar{K}$	$(3.1 \pm 1.7) \times 10^{-5}$	
Γ_{148}	$\gamma f_0(1500)$	$(9.2 \pm 1.9) \times 10^{-5}$	
Γ_{149}	$\gamma f_2'(1525)$	$(3.3 \pm 0.8) \times 10^{-5}$	
Γ_{150}	$\gamma f_0(1710)$		
Γ_{151}	$\gamma f_0(1710) \rightarrow \gamma\pi\pi$	$(3.5 \pm 0.6) \times 10^{-5}$	
Γ_{152}	$\gamma f_0(1710) \rightarrow \gamma K \bar{K}$	$(6.6 \pm 0.7) \times 10^{-5}$	
Γ_{153}	$\gamma f_0(2100) \rightarrow \gamma\pi\pi$	$(4.8 \pm 1.0) \times 10^{-6}$	
Γ_{154}	$\gamma f_0(2200) \rightarrow \gamma K \bar{K}$	$(3.2 \pm 1.0) \times 10^{-6}$	
Γ_{155}	$\gamma f_J(2220) \rightarrow \gamma\pi\pi$	$< 5.8 \times 10^{-6}$	CL=90%
Γ_{156}	$\gamma f_J(2220) \rightarrow \gamma K \bar{K}$	$< 9.5 \times 10^{-6}$	CL=90%
Γ_{157}	$\gamma\gamma$	$< 1.5 \times 10^{-4}$	CL=90%
Γ_{158}	$\gamma\eta$	$(9.2 \pm 1.8) \times 10^{-7}$	
Γ_{159}	$\gamma\eta\pi^+\pi^-$	$(8.7 \pm 2.1) \times 10^{-4}$	
Γ_{160}	$\gamma\eta(1405)$		
Γ_{161}	$\gamma\eta(1405) \rightarrow \gamma K \bar{K}\pi$	$< 9 \times 10^{-5}$	CL=90%
Γ_{162}	$\gamma\eta(1405) \rightarrow \eta\pi^+\pi^-$	$(3.6 \pm 2.5) \times 10^{-5}$	
Γ_{163}	$\gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^+\pi^-\pi^0$	$< 5.0 \times 10^{-7}$	CL=90%
Γ_{164}	$\gamma\eta(1475)$		
Γ_{165}	$\gamma\eta(1475) \rightarrow K \bar{K}\pi$	$< 1.4 \times 10^{-4}$	CL=90%
Γ_{166}	$\gamma\eta(1475) \rightarrow \eta\pi^+\pi^-$	$< 8.8 \times 10^{-5}$	CL=90%
Γ_{167}	$\gamma 2(\pi^+\pi^-)$	$(4.0 \pm 0.6) \times 10^{-4}$	
Γ_{168}	$\gamma K^{*0} K^+ \pi^- + \text{c.c.}$	$(3.7 \pm 0.9) \times 10^{-4}$	
Γ_{169}	$\gamma K^{*0} \bar{K}^{*0}$	$(2.4 \pm 0.7) \times 10^{-4}$	
Γ_{170}	$\gamma K_S^0 K^+ \pi^- + \text{c.c.}$	$(2.6 \pm 0.5) \times 10^{-4}$	
Γ_{171}	$\gamma K^+ K^- \pi^+ \pi^-$	$(1.9 \pm 0.5) \times 10^{-4}$	
Γ_{172}	$\gamma p \bar{p}$	$(3.9 \pm 0.5) \times 10^{-5}$	S=2.0
Γ_{173}	$\gamma f_2(1950) \rightarrow \gamma p \bar{p}$	$(1.20 \pm 0.22) \times 10^{-5}$	
Γ_{174}	$\gamma f_2(2150) \rightarrow \gamma p \bar{p}$	$(7.2 \pm 1.8) \times 10^{-6}$	
Γ_{175}	$\gamma X(1835) \rightarrow \gamma p \bar{p}$	$(4.6^{+1.8}_{-4.0}) \times 10^{-6}$	
Γ_{176}	$\gamma X \rightarrow \gamma p \bar{p}$	[a] $< 2 \times 10^{-6}$	CL=90%
Γ_{177}	$\gamma\pi^+\pi^- p \bar{p}$	$(2.8 \pm 1.4) \times 10^{-5}$	
Γ_{178}	$\gamma 2(\pi^+\pi^-) K^+ K^-$	$< 2.2 \times 10^{-4}$	CL=90%
Γ_{179}	$\gamma 3(\pi^+\pi^-)$	$< 1.7 \times 10^{-4}$	CL=90%
Γ_{180}	$\gamma K^+ K^- K^+ K^-$	$< 4 \times 10^{-5}$	CL=90%
Γ_{181}	$\gamma\gamma J/\psi$	$(3.1^{+1.0}_{-1.2}) \times 10^{-4}$	
Γ_{182}	$e^+ e^- \chi_{c0}(1P)$	$(1.06 \pm 0.24) \times 10^{-3}$	
Γ_{183}	$e^+ e^- \chi_{c1}(1P)$	$(8.5 \pm 0.6) \times 10^{-4}$	
Γ_{184}	$e^+ e^- \chi_{c2}(1P)$	$(7.0 \pm 0.8) \times 10^{-4}$	

Weak decays

$\Gamma_{185} D^0 e^+ e^- + \text{c.c.} < 1.4 \times 10^{-7} \text{ CL}=90\%$

Other decays

$\Gamma_{186} \text{ invisible} < 1.6 \% \text{ CL}=90\%$

[a] For a narrow resonance in the range $2.2 < M(X) < 2.8 \text{ GeV}$.

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 84 branching ratios uses 247 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 376.9$ for 198 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x_7	3									
x_8	1	0								
x_{11}	29	11	2							
x_{12}	28	6	1	48						
x_{13}	13	4	1	36	15					
x_{19}	0	1	0	5	3	2				
x_{139}	1	0	0	2	1	1	0			
x_{140}	1	0	0	2	1	1	0	0		
x_{141}	1	0	0	3	1	1	0	0	0	
Γ	-81	-4	-1	-38	-34	-16	-8	-1	-1	-1
	x_6	x_7	x_8	x_{11}	x_{12}	x_{13}	x_{19}	x_{139}	x_{140}	x_{141}

$\psi(2S)$ PARTIAL WIDTHS

$\Gamma(\text{hadrons})$

Γ_1

VALUE (keV) DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

258 ± 26	BAI	02B	BES2	$e^+ e^-$
224 ± 56	LUTH	75	MRK1	$e^+ e^-$

$\Gamma(e^+ e^-)$

Γ_6

VALUE (keV) DOCUMENT ID TECN COMMENT

2.33 ± 0.04 OUR FIT

2.29 ± 0.06 OUR AVERAGE

$2.23 \pm 0.10 \pm 0.02$	¹ ABLIKIM	15V	BES3	$4.0\text{--}4.4 e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$
$2.338 \pm 0.037 \pm 0.096$	ABLIKIM	08B	BES2	$e^+ e^- \rightarrow \text{hadrons}$

2.330 ± 0.036 ± 0.110	ABLIKIM	06L	BES2	$e^+e^- \rightarrow$ hadrons
2.44 ± 0.21	² BAI	02B	BES2	e^+e^-
2.14 ± 0.21	ALEXANDER	89	RVUE	See Υ mini-review
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.0 ± 0.3	BRANDELIK	79C	DASP	e^+e^-
2.1 ± 0.3	³ LUTH	75	MRK1	e^+e^-

¹ABLIKIM 15V reports $2.213 \pm 0.018 \pm 0.099$ keV from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+e^-)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.95 \pm 0.45) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.67 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channel, assuming $\Gamma_e = \Gamma_\mu = \Gamma_\tau/0.38847$.

³From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.

$\Gamma(\gamma\gamma)$ Γ_{157}

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<43	90	BRANDELIK	79C	DASP e^+e^-

$\psi(2S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel(i) in the e^+e^- annihilation. We list only data that have not been used to determine the partial width $\Gamma(i)$ or the branching ratio $\Gamma(i)/\text{total}$.

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_6/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
2.233 ± 0.015 ± 0.042	¹ ANASHIN	12	KEDR $e^+e^- \rightarrow$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.2 ± 0.4	ABRAMS	75	MRK1	e^+e^-
-----------	--------	----	------	----------

¹ANASHIN 12 reports the value $2.233 \pm 0.015 \pm 0.037 \pm 0.020$ keV, where the third uncertainty is due to assumptions on the interference between the resonance and hadronic continuum. We combined the two systematic uncertainties.

$\Gamma(\tau^+\tau^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_8\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.0 ± 2.6	79	¹ ANASHIN	07	KEDR $e^+e^- \rightarrow \psi(2S) \rightarrow \tau^+\tau^-$
-----------	----	----------------------	----	---

¹Using $\psi(2S)$ total width of 337 ± 13 keV. Systematic errors not evaluated.

$\Gamma(J/\psi(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{11}\Gamma_6/\Gamma$


VALUE (keV) EVTS DOCUMENT ID TECN COMMENT

0.809±0.013 OUR FIT

0.837±0.025 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

0.837±0.028±0.005		¹ LEES	12E	BABR	10.6	$e^+e^- \rightarrow 2\pi^+2\pi^-\gamma$
0.852±0.010±0.026	19.5k	ADAM	06	CLEO	3.773	$e^+e^- \rightarrow \gamma\psi(2S)$
0.68 ±0.09		² BAI	98E	BES		e^+e^-
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
0.88 ±0.08 ±0.03	256	³ AUBERT	07AU	BABR	10.6	$e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$
0.755±0.048±0.004	544	⁴ AUBERT	05D	BABR	10.6	$e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-\gamma$

WEIGHTED AVERAGE
0.837±0.025 (Error scaled by 1.3)



$\Gamma(J/\psi(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ (keV)

- ¹ LEES 12E reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = (49.9 \pm 1.3 \pm 1.0) \times 10^{-3}$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ² The value of $\Gamma(e^+e^-)$ quoted in BAI 98E is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6) \times 10^{-2}$ and $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1203 \pm 0.0038$. Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.
- ³ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)] = 0.0186 \pm 0.0012 \pm 0.0011$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0) = (2.11 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ⁴ AUBERT 05D reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = 0.0450 \pm 0.0018 \pm 0.0022$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their

experiment's error and our second error is the systematic error from using our best value. Superseded by LEES 12E.

$\Gamma(J/\psi(1S)\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{12}\Gamma_6/\Gamma$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.425±0.009 OUR FIT				
0.411±0.008±0.018	3.6k±96	ADAM	06	CLEO 3.773 e ⁺ e ⁻ → γψ(2S)

$\Gamma(J/\psi(1S)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{13}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
78.7± 1.6 OUR FIT				
87 ± 9 OUR AVERAGE				
83 ±25 ±5	14	¹ AUBERT	07AU	BABR 10.6 e ⁺ e ⁻ → J/ψπ ⁺ π ⁻ π ⁰ γ
88 ± 6 ±7	291 ± 24	ADAM	06	CLEO 3.773 e ⁺ e ⁻ → γψ(2S)
¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow J/\psi\eta) \cdot B(J/\psi \rightarrow \mu^+\mu^-) \cdot B(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.11 \pm 0.33 \pm 0.07$ eV.				

$\Gamma(J/\psi(1S)\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{14}\Gamma_6/\Gamma$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<8	90	<37	ADAM	06	CLEO 3.773 e ⁺ e ⁻ → γψ(2S)

$\Gamma(p\bar{p}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{19}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.672±0.023 OUR FIT				
0.63 ±0.05 OUR AVERAGE				Error includes scale factor of 1.2.
0.67 ±0.12 ±0.02	43	¹ LEES	130	BABR e ⁺ e ⁻ → p \bar{p} γ
0.74 ±0.07 ±0.04	142	² LEES	13Y	BABR e ⁺ e ⁻ → p \bar{p} γ
0.579±0.038±0.036	2.7k	ANDREOTTI	07	E835 p \bar{p} → e ⁺ e ⁻ , J/ψX
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.70 ±0.17 ±0.03	22	³ AUBERT	06B	BABR e ⁺ e ⁻ → p \bar{p} γ
¹ ISR photon reconstructed in the detector				
² ISR photon undetected				
³ Superseded by LEES 130				

$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{26}\Gamma_6/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
1.5±0.4±0.1	AUBERT	07BD	BABR 10.6 e ⁺ e ⁻ → Λ $\bar{\Lambda}$ γ

$\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{62}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.2±3.3±1.3	43	AUBERT	06D	BABR 10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻ π ⁰)γ

$\Gamma(\pi^0\pi^0K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{71}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.60±0.31±0.03	17	LEES	12F	BABR 10.6 e ⁺ e ⁻ → π ⁰ π ⁰ K ⁺ K ⁻ γ

$\Gamma(K^+ K^- 2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{78}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.4±2.1±0.3	26	AUBERT	06D BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ 2(π ⁺ π ⁻)γ

$\Gamma(\pi^+ \pi^- K^+ K^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{72}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.92±0.30±0.06	133	LEES	12F BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.56±0.42±0.16	85	¹ AUBERT	07AK BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ
----------------	----	---------------------	-----------	--

¹ Superseded by LEES 12F.

$\Gamma(\pi^0 \pi^0 K_S^0 K_L^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{73}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.92±1.27±0.15	14	LEES	17A BABR	e ⁺ e ⁻ → K _S ⁰ K _L ⁰ π ⁰ π ⁰ γ

$\Gamma(K_S^0 K_L^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{112}\Gamma_6/\Gamma$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.7	90	8	LEES	17A BABR	e ⁺ e ⁻ → K _S ⁰ K _L ⁰ π ⁰ γ

$\Gamma(K_S^0 K_L^0 \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{113}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.14±1.08±0.16	16	LEES	17A BABR	e ⁺ e ⁻ → K _S ⁰ K _L ⁰ ηγ

$\Gamma(\phi f_0(980) \rightarrow \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{117}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.345±0.128±0.004	12	¹ LEES	12F BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.345±0.168±0.004	6 ± 3	² AUBERT	07AK BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ
-------------------	-------	---------------------	-----------	--

¹ LEES 12F reports [$\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}$] × [B(φ(1020) → K⁺K⁻)] = 0.17 ± 0.06 ± 0.02 eV which we divide by our best value B(φ(1020) → K⁺K⁻) = (49.2 ± 0.5) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Superseded by LEES 12F. AUBERT 07AK reports [$\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}$] × [B(φ(1020) → K⁺K⁻)] = 0.17 ± 0.08 ± 0.02 eV which we divide by our best value B(φ(1020) → K⁺K⁻) = (49.2 ± 0.5) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(K^+ K^-)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{118}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.22±0.10±0.02	13	LEES	12F BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ K ⁺ K ⁻ γ

$$\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{116}\Gamma_6/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

0.55±0.19±0.01	19	¹ LEES	12F BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ γ
-----------------------	----	-------------------	----------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.57±0.23±0.01	10	² AUBERT,BE	06D BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ γ
----------------	----	------------------------	----------	--

¹ LEES 12F reports [$\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B($\phi(1020) \rightarrow K^+K^-$)] = 0.27 ± 0.09 ± 0.02 eV which we divide by our best value B($\phi(1020) \rightarrow K^+K^-$) = (49.2 ± 0.5) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Superseded by LEES 12F. AUBERT,BE 06D reports [$\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B($\phi(1020) \rightarrow K^+K^-$)] = 0.28 ± 0.11 ± 0.02 eV which we divide by our best value B($\phi(1020) \rightarrow K^+K^-$) = (49.2 ± 0.5) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(2(\pi^+\pi^-)\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{17}\Gamma_6/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

29.7±2.2±1.8	410	AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)π ⁰ γ
---------------------	-----	--------	-----------	--

$$\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{67}\Gamma_6/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

3.01±0.84±0.02	37	¹ AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → ωπ ⁺ π ⁻ γ
-----------------------	----	---------------------	-----------	---

¹ AUBERT 07AU reports [$\Gamma(\psi(2S) \rightarrow \omega\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B($\omega(782) \rightarrow \pi^+\pi^-\pi^0$)] = 2.69 ± 0.73 ± 0.16 eV which we divide by our best value B($\omega(782) \rightarrow \pi^+\pi^-\pi^0$) = (89.2 ± 0.7) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(2(\pi^+\pi^-\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{65}\Gamma_6/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

2.87±1.41±0.01	16	¹ AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)ηγ
-----------------------	----	---------------------	-----------	--

¹ AUBERT 07AU reports [$\Gamma(\psi(2S) \rightarrow 2(\pi^+\pi^-\eta) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B($\eta \rightarrow 2\gamma$)] = 1.13 ± 0.55 ± 0.08 eV which we divide by our best value B($\eta \rightarrow 2\gamma$) = (39.41 ± 0.20) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{85}\Gamma_6/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

4.4±1.3±0.3	32	AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ π ⁰ γ
--------------------	----	--------	-----------	---

$$\Gamma(K^+K^-\pi^+\pi^-\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{76}\Gamma_6/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

3.04±1.79±0.02	7	¹ AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ ηγ
-----------------------	---	---------------------	-----------	---

¹ AUBERT 07AU reports [$\Gamma(\psi(2S) \rightarrow K^+K^-\pi^+\pi^-\eta) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B($\eta \rightarrow 2\gamma$)] = 1.2 ± 0.7 ± 0.1 eV which we divide by our best value B($\eta \rightarrow 2\gamma$) = (39.41 ± 0.20) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{103}\Gamma_6/\Gamma$

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-------------------	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.147 \pm 0.035 \pm 0.005$	66	¹ LEES	15J BABR	$e^+e^- \rightarrow K^+K^-\gamma$
$0.197 \pm 0.035 \pm 0.005$	66	² LEES	15J BABR	$e^+e^- \rightarrow K^+K^-\gamma$
$0.35 \pm 0.14 \pm 0.03$	11	³ LEES	13Q BABR	$e^+e^- \rightarrow K^+K^-\gamma$

¹ $\sin\phi > 0$.

² $\sin\phi < 0$.

³ Interference with non-resonant K^+K^- production not taken into account.

 $\psi(2S)$ BRANCHING RATIOS $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$ Γ_1/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

0.9785 ± 0.0013 OUR AVERAGE

0.9779 ± 0.0015	¹ BAI	02B BES2	e^+e^-
0.981 ± 0.003	¹ LUTH	75 MRK1	e^+e^-

¹ Includes cascade decay into $J/\psi(1S)$.

 $\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}}$ Γ_2/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

0.0173 ± 0.0014 OUR AVERAGE Error includes scale factor of 1.5.

0.0166 ± 0.0010	^{1,2} SETH	04 RVUE	e^+e^-
0.0199 ± 0.0019	¹ BAI	02B BES2	e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.029 ± 0.004	¹ LUTH	75 MRK1	e^+e^-
-------------------	-------------------	---------	----------

¹ Included in $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$.

² Using $B(\psi(2S) \rightarrow \ell^+\ell^-) = (0.73 \pm 0.04)\%$ from RPP-2002 and $R = 2.28 \pm 0.04$ determined by a fit to data from BAI 00 and BAI 02C.

 $\Gamma(g g g)/\Gamma_{\text{total}}$ Γ_3/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

10.58 ± 1.62 2.9 M ¹ LIBBY 09 CLEO $\psi(2S) \rightarrow \text{hadrons}$

¹ Calculated using $\Gamma(\gamma g g)/\Gamma(g g g) = 0.097 \pm 0.026 \pm 0.016$ from LIBBY 09, $B(\psi(2S) \rightarrow XJ/\psi)$ relative and absolute branching fractions from MENDEZ 08, $B(\psi(2S) \rightarrow \gamma\eta_c)$ from MITCHELL 09, and $B(\psi(2S) \rightarrow \text{virtual } \gamma \rightarrow \text{hadrons})$, $B(\psi(2S) \rightarrow \gamma\chi_{cJ})$, and $B(\psi(2S) \rightarrow \ell^+\ell^-)$ from PDG 08. The statistical error is negligible and the systematic error is largely uncorrelated with that of $\Gamma(\gamma g g)/\Gamma_{\text{total}}$ LIBBY 09 measurement.

 $\Gamma(\gamma g g)/\Gamma_{\text{total}}$ Γ_4/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

1.025 ± 0.288 200 k ¹ LIBBY 09 CLEO $\psi(2S) \rightarrow \gamma + \text{hadrons}$

¹ Calculated using $\Gamma(\gamma g g)/\Gamma(g g g) = 0.097 \pm 0.026 \pm 0.016$ from LIBBY 09. The statistical error is negligible and the systematic error is largely uncorrelated with that of $\Gamma(g g g)/\Gamma_{\text{total}}$ LIBBY 09 measurement.

$\Gamma(\gamma g g)/\Gamma(g g g)$ Γ_4/Γ_3

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$9.7 \pm 2.6 \pm 1.6$	2.9 M	LIBBY	09	CLEO $\psi(2S) \rightarrow (\gamma +)$ hadrons

 $\Gamma(\text{light hadrons})/\Gamma_{\text{total}}$ Γ_5/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.154 ± 0.015	¹ MENDEZ	08	CLEO $e^+e^- \rightarrow \psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.169 ± 0.026	² ADAM	05A	CLEO $e^+e^- \rightarrow \psi(2S)$
¹ Uses $B(\psi(2S) \rightarrow J/\psi X)$ from MENDEZ 08 and other branching fractions from PDG 07.			
² Uses $B(J/\psi X)$ from ADAM 05A, $B(\chi_{cJ}\gamma)$, $B(\eta_c\gamma)$ from ATHAR 04 and $B(\ell^+\ell^-)$ from PDG 04. Superseded by MENDEZ 08.			

 $\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_6/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
79.3 ± 1.7 OUR FIT			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
88 ± 13	¹ FELDMAN	77	RVUE e^+e^-
¹ From an overall fit assuming equal partial widths for e^+e^- and $\mu^+\mu^-$. For a measurement of the ratio see the entry $\Gamma(\mu^+\mu^-)/\Gamma(e^+e^-)$ below. Includes LUTH 75, HILGER 75, BURMESTER 77.			

 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_7/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>
80 ± 6 OUR FIT	

 $\Gamma(\mu^+\mu^-)/\Gamma(e^+e^-)$ Γ_7/Γ_6

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.00 ± 0.08 OUR FIT			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.89 ± 0.16	BOYARSKI	75C	MRK1 e^+e^-

 $\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
31 ± 4 OUR FIT			
$30.8 \pm 2.1 \pm 3.8$	¹ ABLIKIM	06W	BES $e^+e^- \rightarrow \psi(2S)$
¹ Computed using PDG 02 value of $B(\psi(2S) \rightarrow \text{hadrons}) = 0.9810 \pm 0.0030$ to estimate the total number of $\psi(2S)$ events.			

————— **DECAYS INTO $J/\psi(1S)$ AND ANYTHING** —————

 $\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$ Γ_9/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.614 ± 0.006 OUR FIT				
0.55 ± 0.07 OUR AVERAGE				
0.51 ± 0.12		BRANDELIK	79C	DASP $e^+e^- \rightarrow \mu^+\mu^-X$
0.57 ± 0.08		ABRAMS	75B	MRK1 $e^+e^- \rightarrow \mu^+\mu^-X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.6254 \pm 0.0016 \pm 0.0155$	1.1M	¹ MENDEZ	08	CLEO $\psi(2S) \rightarrow \ell^+\ell^-X$
$0.5950 \pm 0.0015 \pm 0.0190$	151k	ADAM	05A	CLEO Repl. by MENDEZ 08
¹ Not independent from other measurements of MENDEZ 08.				

$\Gamma(e^+ e^-)/\Gamma(J/\psi(1S)\text{anything})$

$$\Gamma_6/\Gamma_9 = \Gamma_6/(\Gamma_{11}+\Gamma_{12}+\Gamma_{13}+0.343\Gamma_{140}+0.190\Gamma_{141})$$


VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.292 ± 0.026				OUR FIT

1.28 ± 0.04 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.

1.22 ± 0.02 ± 0.05	5097 ± 73	¹ ANDREOTTI 05	E835	$\rho\bar{p} \rightarrow \psi(2S) \rightarrow e^+e^-$
1.28 ± 0.03 ± 0.02		¹ AMBROGIANI 00A	E835	$\rho\bar{p} \rightarrow \psi(2S)$
1.44 ± 0.08 ± 0.02		¹ ARMSTRONG 97	E760	$\bar{p}p \rightarrow \psi(2S)$

¹ Using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.

WEIGHTED AVERAGE
1.28 ± 0.04 (Error scaled by 1.6)



$\Gamma(\mu^+ \mu^-)/\Gamma(J/\psi(1S)\text{anything})$

$$\Gamma_7/\Gamma_9 = \Gamma_7/(\Gamma_{11}+\Gamma_{12}+\Gamma_{13}+0.343\Gamma_{140}+0.190\Gamma_{141})$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0130 ± 0.0010			OUR FIT

0.014 ± 0.003 HILGER 75 SPEC e^+e^-

$\Gamma(J/\psi(1S)\text{neutrals})/\Gamma_{\text{total}}$

Γ_{10}/Γ

VALUE	DOCUMENT ID
0.2537 ± 0.0032	OUR FIT

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{11}/Γ

VALUE EVTS DOCUMENT ID TECN COMMENT

0.3467 ± 0.0030 OUR FIT

0.348 ± 0.005 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.


0.3498 ± 0.0002 ± 0.0045	20M	ABLIKIM	13R	BES3	$\psi(2S) \rightarrow J/\psi\pi^+\pi^-$
0.3504 ± 0.0007 ± 0.0077	565k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+\ell^-\pi^+\pi^-$
0.323 ± 0.014		BAI	02B	BES2	e^+e^-
0.32 ± 0.04		ABRAMS	75B	MRK1	$e^+e^- \rightarrow J/\psi\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.3354 ± 0.0014 ± 0.0110	60k	¹ ADAM	05A	CLEO	Repl. by MENDEZ 08
--------------------------	-----	-------------------	-----	------	--------------------

¹ Not independent from other values reported by ADAM 05A.

WEIGHTED AVERAGE
0.348 ± 0.005 (Error scaled by 1.3)



$\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

Γ_6/Γ_{11}

VALUE DOCUMENT ID TECN COMMENT

0.0229 ± 0.0005 OUR FIT

0.0252 ± 0.0028 ± 0.0011	¹ AUBERT	02B	BABR	e^+e^-
---------------------------------	---------------------	-----	------	----------

¹ Using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.

$\Gamma(\mu^+\mu^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

Γ_7/Γ_{11}

VALUE DOCUMENT ID TECN COMMENT

0.0230 ± 0.0017 OUR FIT

0.0228 ± 0.0018 OUR AVERAGE

0.0230 ± 0.0020 ± 0.0012	¹ AAIJ	16Y	LHCB	$\Lambda_b^0 \rightarrow \psi(2S)X$
0.0216 ± 0.0026 ± 0.0014	² AUBERT	02B	BABR	e^+e^-
0.0327 ± 0.0077 ± 0.0072	² GRIBUSHIN	96	FMPS	515 $\pi^- \text{Be} \rightarrow 2\mu X$

¹ Using $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033) \times 10^{-2}$.

² Using $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = (5.88 \pm 0.10) \times 10^{-2}$.

$\Gamma(\tau^+ \tau^-) / \Gamma(J/\psi(1S) \pi^+ \pi^-)$ Γ_8 / Γ_{11}

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
8.8 ± 1.1 OUR FIT			
8.73 ± 1.39 ± 1.57	BAI	02	BES $e^+ e^-$

$\Gamma(J/\psi(1S) \pi^+ \pi^-) / \Gamma(J/\psi(1S) \text{anything})$ Γ_{11} / Γ_9

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.5645 ± 0.0026 OUR FIT				

0.554 ± 0.008 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

0.5604 ± 0.0009 ± 0.0062	565k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \pi^+ \pi^-$
0.525 ± 0.009 ± 0.022	4k	ANDREOTTI	05	E835	$\psi(2S) \rightarrow J/\psi X$
0.536 ± 0.007 ± 0.016	20k	^{1,2} ABLIKIM	04B	BES	$\psi(2S) \rightarrow J/\psi X$
0.496 ± 0.037		ARMSTRONG	97	E760	$\bar{p} p \rightarrow \psi(2S)$


• • • We do not use the following data for averages, fits, limits, etc. • • •

0.5637 ± 0.0027 ± 0.0046 60k ADAM 05A CLEO Repl. by MENDEZ 08

¹ From a fit to the J/ψ recoil mass spectra.

² ABLIKIM 04B quotes $B(\psi(2S) \rightarrow J/\psi X) / B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-)$.

WEIGHTED AVERAGE
0.554 ± 0.008 (Error scaled by 1.3)



$\Gamma(J/\psi(1S) \pi^+ \pi^-) / \Gamma(J/\psi(1S) \text{anything})$ Γ_{11} / Γ_9

$\Gamma(J/\psi(1S) \text{neutrals}) / \Gamma(J/\psi(1S) \pi^+ \pi^-)$
 $\Gamma_{10} / \Gamma_{11} = (0.9761 \Gamma_{12} + 0.719 \Gamma_{13} + 0.343 \Gamma_{140} + 0.190 \Gamma_{141}) / \Gamma_{11}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.732 ± 0.008 OUR FIT			
0.73 ± 0.09	TANENBAUM	76	MRK1 $e^+ e^-$

$$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma_{\text{total}} \qquad \Gamma_{12}/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.1823 ± 0.0031 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.1769 ± 0.0008 ± 0.0053	61k	¹ MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$
0.1652 ± 0.0014 ± 0.0058	13.4k	² ADAM	05A	CLEO	Repl. by MENDEZ 08

¹ Not independent from other measurements of MENDEZ 08.

² Not independent from other values reported by ADAM 05A.

$$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\text{anything}) \qquad \Gamma_{12}/\Gamma_9$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.2968 ± 0.0031 OUR FIT

0.320 ± 0.012 OUR AVERAGE

0.300 ± 0.008 ± 0.022	1655 ± 44	ANDREOTTI	05	E835	$\psi(2S) \rightarrow J/\psi X$
0.328 ± 0.013 ± 0.008		AMBROGIANI	00A	E835	$p\bar{p} \rightarrow \psi(2S)$
0.323 ± 0.033		ARMSTRONG	97	E760	$\bar{p}p \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.2829 ± 0.0012 ± 0.0056	61k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$
0.2776 ± 0.0025 ± 0.0043	13.4k	ADAM	05A	CLEO	Repl. by MENDEZ 08

$$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-) \qquad \Gamma_{12}/\Gamma_{11}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.526 ± 0.008 OUR FIT

0.513 ± 0.022 OUR AVERAGE Error includes scale factor of 2.2.

0.5047 ± 0.0022 ± 0.0102	61k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$
0.570 ± 0.009 ± 0.026	14k	¹ ABLIKIM	04B	BES	$\psi(2S) \rightarrow J/\psi X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.4924 ± 0.0047 ± 0.0086	73k	^{2,3} ADAM	05A	CLEO	Repl. by MENDEZ 08
0.571 ± 0.018 ± 0.044		⁴ ANDREOTTI	05	E835	$\psi(2S) \rightarrow J/\psi X$
0.53 ± 0.06		TANENBAUM	76	MRK1	$e^+ e^-$
0.64 ± 0.15		⁵ HILGER	75	SPEC	$e^+ e^-$

¹ From a fit to the J/ψ recoil mass spectra.

² Not independent from other values reported by ADAM 05A.

³ Using 13,217 $J/\psi\pi^0\pi^0$ and 60,010 $J/\psi\pi^+\pi^-$ events.

⁴ Not independent from other values reported by ANDREOTTI 05.

⁵ Ignoring the $J/\psi(1S)\eta$ and $J/\psi(1S)\gamma\gamma$ decays.

$$\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}} \qquad \Gamma_{13}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

33.7 ± 0.5 OUR FIT


32.9 ± 1.7 OUR AVERAGE Error includes scale factor of 2.1. See the ideogram below.

33.75 ± 0.17 ± 0.86	68.2k	ABLIKIM	12M	BES3	$e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$
29.8 ± 0.9 ± 2.3	5.7k	BAI	04I	BES2	$\psi(2S) \rightarrow J/\psi\gamma\gamma$
25.5 ± 2.9	386	¹ OREGLIA	80	CBAL	$e^+ e^- \rightarrow J/\psi 2\gamma$
45 ± 12	17	² BRANDELIK	79B	DASP	$e^+ e^- \rightarrow J/\psi 2\gamma$
42 ± 6	164	² BARTEL	78B	CNTR	$e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

34.3 ± 0.4 ± 0.9	18.4k	³ MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \eta$
32.5 ± 0.6 ± 1.1	2.8k	⁴ ADAM	05A	CLEO	Repl. by MENDEZ 08
43 ± 8	44	TANENBAUM	76	MRK1	$e^+ e^-$

- ¹ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.
- ² Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.
- ³ Not independent from other measurements of MENDEZ 08.
- ⁴ Not independent from other values reported by ADAM 05A.




$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\text{anything})$

Γ_{13}/Γ_9

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0549±0.0008 OUR FIT				
0.058 ±0.007 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
0.050 ±0.006 ±0.003	298 ± 20	ANDREOTTI 05	E835	$\psi(2S) \rightarrow J/\psi X$
0.072 ±0.009		AMBROGIANI 00A	E835	$p\bar{p} \rightarrow \psi(2S)$
0.061 ±0.015		ARMSTRONG 97	E760	$\bar{p}p \rightarrow \psi(2S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.0549±0.0006±0.0009	18.4k	¹ MENDEZ	08 CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \eta$
0.0546±0.0010±0.0007	2.8k	ADAM	05A CLEO	Repl. by MENDEZ 08

¹ Not independent from other measurements of MENDEZ 08.

WEIGHTED AVERAGE
 0.058 ± 0.007 (Error scaled by 1.4)



$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

Γ_{13}/Γ_{11}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0972 ± 0.0014				OUR FIT
0.0979 ± 0.0018				OUR AVERAGE
$0.0979 \pm 0.0010 \pm 0.0015$	18.4k	MENDEZ 08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \eta$
$0.098 \pm 0.005 \pm 0.010$	2k	¹ ABLIKIM 04B	BES	$\psi(2S) \rightarrow J/\psi X$
0.091 ± 0.021		² HIMEL 80	MRK2	$e^+ e^- \rightarrow \psi(2S) X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$0.0968 \pm 0.0019 \pm 0.0013$	2.8k	³ ADAM 05A	CLEO	Repl. by MENDEZ 08
$0.095 \pm 0.007 \pm 0.007$		⁴ ANDREOTTI 05	E835	$\psi(2S) \rightarrow J/\psi X$

¹ From a fit to the J/ψ recoil mass spectra.

² The value for $B(\psi(2S) \rightarrow J/\psi(1S)\eta)$ reported in HIMEL 80 is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.138 \pm 0.018$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = (0.1181 \pm 0.0020)$.

³ Not independent from other values reported by ADAM 05A.

⁴ Not independent from other values reported by ANDREOTTI 05.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma_{\text{total}}$

Γ_{14}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
12.68 ± 0.32				OUR AVERAGE
$12.6 \pm 0.2 \pm 0.3$	4.1k	ABLIKIM 12M	BES3	$e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$
$13.3 \pm 0.8 \pm 0.3$	530	MENDEZ 08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\gamma$
$14.3 \pm 1.4 \pm 1.2$	280	BAI 04I	BES2	$\psi(2S) \rightarrow J/\psi \gamma \gamma$
14 ± 6	7	HIMEL 80	MRK2	$e^+ e^-$
$9 \pm 2 \pm 1$	23	¹ OREGLIA 80	CBAL	$\psi(2S) \rightarrow J/\psi 2\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$13 \pm 1 \pm 1$	88	ADAM 05A	CLEO	Repl. by MENDEZ 08

¹ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\text{anything})$

$\Gamma_{14}/\Gamma_9 = \Gamma_{14}/(\Gamma_{11} + \Gamma_{12} + \Gamma_{13} + 0.343\Gamma_{140} + 0.190\Gamma_{141})$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$0.213 \pm 0.012 \pm 0.003$	527	¹ MENDEZ	08	CLEO $e^+ e^- \rightarrow J/\psi \gamma \gamma$
$0.22 \pm 0.02 \pm 0.01$		² ADAM	05A	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi \gamma \gamma$

¹ Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.
² Not independent from other values reported by ADAM 05A.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\pi^+ \pi^-)$

Γ_{14}/Γ_{11}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$0.380 \pm 0.022 \pm 0.005$	527	¹ MENDEZ	08	CLEO $e^+ e^- \rightarrow J/\psi \gamma \gamma$
$0.39 \pm 0.04 \pm 0.01$		² ADAM	05A	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi \gamma \gamma$

¹ Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.
² Not independent from other values reported by ADAM 05A.

———— HADRONIC DECAYS ————

$\Gamma(\pi^0 h_c(1P))/\Gamma_{\text{total}}$

Γ_{15}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.6 ± 1.3 OUR AVERAGE				
$9.0 \pm 1.5 \pm 1.3$	3k	¹ GE	11	CLEO $\psi(2S) \rightarrow \pi^0 \text{ anything}$
$8.4 \pm 1.3 \pm 1.0$	11k	ABLIKIM	10B	BES3 $\psi(2S) \rightarrow \pi^0 h_c$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
seen	92^{+23}_{-22}	ADAMS	09	CLEO $\psi(2S) \rightarrow 2\pi^+ 2\pi^- 2\pi^0$
seen	1282	DOBBS	08A	CLEO $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
seen	168 ± 40	ROSNER	05	CLEO $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$

¹ Assuming a width $\Gamma(h_c(1P)) = 0.86 \text{ MeV} \equiv \Gamma_0$, a measured dependence of the central value of $B = (7.6 + 1.4 \times \Gamma(h_c(1P)/\Gamma_0) \times 10^{-4})$, and with a systematic error that accounts for the width variation range 0.43–1.29 MeV.

$\Gamma(3(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}$

Γ_{16}/Γ


VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
35 ± 16	6	FRANKLIN	83	MRK2 $e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(2(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}$

Γ_{17}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
29 ± 10 OUR AVERAGE				
$24.9 \pm 0.7 \pm 3.6$	2173	ABLIKIM	07D	BES2 $e^+ e^- \rightarrow \psi(2S)$
$127 \pm 12 \pm 2$	410	¹ AUBERT	07AU	BABR $10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-)\pi^0 \gamma$
$26.1 \pm 0.7 \pm 3.0$	1703	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-)\pi^0$
30 ± 8	42	FRANKLIN	83	MRK2 $e^+ e^-$

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0) / \Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (297 \pm 22 \pm 18) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.



$\Gamma(\rho a_2(1320)) / \Gamma_{\text{total}}$

Γ_{18} / Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.55±0.73±0.47		112 ± 31	BAI	04C BES2	$\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<2.3		90	BAI	98J BES	$e^+ e^-$

$\Gamma(\rho \bar{p}) / \Gamma_{\text{total}}$

Γ_{19} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.88±0.10 OUR FIT					
3.00±0.13 OUR AVERAGE	Error includes scale factor of 1.1.				
3.08±0.05±0.18	4.5k	¹ DOBBS	14	$e^+ e^- \rightarrow \psi(2S) \rightarrow \rho \bar{p}$	
3.36±0.09±0.25	1.6k	ABLIKIM	07C BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow \rho \bar{p}$	
2.87±0.12±0.15	557	PEDLAR	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow \rho \bar{p}$	
1.4 ±0.8	4	BRANDELIK	79C DASP	$e^+ e^- \rightarrow \psi(2S) \rightarrow \rho \bar{p}$	
2.3 ±0.7		FELDMAN	77 MRK1	$e^+ e^- \rightarrow \psi(2S) \rightarrow \rho \bar{p}$	

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(p\bar{p})/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_{19}/Γ_{11}

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
8.31 ± 0.28 OUR FIT			
$6.98 \pm 0.49 \pm 0.97$	BAI	01	BES $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$

$\Gamma(\Delta^{++}\bar{\Delta}^{--})/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$12.8 \pm 1.0 \pm 3.4$	157	¹ BAI	01	BES $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

¹ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.29	90	¹ ABLIKIM	13F	BES3 $\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<12	90	² ABLIKIM	07H	BES2 $e^+e^- \rightarrow \psi(2S)$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\pi^0 \rightarrow \gamma\gamma) = 98.8\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.4\%$.

$\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.48 \pm 0.34 \pm 0.19$		60	¹ ABLIKIM	13F	BES3 $\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<4.9	90		² ABLIKIM	07H	BES2 $e^+e^- \rightarrow \psi(2S)$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.31\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$.

$\Gamma(\Lambda\bar{p}K^+)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.0 \pm 0.1 \pm 0.1$	74.0	BRIERE	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow$ $p\bar{p}K^+\pi^-$

$\Gamma(\Lambda\bar{p}K^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.8 \pm 0.3 \pm 0.3$	45.8	BRIERE	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow$ $p\bar{p}K^+\pi^+\pi^-\pi^-$

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.8 \pm 0.4 \pm 0.5$	73.4	BRIERE	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow$ $p\bar{p}2(\pi^+\pi^-)$

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$

Γ_{26}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.81±0.13 OUR AVERAGE			Error includes scale factor of 1.4. See the ideogram below.		
3.97±0.02±0.12	31k		ABLIKIM	17L BES3	$e^+e^- \rightarrow \Lambda\bar{\Lambda}$
3.71±0.05±0.15	6.5k		¹ DOBBS	17	$e^+e^- \rightarrow \Lambda\bar{\Lambda}$
3.39±0.20±0.32	337		ABLIKIM	07C BES	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
6.4 ±1.8 ±0.1			² AUBERT	07BD BABR	10.6 $e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$
3.28±0.23±0.25	208		PEDLAR	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
3.75±0.09±0.23	1.9k		^{1,3} DOBBS	14	$e^+e^- \rightarrow \Lambda\bar{\Lambda}$
1.81±0.20±0.27	80		⁴ BAI	01 BES	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
< 4	90		FELDMAN	77 MRK1	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$


¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² AUBERT 07BD reports $[\Gamma(\psi(2S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (15 \pm 4 \pm 1) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Superseded by DOBBS 17.

⁴ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

WEIGHTED AVERAGE
3.81±0.13 (Error scaled by 1.4)



$\Gamma(\Lambda\bar{\Sigma}^+\pi^- + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{27}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.40±0.03±0.13	2.8k	ABLIKIM	13W BES3	$\psi(2S) \rightarrow \text{hadrons}$

$\Gamma(\Lambda\bar{\Sigma}^-\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.54±0.04±0.13	2.8k	ABLIKIM	13W BES3	$\psi(2S) \rightarrow \text{hadrons}$

 $\Gamma(\Lambda\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	COMMENT
1.23±0.23±0.08	30	¹ DOBBS 17	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration. $\Gamma(\Sigma^0\bar{p}K^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.67±0.13±0.12	276	¹ ABLIKIM 13D	BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{p}K^+$

¹ Using $B(\Lambda \rightarrow p\pi^-) = 63.9\%$, and $B(\Sigma^0 \rightarrow \Lambda\gamma) = 100\%$. $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.32±0.12 OUR AVERAGE				

2.31±0.06±0.10 1.9k ¹ DOBBS 17 $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ 2.57±0.44±0.68 35 PEDLAR 05 CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.51±0.15±0.16 281 ^{1,2} DOBBS 14 $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ ¹ Using CLEO-c data but not authored by the CLEO Collaboration.² Superseded by DOBBS 17. $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.35±0.09 OUR AVERAGE				Error includes scale factor of 1.1.

2.44±0.03±0.11 7k ABLIKIM 17L BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ 2.22±0.05±0.11 2.6k ¹ DOBBS 17 $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ 2.35±0.36±0.32 59 ABLIKIM 07C BES $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ 2.63±0.35±0.21 58 PEDLAR 05 CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.25±0.11±0.16 439 ^{1,2} DOBBS 14 $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ 1.2 ±0.4 ±0.4 8 ³ BAI 01 BES $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ ¹ Using CLEO-c data but not authored by the CLEO Collaboration.² Superseded by DOBBS 17.³ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$. $\Gamma(\Sigma(1385)^+\bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
8.5±0.7 OUR AVERAGE				

8.4±0.5±0.5 1.5k ABLIKIM 16L BES3 $\psi(2S) \rightarrow \Sigma(1385)^+\bar{\Sigma}(1385)^-$ 11 ±3 ±3 14 ¹ BAI 01 BES $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$ ¹ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

$\Gamma(\Sigma(1385)^-\bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$8.5 \pm 0.6 \pm 0.6$	1.4K	ABLIKIM 16L	BES3	$\psi(2S) \rightarrow \Sigma(1385)^-\bar{\Sigma}(1385)^+$

$\Gamma(\Sigma(1385)^0\bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.69 \pm 0.05 \pm 0.05$	2.2k	ABLIKIM 17E	BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

$\Gamma(\Xi^-\bar{\Xi}^+)/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.87 ± 0.11 OUR AVERAGE			Error includes scale factor of 1.1.		
$3.03 \pm 0.05 \pm 0.14$	3.6k	¹ DOBBS 17			$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
$2.78 \pm 0.05 \pm 0.14$	5k	ABLIKIM 16L	BES3		$\psi(2S) \rightarrow \Xi^-\bar{\Xi}^+$
$3.03 \pm 0.40 \pm 0.32$	67	ABLIKIM 07C	BES		$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
$2.38 \pm 0.30 \pm 0.21$	63	PEDLAR 05	CLEO		$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.66 \pm 0.12 \pm 0.20$	548	^{1,2} DOBBS 14			$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
$0.94 \pm 0.27 \pm 0.15$	12	³ BAI 01	BES		$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
<2	90	FELDMAN 77	MRK1		$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Superseded by DOBBS 17.

³ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

$\Gamma(\Xi^0\bar{\Xi}^0)/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.3 ± 0.4 OUR AVERAGE		Error includes scale factor of 4.2.		
$2.73 \pm 0.03 \pm 0.13$	11k	ABLIKIM 17E	BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
$1.97 \pm 0.06 \pm 0.11$	1.2k	¹ DOBBS 17		$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
$2.75 \pm 0.64 \pm 0.61$	19	PEDLAR 05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.02 \pm 0.19 \pm 0.15$	112	^{1,2} DOBBS 14			$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
--------------------------	-----	-------------------------	--	--	---

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Superseded by DOBBS 17.

$\Gamma(\Xi(1530)^0\bar{\Xi}(1530)^0)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
5.2 ± 0.3 ^{+3.2}_{-1.2}		527	¹ ABLIKIM 13S	BES3	$\psi(2S) \rightarrow \eta p \bar{p}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<32	90	PEDLAR 05	CLEO		$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons
< 8.1	90	² BAI 01	BES		$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

¹ With $N(1535)$ decaying to $p\eta$.

² Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

$\Gamma(K^- \Lambda_{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.86 \pm 0.27 \pm 0.32$	236	ABLIKIM 15I	BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Lambda_{\Xi}^+ + \text{c.c.}$

 $\Gamma(\Xi(1690)^- \Xi^+ \rightarrow K^- \Lambda_{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.21 \pm 1.48 \pm 0.57$	74	ABLIKIM 15I	BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Lambda_{\Xi}^+ + \text{c.c.}$

 $\Gamma(\Xi(1820)^- \Xi^+ \rightarrow K^- \Lambda_{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$12.03 \pm 2.94 \pm 1.22$	136	ABLIKIM 15I	BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Lambda_{\Xi}^+ + \text{c.c.}$

 $\Gamma(K^- \Sigma^0 \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.67 \pm 0.33 \pm 0.28$	142	ABLIKIM 15I	BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Sigma^0 \Xi^+ + \text{c.c.}$

 $\Gamma(\Omega^- \bar{\Omega}^+)/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$0.52 \pm 0.03 \pm 0.03$		326	¹ DOBBS 17		$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.47 \pm 0.09 \pm 0.05$		27	^{1,2} DOBBS 14		$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
<1.5	90		ABLIKIM 12Q	BES2	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
<1.6	90		PEDLAR 05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
<0.73	90		³ BAI 01	BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.² Superseded by DOBBS 17.³ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$. $\Gamma(\pi^0 \rho \bar{\rho})/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.53 ± 0.07 OUR AVERAGE				
$1.65 \pm 0.03 \pm 0.15$	4.5k	ABLIKIM 13A	BES3	$\psi(2S) \rightarrow \rho \bar{\rho} \pi^0$
$1.54 \pm 0.06 \pm 0.06$	948	ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \pi^0 \rho \bar{\rho}$
$1.32 \pm 0.10 \pm 0.15$	256	¹ ABLIKIM 05E	BES2	$e^+ e^- \rightarrow \psi(2S) \rightarrow \rho \bar{\rho} \gamma \gamma$
1.4 ± 0.5	9	FRANKLIN 83	MRK2	$e^+ e^-$

¹ Computed using $B(\pi^0 \rightarrow \gamma \gamma) = (98.80 \pm 0.03)\%$. $\Gamma(N(940) \bar{p} + \text{c.c.} \rightarrow \pi^0 \rho \bar{\rho})/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
6.42 ± 0.20 $\begin{smallmatrix} +1.78 \\ -1.28 \end{smallmatrix}$	1.9k	¹ ABLIKIM 13A	BES3	$\psi(2S) \rightarrow \rho \bar{\rho} \pi^0$

¹ From a fit of $\pi^0 \rho \bar{\rho}$ data to eight distinct intermediate $N \bar{p}$ resonant states.

$\Gamma(N(1440)\bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p})/\Gamma_{\text{total}}$ **Γ_{46}/Γ**

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

7.3 $^{+1.7}_{-1.5}$ OUR AVERAGE Error includes scale factor of 2.5.

3.58 ± 0.25 $^{+1.59}_{-0.84}$	1.1k	¹ ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p \bar{p} \pi^0$
8.1 ± 0.7 ± 0.3	474	² ALEXANDER	10	CLEO $\psi(2S) \rightarrow \pi^0 p \bar{p}$

¹ From a fit of $\pi^0 p \bar{p}$ data to eight distinct intermediate $N \bar{p}$ resonant states.² From a fit of the $p \bar{p}$ and $p \pi^0$ mass distributions to a combination of $N(1440)\bar{p}$, $\pi^0 f_0(2100)$, and two other broad, unestablished resonances. **$\Gamma(N(1520)\bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p})/\Gamma_{\text{total}}$** **$\Gamma_{47}/\Gamma$**

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.64 ± 0.05 $^{+0.22}_{-0.17}$	0.2k	¹ ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p \bar{p} \pi^0$
------------------------------------	------	----------------------	-----	---

¹ From a fit of $\pi^0 p \bar{p}$ data to eight distinct intermediate $N \bar{p}$ resonant states. **$\Gamma(N(1535)\bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p})/\Gamma_{\text{total}}$** **$\Gamma_{48}/\Gamma$**

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.47 ± 0.28 $^{+0.99}_{-0.97}$	0.7k	¹ ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p \bar{p} \pi^0$
------------------------------------	------	----------------------	-----	---

¹ From a fit of $\pi^0 p \bar{p}$ data to eight distinct intermediate $N \bar{p}$ resonant states. **$\Gamma(N(1650)\bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p})/\Gamma_{\text{total}}$** **$\Gamma_{49}/\Gamma$**

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

3.76 ± 0.28 $^{+1.37}_{-1.66}$	1.1k	¹ ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p \bar{p} \pi^0$
------------------------------------	------	----------------------	-----	---

¹ From a fit of $\pi^0 p \bar{p}$ data to eight distinct intermediate $N \bar{p}$ resonant states. **$\Gamma(N(1720)\bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p})/\Gamma_{\text{total}}$** **$\Gamma_{50}/\Gamma$**

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.79 ± 0.10 $^{+0.24}_{-0.71}$	0.5k	¹ ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p \bar{p} \pi^0$
------------------------------------	------	----------------------	-----	---

¹ From a fit of $\pi^0 p \bar{p}$ data to eight distinct intermediate $N \bar{p}$ resonant states. **$\Gamma(N(2300)\bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p})/\Gamma_{\text{total}}$** **$\Gamma_{51}/\Gamma$**

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.62 ± 0.28 $^{+1.12}_{-0.64}$	0.9k	¹ ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p \bar{p} \pi^0$
------------------------------------	------	----------------------	-----	---

¹ From a fit of $\pi^0 p \bar{p}$ data to eight distinct intermediate $N \bar{p}$ resonant states. **$\Gamma(N(2570)\bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p})/\Gamma_{\text{total}}$** **$\Gamma_{52}/\Gamma$**

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.13 ± 0.08 $^{+0.40}_{-0.30}$	0.8k	¹ ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p \bar{p} \pi^0$
------------------------------------	------	----------------------	-----	---

¹ From a fit of $\pi^0 p \bar{p}$ data to eight distinct intermediate $N \bar{p}$ resonant states.

$\Gamma(\pi^0 f_0(2100) \rightarrow \pi^0 p\bar{p})/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.1±0.4±0.1	76	¹ ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \pi^0 p\bar{p}$

¹ From a fit of the $p\bar{p}$ and $p\pi^0$ mass distributions to a combination of $N_1^*(1440)\bar{p}$, $\pi^0 f_0(2100)$, and two other broad, unestablished resonances.

 $\Gamma(\eta p\bar{p})/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
6.0±0.4 OUR AVERAGE				
6.4±0.2±0.6	679	¹ ABLIKIM 13S	BES3	$\psi(2S) \rightarrow \eta p\bar{p}$
5.6±0.6±0.3	154	¹ ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \eta p\bar{p}$
5.8±1.1±0.7	44.8 ± 8.5	² ABLIKIM 05E	BES2	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\gamma\gamma$
8 ±3 ±3	9.8	BRIERE 05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$

¹ With $N(1535)$ decaying to $p\eta$.

² Computed using $B(\eta \rightarrow \gamma\gamma) = (39.43 \pm 0.26)\%$.

 $\Gamma(\eta f_0(2100) \rightarrow \eta p\bar{p})/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.2±0.4±0.1	31	¹ ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \eta p\bar{p}$

¹ From a fit of the $p\bar{p}$ and $p\eta$ distributions to a combination of $N^*(1535)\bar{p}$ and $\eta f_0(2100)$.

 $\Gamma(N(1535)\bar{p} \rightarrow \eta p\bar{p})/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
4.4±0.6±0.3	123	¹ ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \eta p\bar{p}$

¹ From a fit of the $p\bar{p}$ and $p\eta$ distributions to a combination of $N^*(1535)\bar{p}$ and $\eta f_0(2100)$.

 $\Gamma(\omega p\bar{p})/\Gamma_{\text{total}}$ Γ_{57}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.69±0.21 OUR AVERAGE				
0.6 ±0.2 ±0.2	21.2	BRIERE 05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$
0.8 ±0.3 ±0.1	14.9 ± 0.1	¹ BAI 03B	BES	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

 $\Gamma(\phi p\bar{p})/\Gamma_{\text{total}}$ Γ_{58}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.24	90	BRIERE 05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.26	90	¹ BAI 03B	BES	$\psi(2S) \rightarrow K^+K^-p\bar{p}$
-------	----	----------------------	-----	---------------------------------------

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

$\Gamma(\pi^+ \pi^- \rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{59}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.0 ± 0.4	OUR AVERAGE			
$5.9 \pm 0.2 \pm 0.4$	904.5	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \rho\bar{\rho}\pi^+\pi^-$
8 ± 2		¹ TANENBAUM	78	MRK1 $e^+ e^-$

¹ Assuming entirely strong decay.

$\Gamma(\rho\bar{\rho}\pi^- \text{ or c.c.})/\Gamma_{\text{total}}$ Γ_{60}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.48 ± 0.17	OUR AVERAGE			
$2.45 \pm 0.11 \pm 0.21$	851	ABLIKIM	06i	BES2 $e^+ e^- \rightarrow \rho\pi^- X$
$2.52 \pm 0.12 \pm 0.22$	849	ABLIKIM	06i	BES2 $e^+ e^- \rightarrow \bar{\rho}\pi^+ X$

$\Gamma(\rho\bar{\rho}\pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{61}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.18 \pm 0.50 \pm 0.50$	135 ± 21	ABLIKIM	06i	BES2 $e^+ e^- \rightarrow \rho\pi^- \pi^0 X$

$\Gamma(\eta\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{63}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

$\Gamma(\eta\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{64}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$9.5 \pm 0.7 \pm 1.5$		¹ BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadr}$
$10.3 \pm 0.8 \pm 1.4$	201.7	² BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \eta 3\pi(\eta \rightarrow \gamma\gamma)$
$8.1 \pm 1.4 \pm 1.6$	50.0	² BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \eta 3\pi(\eta \rightarrow 3\pi)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Average of $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow 3\pi$.

² Not independent from other values reported by BRIERE 05.

$\Gamma(2(\pi^+ \pi^-)\eta)/\Gamma_{\text{total}}$ Γ_{65}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.2 \pm 0.6 \pm 0.1$	16	¹ AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-)\eta\gamma$

¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow 2(\pi^+ \pi^-)\eta) \cdot B(\eta \rightarrow \gamma\gamma) = 1.2 \pm 0.7 \pm 0.1 \text{ eV}$.

$\Gamma(\eta'\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{66}/Γ


VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.5 \pm 1.6 \pm 1.3$	12.8	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadr}$

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{67}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.3±1.2 OUR AVERAGE				Error includes scale factor of 2.1. See the ideogram below.
8.4±0.5±1.2	386	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$
12.2±2.2±0.7	37	¹ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
8.2±0.5±0.7	391	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
4.8±0.6±0.7	100 ± 22	² BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow \omega\pi^+\pi^-) \cdot B(\omega \rightarrow 3\pi) = 2.69 \pm 0.73 \pm 0.16$ eV.				
² Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.				

WEIGHTED AVERAGE
7.3±1.2 (Error scaled by 2.1)



$\Gamma(b_1^\pm\pi^\mp)/\Gamma_{\text{total}}$

Γ_{68}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.0 ± 0.6 OUR AVERAGE				Error includes scale factor of 1.1.
5.1 ± 0.6 ± 0.8	202	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$
4.18 ^{+0.43} _{-0.42} ± 0.92	170	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$
3.2 ± 0.6 ± 0.5	61 ± 11	^{1,2} BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
5.2 ± 0.8 ± 1.0		¹ BAI	99C BES	Repl. by BAI 03B
¹ Assuming $B(b_1 \rightarrow \omega\pi) = 1$.				
² Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.				

$\Gamma(b_1^0 \pi^0)/\Gamma_{\text{total}}$					Γ_{69}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
$2.35^{+0.47}_{-0.42} \pm 0.40$	45	ADAM	05	CLEO	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$					Γ_{70}/Γ
VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.2 ± 0.4	OUR AVERAGE				
$2.3 \pm 0.5 \pm 0.4$		57	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$
$2.05 \pm 0.41 \pm 0.38$		62 ± 12	BAI	04C BES2	$\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 1.5		90	¹ BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$
< 1.7		90	BAI	98J BES	Repl. by BAI 03B

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

$\Gamma(\pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}$					Γ_{72}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
7.3 ± 0.5	OUR AVERAGE				
$8.1 \pm 1.3 \pm 0.3$	133	LEES	12F BABR	10.6	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
$7.1 \pm 0.3 \pm 0.4$	817.2	BRIERE	05 CLEO		$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
16 ± 4		¹ TANENBAUM	78 MRK1		$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$11.0 \pm 1.9 \pm 0.2$	85	² AUBERT	07AK BABR	10.6	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ Assuming entirely strong decay.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (2.56 \pm 0.42 \pm 0.16) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho^0 K^+ K^-)/\Gamma_{\text{total}}$					Γ_{74}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
$2.2 \pm 0.2 \pm 0.4$	223.8	BRIERE	05 CLEO		$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$

$\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$					Γ_{75}/Γ
VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$1.86 \pm 0.32 \pm 0.43$		93 ± 16	BAI	04C	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 1.2		90	BAI	98J BES	$e^+ e^-$

$\Gamma(K^+ K^- \pi^+ \pi^- \eta)/\Gamma_{\text{total}}$					Γ_{76}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
$1.3 \pm 0.7 \pm 0.1$	7	¹ AUBERT	07AU BABR	10.6	$e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$

¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow 2(\pi^+ \pi^-) \eta) \cdot B(\eta \rightarrow \gamma \gamma) = 1.2 \pm 0.7 \pm 0.1$ eV.

$\Gamma(K^+ K^- 2(\pi^+ \pi^-) \pi^0) / \Gamma_{\text{total}}$ Γ_{77} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
10.0 ± 2.5 ± 1.8	65	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(K_1(1270)^\pm K^\mp) / \Gamma_{\text{total}}$ Γ_{79} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
10.0 ± 1.8 ± 2.1	¹ BAI	99C BES	$e^+ e^-$

¹ Assuming $B(K_1(1270) \rightarrow K \rho) = 0.42 \pm 0.06$

 $\Gamma(K_S^0 K_S^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{80} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.20 ± 0.25 ± 0.37	83 ± 9	ABLIKIM	050 BES2	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\rho^0 \rho^0) / \Gamma_{\text{total}}$ Γ_{81} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.5 ± 0.1 ± 0.2	61.1	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow \rho \bar{\rho} \pi^+ \pi^-$

 $\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{82} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
6.7 ± 2.5	TANENBAUM 78	MRK1	$e^+ e^-$

 $\Gamma(2(\pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{83} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.4 ± 0.6 OUR AVERAGE	Error includes scale factor of 2.2.			
2.2 ± 0.2 ± 0.2	308	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-)$
4.5 ± 1.0		TANENBAUM 78	MRK1	$e^+ e^-$

 $\Gamma(\rho^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{84} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.2 ± 0.6 OUR AVERAGE	Error includes scale factor of 1.4.			
2.0 ± 0.2 ± 0.4	285.5	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-)$
4.2 ± 1.5		TANENBAUM 78	MRK1	$e^+ e^-$

 $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$ Γ_{85} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
12.6 ± 0.9 OUR AVERAGE				
18.9 ± 5.7 ± 0.3	32	¹ AUBERT	07AU BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \pi^0 \gamma$
11.7 ± 1.0 ± 1.5	597	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
12.7 ± 0.5 ± 1.0	711.6	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (44 \pm 13 \pm 3) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega f_0(1710) \rightarrow \omega K^+ K^-) / \Gamma_{\text{total}}$ Γ_{86} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.9 \pm 2.0 \pm 0.9$	19	ABLIKIM	06G	BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

$\Gamma(K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{87} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$8.6 \pm 1.3 \pm 1.8$	238	ABLIKIM	06G	BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

$\Gamma(K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{88} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$9.6 \pm 2.2 \pm 1.7$	133	ABLIKIM	06G	BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

$\Gamma(K^*(892)^+ K^- \rho^0 + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{89} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.3 \pm 2.2 \pm 1.4$	78	ABLIKIM	06G	BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

$\Gamma(K^*(892)^0 K^- \rho^+ + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{90} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.1 \pm 1.3 \pm 1.2$	125	ABLIKIM	06G	BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

$\Gamma(\eta K^+ K^-, \text{no } \eta\phi) / \Gamma_{\text{total}}$ Γ_{91} / Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$3.08 \pm 0.29 \pm 0.25$	0.3k	¹	ABLIKIM	12L	BES3 $\psi(2S) \rightarrow K^+ K^- \gamma \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<13	90	BRIERE	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
-----	----	--------	----	------	--

¹ Excluding $\eta\phi$.

$\Gamma(\omega K^+ K^-) / \Gamma_{\text{total}}$ Γ_{92} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.62 ± 0.11 OUR AVERAGE		Error includes scale factor of 1.1.		
$1.56 \pm 0.04 \pm 0.11$	2.8k	ABLIKIM	14G	BES3 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
$2.38 \pm 0.37 \pm 0.29$	78	ABLIKIM	06G	BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
$1.9 \pm 0.3 \pm 0.3$	76.8	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
$1.5 \pm 0.3 \pm 0.2$	23	¹ BAI	03B	BES $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

$\Gamma(\omega K^*(892)^+ K^- + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{93} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
20.7 ± 2.6 OUR AVERAGE				
$18.9 \pm 2.9 \pm 2.2$	396	ABLIKIM	13M	BES3 $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$
$22.6 \pm 3.0 \pm 2.4$	535	ABLIKIM	13M	BES3 $\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

$$\Gamma(\omega K_2^*(1430)^+ K^- + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{94}/\Gamma$$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.1 ± 1.2 OUR AVERAGE				
6.39 ± 1.50 ± 0.78	128	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$
5.86 ± 1.61 ± 0.83	143	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

$$\Gamma(\omega \bar{K}^*(892)^0 K^0)/\Gamma_{\text{total}} \quad \Gamma_{95}/\Gamma$$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
16.8 ± 2.5 ± 1.6	356	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$

$$\Gamma(\omega \bar{K}_2^*(1430)^0 K^0)/\Gamma_{\text{total}} \quad \Gamma_{96}/\Gamma$$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.82 ± 2.08 ± 0.72	116	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$

$$\Gamma(\omega X(1440) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{97}/\Gamma$$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.60 ± 0.27 ± 0.24	109	¹ ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$

¹ X(1440) compatible with $\eta(1405)$ and $\eta(1475)$. A $f_1(1420)$ is also possible.

$$\Gamma(\omega X(1440) \rightarrow \omega K^+ K^- \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{98}/\Gamma$$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.09 ± 0.20 ± 0.16	82	¹ ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

¹ X(1440) compatible with $\eta(1405)$ and $\eta(1475)$. A $f_1(1420)$ is also possible.

$$\Gamma(\omega f_1(1285) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{99}/\Gamma$$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.302 ± 0.098 ± 0.027	22	¹ ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$

¹ Statistical significance 4.5 σ . This measurement is equivalent to a limit of $< 0.478 \times 10^{-5}$ at 90% C.L.

$$\Gamma(\omega f_1(1285) \rightarrow \omega K^+ K^- \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{100}/\Gamma$$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.125 ± 0.070 ± 0.013	10	¹ ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

¹ Statistical significance 3.2 σ . This measurement is equivalent to a limit of $< 0.221 \times 10^{-5}$ at 90% C.L.

$$\Gamma(3(\pi^+ \pi^-))/\Gamma_{\text{total}} \quad \Gamma_{101}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.5 ± 2.0 OUR AVERAGE				Error includes scale factor of 2.8.

5.45 ± 0.42 ± 0.87	671	ABLIKIM	05H BES2	$e^+ e^- \rightarrow \psi(2S) \rightarrow 3(\pi^+ \pi^-)$
--------------------	-----	---------	----------	---

1.5 ± 1.0		¹ TANENBAUM	78 MRK1	$e^+ e^-$
-----------	--	------------------------	---------	-----------

¹ Assuming entirely strong decay.

$\Gamma(\rho\bar{\rho}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$					Γ_{102}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
$7.3 \pm 0.4 \pm 0.6$	434.9	BRIERE	05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{\rho}\pi^+\pi^-\pi^0$

$\Gamma(K^+K^-)/\Gamma_{\text{total}}$					Γ_{103}/Γ
VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$7.48 \pm 0.23 \pm 0.39$		1.3k	¹ METREVELI	12	$\psi(2S) \rightarrow K^+K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$6.2 \pm 1.5 \pm 0.2$	66	^{2,3} LEES	15J	BABR	$e^+e^- \rightarrow K^+K^-\gamma$
$8.3 \pm 1.5 \pm 0.2$	66	^{3,4} LEES	15J	BABR	$e^+e^- \rightarrow K^+K^-\gamma$
$6.3 \pm 0.6 \pm 0.3$		⁵ DOBBS	06A	CLEO	e^+e^-
10 ± 7		⁵ BRANDELIK	79C	DASP	e^+e^-
< 5	90	FELDMAN	77	MRK1	e^+e^-

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² $\sin\phi > 0$.

³ Using $\Gamma(\psi(2S) \rightarrow e^+e^-) = (2.37 \pm 0.04) \text{ keV}$.

⁴ $\sin\phi < 0$.

⁵ Interference with non-resonant K^+K^- production not taken into account.

$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$					Γ_{104}/Γ
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
5.34 ± 0.33 OUR AVERAGE					
$5.28 \pm 0.25 \pm 0.34$	478 ± 23	¹ METREVELI	12		$\psi(2S) \rightarrow K_S^0 K_L^0$
$5.8 \pm 0.8 \pm 0.4$		DOBBS	06A	CLEO	e^+e^-
$5.24 \pm 0.47 \pm 0.48$	156 ± 14	² BAI	04B	BES2	$\psi(2S) \rightarrow K_S^0 K_L^0 \rightarrow \pi^+\pi^-X$

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.


² Using $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6860 \pm 0.0027$.

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$					Γ_{105}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.01 ± 0.17 OUR AVERAGE					Error includes scale factor of 1.7. See the ideogram below.
$2.14 \pm 0.03^{+0.12}_{-0.11}$	7k	¹ ABLIKIM	12H	BES3	$e^+e^- \rightarrow \psi(2S)$
$1.81 \pm 0.18 \pm 0.19$	260 ± 19	² ABLIKIM	05J	BES2	$e^+e^- \rightarrow \psi(2S)$
$1.88^{+0.16}_{-0.15} \pm 0.28$	194	ADAM	05	CLEO	$e^+e^- \rightarrow \psi(2S)$
0.85 ± 0.46	4	FRANKLIN	83	MRK2	$e^+e^- \rightarrow \text{hadrons}$

¹ From $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$ events directly. The quoted systematic error includes a contribution of 4% (added in quadrature) from the uncertainty on the number of $\psi(2S)$ events.

² From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.

WEIGHTED AVERAGE
 2.01 ± 0.17 (Error scaled by 1.7)



$\Gamma(\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ **Γ_{106}/Γ**

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
1.94 ± 0.25 $+1.15$ -0.34	¹ ABLIKIM	05J BES2	$\psi(2S) \rightarrow \rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0$

¹ From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.

$\Gamma(\rho(770)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ **Γ_{107}/Γ**

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.32 ± 0.12 OUR AVERAGE					Error includes scale factor of 1.8.
$0.51 \pm 0.07 \pm 0.11$			¹ ABLIKIM	05J BES2	$\psi(2S) \rightarrow \rho(770)\pi \rightarrow \pi^+\pi^-\pi^0$
$0.24^{+0.08}_{-0.07} \pm 0.02$		22	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.83	90	1	FRANKLIN	83 MRK2	e^+e^-
<10	90		BARTEL	76 CNTR	e^+e^-
<10	90		² ABRAMS	75 MRK1	e^+e^-

¹ From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.

² Final state $\rho^0\pi^0$.

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{108}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.78 ± 0.26 OUR AVERAGE					
$0.76 \pm 0.25 \pm 0.06$		30	¹ METREVELI 12		$\psi(2S) \rightarrow \pi^+\pi^-$
8 ± 5			BRANDELIK 79C	DASP	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2.1	90		DOBBS 06A	CLEO	$e^+e^- \rightarrow \psi(2S)$
<5	90		FELDMAN 77	MRK1	e^+e^-

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration. Using $\psi(3770) \rightarrow \pi^+\pi^-$ for continuum subtraction.

$\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{109}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<3.1	90	¹ BAI 99C	BES	e^+e^-

¹ Assuming $B(K_1(1400) \rightarrow K^*\pi) = 0.94 \pm 0.06$

$\Gamma(K_2^*(1430)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{110}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.12 \pm 0.62 \pm_{-0.61}^{+1.13}$	251 ± 22	ABLIKIM 12L	BES3	$e^+e^- \rightarrow \psi(2S)$

$\Gamma(K^+K^-\pi^0)/\Gamma_{\text{total}}$ Γ_{111}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$4.07 \pm 0.16 \pm 0.26$		0.9k	ABLIKIM 12L	BES3	$e^+e^- \rightarrow \psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<8.9	90	1	FRANKLIN 83	MRK2	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(K^+K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{114}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.9 ± 0.4 OUR AVERAGE Error includes scale factor of 1.2.					
$3.18 \pm 0.30 \pm_{-0.31}^{+0.26}$		0.2k	ABLIKIM 12L	BES3	$e^+e^- \rightarrow \psi(2S)$
$2.9 \pm_{-1.7}^{+1.3} \pm 0.4$		9.6 ± 4.2	ABLIKIM 05I	BES2	$e^+e^- \rightarrow \psi(2S)$
$1.3 \pm_{-0.7}^{+1.0} \pm 0.3$		7	ADAM 05	CLEO	$e^+e^- \rightarrow \psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<5.4	90		FRANKLIN 83	MRK2	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(K^*(892)^0\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{115}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
10.9 ± 2.0 OUR AVERAGE				
$13.3 \pm_{-2.8}^{+2.4} \pm 1.7$	65.6 ± 9.0	ABLIKIM 05I	BES2	$e^+e^- \rightarrow \psi(2S)$
$9.2 \pm_{-2.2}^{+2.7} \pm 0.9$	25	ADAM 05	CLEO	$e^+e^- \rightarrow \psi(2S)$

$\Gamma(K^+ K^*(892)^- + c.c.) / \Gamma(K^*(892)^0 \bar{K}^0 + c.c.)$ $\Gamma_{114} / \Gamma_{115}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.16 ± 0.06 OUR AVERAGE			
0.22 ^{+0.10} _{-0.14}	ABLIKIM	05I	BES2 $e^+ e^- \rightarrow \psi(2S)$
0.14 ^{+0.08} _{-0.06}	ADAM	05	CLEO $e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\phi \pi^+ \pi^-) / \Gamma_{total}$ Γ_{116} / Γ


VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
1.18 ± 0.26 OUR AVERAGE				Error includes scale factor of 1.5. See the ideogram below.
2.3 ± 0.8 ± 0.1	19 ± 6	LEES	12F BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
0.9 ± 0.2 ± 0.1	47.6	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
1.5 ± 0.2 ± 0.2	51.5 ± 8.3	¹ BAI	03B BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2.44 ± 0.96 ± 0.04	10 ± 4	^{2,3} AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi \pi^+ \pi^-) / \Gamma_{total}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (0.57 \pm 0.22 \pm 0.04) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.

WEIGHTED AVERAGE
1.18 ± 0.26 (Error scaled by 1.5)



$\Gamma(\phi f_0(980) \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{117} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.75 ± 0.33 OUR AVERAGE				Error includes scale factor of 1.6.
1.5 ± 0.5 ± 0.1	12 ± 4	LEES	12F BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
0.6 ± 0.2 ± 0.1	18.4 ± 6.4	¹ BAI	03B BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.46 ± 0.71 ± 0.02	6 ± 3	^{2,3} AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

² Superseded by LEES 12F. AUBERT 07AK reports [$\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}$] × [$\Gamma(\psi(2S) \rightarrow e^+ e^-)$] = $(0.34 \pm 0.16 \pm 0.04) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.

$\Gamma(2(K^+ K^-)) / \Gamma_{\text{total}}$ Γ_{118} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.63 ± 0.13 OUR AVERAGE				
0.9 ± 0.4 ± 0.1	13	LEES	12F BABR	10.6 $e^+ e^- \rightarrow 2(K^+ K^-) \gamma$
0.6 ± 0.1 ± 0.1	59.2	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$

$\Gamma(\phi K^+ K^-) / \Gamma_{\text{total}}$ Γ_{119} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.70 ± 0.16 OUR AVERAGE				
0.8 ± 0.2 ± 0.1	36.8	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$
0.6 ± 0.2 ± 0.1	16.1 ± 5.0	¹ BAI	03B BES	$\psi(2S) \rightarrow 2(K^+ K^-)$

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

$\Gamma(2(K^+ K^-) \pi^0) / \Gamma_{\text{total}}$ Γ_{120} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.1 ± 0.2 ± 0.2	44.7	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-) \pi^0$

$\Gamma(\phi \eta) / \Gamma_{\text{total}}$ Γ_{121} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.10 ± 0.31 OUR AVERAGE				
3.14 ± 0.23 ± 0.23	0.2k	ABLIKIM	12L BES3	$e^+ e^- \rightarrow \psi(2S)$
2.0 $^{+1.5}_{-1.1}$ ± 0.4	6	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$
3.3 ± 1.1 ± 0.5	17	ABLIKIM	04K BES	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\phi \eta') / \Gamma_{\text{total}}$ Γ_{122} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.1 ± 1.4 ± 0.7	8	¹ ABLIKIM	04K BES	$e^+ e^- \rightarrow \psi(2S)$

¹ Calculated combining $\eta' \rightarrow \gamma \rho$ and $\eta \pi^+ \pi^-$ channels.

$\Gamma(\omega\eta')/\Gamma_{\text{total}}$ **Γ_{123}/Γ**

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.2^{+2.4}_{-2.0} \pm 0.7$	4	¹ ABLIKIM	04K	BES $e^+e^- \rightarrow \psi(2S)$

¹ Calculated combining $\eta' \rightarrow \gamma\rho$ and $\eta\pi^+\pi^-$ channels.

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ **Γ_{124}/Γ**

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.1 ± 0.6 OUR AVERAGE				
$2.5^{+1.2}_{-1.0} \pm 0.2$	14	ADAM	05	CLEO $e^+e^- \rightarrow \psi(2S)$
$1.87^{+0.68}_{-0.62} \pm 0.28$	14	ABLIKIM	04L	BES $e^+e^- \rightarrow \psi(2S)$

$\Gamma(\rho\eta')/\Gamma_{\text{total}}$ **Γ_{125}/Γ**

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.87^{+1.64}_{-1.11} \pm 0.33$	2	ABLIKIM	04L	BES $e^+e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.02 \pm 0.11 \pm 0.24$	143	¹ ABLIKIM	17AK	BES3 $e^+e^- \rightarrow \psi(2S)$
$0.569 \pm 0.128 \pm 0.236$	80	² ABLIKIM	17AK	BES3 $e^+e^- \rightarrow \psi(2S)$

¹ Destructive-interference solution of a partial wave analysis of the decay $\psi(2S) \rightarrow \pi^+\pi^-\eta'$.

² Constructive-interference solution of a partial wave analysis of the decay $\psi(2S) \rightarrow \pi^+\pi^-\eta'$.

$\Gamma(\rho\eta)/\Gamma_{\text{total}}$ **Γ_{126}/Γ**

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.2 ± 0.6 OUR AVERAGE				Error includes scale factor of 1.1.
$3.0^{+1.1}_{-0.9} \pm 0.2$	18	ADAM	05	CLEO $e^+e^- \rightarrow \psi(2S)$
$1.78^{+0.67}_{-0.62} \pm 0.17$	13	ABLIKIM	04L	BES $e^+e^- \rightarrow \psi(2S)$

$\Gamma(\omega\eta)/\Gamma_{\text{total}}$ **Γ_{127}/Γ**

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.1	90	ADAM	05	CLEO $e^+e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.1	90	ABLIKIM	04K	BES $e^+e^- \rightarrow \psi(2S)$
------	----	---------	-----	-----------------------------------

$\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$ **Γ_{128}/Γ**

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.04	90	ABLIKIM	12L	BES3 $e^+e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.7	90	ADAM	05	CLEO $e^+e^- \rightarrow \psi(2S)$
------	----	------	----	------------------------------------

<0.4	90	ABLIKIM	04K	BES $e^+e^- \rightarrow \psi(2S)$
------	----	---------	-----	-----------------------------------

$\Gamma(\eta_c \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$						Γ_{129}/Γ
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT		
<1.0	90	PEDLAR	07	CLEO	$e^+ e^- \rightarrow \psi(2S)$	
$\Gamma(p\bar{p}K^+K^-)/\Gamma_{\text{total}}$						Γ_{130}/Γ
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT		
$2.7 \pm 0.6 \pm 0.4$	30.1	BRIERE	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$	
$\Gamma(\bar{\Lambda}nK_S^0 + \text{c.c.})/\Gamma_{\text{total}}$						Γ_{131}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
$0.81 \pm 0.11 \pm 0.14$	50	¹ ABLIKIM	08C	BES2	$e^+ e^- \rightarrow J/\psi$	
¹ Using $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$.						
$\Gamma(\phi f_2'(1525))/\Gamma_{\text{total}}$						Γ_{132}/Γ
VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.44 \pm 0.12 \pm 0.11$	20 ± 6	BAI	04C		$\psi(2S) \rightarrow 2(K^+K^-)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.45	90	BAI	98J	BES	$e^+ e^- \rightarrow 2(K^+K^-)$	
$\Gamma(\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}}$						Γ_{133}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<0.88	90	BAI	04G	BES2	$e^+ e^-$	
$\Gamma(\Theta(1540)K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$						Γ_{134}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<1.0	90	BAI	04G	BES2	$e^+ e^-$	
$\Gamma(\Theta(1540)K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$						Γ_{135}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<0.70	90	BAI	04G	BES2	$e^+ e^-$	
$\Gamma(\bar{\Theta}(1540)K^+ n \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$						Γ_{136}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<2.6	90	BAI	04G	BES2	$e^+ e^-$	
$\Gamma(\bar{\Theta}(1540)K_S^0 p \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$						Γ_{137}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<0.60	90	BAI	04G	BES2	$e^+ e^-$	
$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$						Γ_{138}/Γ
VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT			
<0.046	¹ BAI	04D	BES	$e^+ e^-$		
¹ Forbidden by CP .						

————— RADIATIVE DECAYS —————

$\Gamma(\gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$ Γ_{139}/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
9.79 ± 0.20				OUR FIT
9.33 ± 0.26				OUR AVERAGE
9.389 ± 0.014 ± 0.332	4.7M	ABLIKIM	17U	BES3 $e^+e^- \rightarrow \gamma X$
9.22 ± 0.11 ± 0.46	72k	ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
9.9 ± 0.5 ± 0.8		¹ GAISER	86	CBAL $e^+e^- \rightarrow \gamma X$
7.2 ± 2.3		¹ BIDDICK	77	CNTR $e^+e^- \rightarrow \gamma X$
7.5 ± 2.6		¹ WHITAKER	76	MRK1 e^+e^-

¹ Angular distribution $(1+\cos^2\theta)$ assumed.

$\Gamma(\gamma\chi_{c1}(1P))/\Gamma_{\text{total}}$ Γ_{140}/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
9.75 ± 0.24				OUR FIT
9.54 ± 0.29				OUR AVERAGE
9.905 ± 0.011 ± 0.353	5.0M	ABLIKIM	17U	BES3 $e^+e^- \rightarrow \gamma X$
9.07 ± 0.11 ± 0.54	76k	ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
9.0 ± 0.5 ± 0.7		¹ GAISER	86	CBAL $e^+e^- \rightarrow \gamma X$
7.1 ± 1.9		² BIDDICK	77	CNTR $e^+e^- \rightarrow \gamma X$

¹ Angular distribution $(1-0.189 \cos^2\theta)$ assumed.

² Valid for isotropic distribution of the photon.


$\Gamma(\gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$ Γ_{141}/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
9.52 ± 0.20				OUR FIT
9.42 ± 0.31				OUR AVERAGE
Error includes scale factor of 1.3. See the ideogram below.				
9.621 ± 0.013 ± 0.272	4.2M	ABLIKIM	17U	BES3 $e^+e^- \rightarrow \gamma X$
9.33 ± 0.14 ± 0.61	79k	ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
8.0 ± 0.5 ± 0.7		¹ GAISER	86	CBAL $e^+e^- \rightarrow \gamma X$
7.0 ± 2.0		² BIDDICK	77	CNTR $e^+e^- \rightarrow \gamma X$

¹ Angular distribution $(1-0.052 \cos^2\theta)$ assumed.

² Valid for isotropic distribution of the photon.

WEIGHTED AVERAGE
 9.42 ± 0.31 (Error scaled by 1.3)



$\Gamma(\gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$ (units 10^{-2})

$[\Gamma(\gamma\chi_{c0}(1P)) + \Gamma(\gamma\chi_{c1}(1P)) + \Gamma(\gamma\chi_{c2}(1P))]/\Gamma_{\text{total}} (\Gamma_{139} + \Gamma_{140} + \Gamma_{141})/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$27.6 \pm 0.3 \pm 2.0$	¹ ATHAR	04	CLEO	$e^+e^- \rightarrow \gamma X$
------------------------	--------------------	----	------	-------------------------------

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.

$\Gamma(\gamma\chi_{c0}(1P))/\Gamma(\gamma\chi_{c1}(1P))$ $\Gamma_{139}/\Gamma_{140}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.02 \pm 0.01 \pm 0.07$	¹ ATHAR	04	CLEO	$e^+e^- \rightarrow \gamma X$
--------------------------	--------------------	----	------	-------------------------------

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.

$\Gamma(\gamma\chi_{c2}(1P))/\Gamma(\gamma\chi_{c1}(1P))$ $\Gamma_{141}/\Gamma_{140}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.03 \pm 0.02 \pm 0.03$	¹ ATHAR	04	CLEO	$e^+e^- \rightarrow \gamma X$
--------------------------	--------------------	----	------	-------------------------------

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.

$\Gamma(\gamma\chi_{c0}(1P))/\Gamma(\gamma\chi_{c2}(1P))$ $\Gamma_{139}/\Gamma_{141}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.99 \pm 0.02 \pm 0.08$	¹ ATHAR	04	CLEO	$e^+e^- \rightarrow \gamma X$
--------------------------	--------------------	----	------	-------------------------------

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.

$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$


Γ_{142}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.34 ± 0.05	OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.		
0.432 ± 0.016 ± 0.060		MITCHELL	09	CLEO $e^+e^- \rightarrow \gamma X$
0.32 ± 0.04 ± 0.06	2.5k	¹ ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
0.28 ± 0.06		² GAISER	86	CBAL $e^+e^- \rightarrow \gamma X$

¹ATHAR 04 used $\Gamma_{\eta_c(1S)} = 24.8 \pm 4.9$ MeV to obtain this result.

²GAISER 86 used $\Gamma_{\eta_c(1S)} = 11.5 \pm 4.5$ MeV to obtain this result.

WEIGHTED AVERAGE
0.34±0.05 (Error scaled by 1.3)



$\Gamma(\gamma\eta_c(2S))/\Gamma_{\text{total}}$

Γ_{143}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
7 ± 2 ± 4		¹ ABLIKIM	12G	BES3 $\psi(2S) \rightarrow \gamma K^0 K \pi, K K \pi^0$
< 8	90	² CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K \bar{K} \pi$
< 20	90	ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
20–130	95	EDWARDS	82C	CBAL $e^+e^- \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ABLIKIM 12G reports $[\Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K\bar{K}\pi)] = (1.30 \pm 0.20 \pm 0.30) \times 10^{-5}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (1.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²CRONIN-HENNESSY 10 reports $[\Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K\bar{K}\pi)] < 14.5 \times 10^{-6}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = 1.9 \times 10^{-2}$. This measurement assumes $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$ Γ_{144}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.04 ± 0.22 OUR AVERAGE Error includes scale factor of 1.4.					
0.95 ± 0.16 ± 0.05		423	ABLIKIM	17X BES3	$\psi(2S) \rightarrow \gamma\pi^0$
1.58 ± 0.40 ± 0.13		37	ABLIKIM	10F BES3	$\psi(2S) \rightarrow \gamma\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 5	90		PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$
< 5400	95		¹ LIBERMAN	75 SPEC	e^+e^-
< 1×10^4	90		WIJK	75 DASP	e^+e^-

¹ Restated by us using $B(\psi(2S) \rightarrow \mu^+\mu^-) = 0.0077$.

$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$ Γ_{145}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.24 ± 0.04 OUR AVERAGE					
1.251 ± 0.022 ± 0.062		56K	ABLIKIM	17X BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta, \gamma\pi^0\pi^0\eta$
1.26 ± 0.03 ± 0.08		2226	¹ ABLIKIM	10F BES3	$\psi(2S) \rightarrow 3\gamma\pi^+\pi^-, 2\gamma\pi^+\pi^-$
1.19 ± 0.08 ± 0.03			PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$
1.24 ± 0.27 ± 0.15		23	ABLIKIM	06R BES2	$e^+e^- \rightarrow \psi(2S)$
1.54 ± 0.31 ± 0.20		~ 43	BAI	98F BES	$\psi(2S) \rightarrow \pi^+\pi^-2\gamma, \pi^+\pi^-3\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 60	90		² BRAUNSCH...	77 DASP	e^+e^-
< 11	90		³ BARTEL	76 CNTR	e^+e^-

¹ Combining the results from $\eta' \rightarrow \pi^+\pi^-\eta$ and $\eta' \rightarrow \pi^+\pi^-\gamma$ decay modes.

² Restated by us using total decay width 228 keV.

³ The value is normalized to the branching ratio for $\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$.

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$ Γ_{146}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.73^{+0.29}_{-0.25} OUR AVERAGE Error includes scale factor of 1.8.					
2.84 ± 0.15 ^{+0.03} _{-0.10}		1.9k	1,2 DOBBS	15	$\psi(2S) \rightarrow \gamma\pi\pi$
2.12 ± 0.19 ± 0.32			3,4 BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.08 ± 0.19 ± 0.33		200.6 ± 18.8	³ BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
2.90 ± 1.08 ± 1.07		29.9 ± 11.1	³ BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^0\pi^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (2.39 \pm 0.09 \pm 0.09) \times 10^{-4}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

⁴ Combining the results from $\pi^+\pi^-$ and $\pi^0\pi^0$ decay modes.

$\Gamma(\gamma f_0(1370) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{147}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.1±1.0±1.4	175	¹ DOBBS	15	$\psi(2S) \rightarrow \gamma K \bar{K}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}}$ Γ_{148}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
9.2±1.8±0.6	274	^{1,2} DOBBS	15	$\psi(2S) \rightarrow \gamma \pi \pi$

¹ DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f_0(1500))/\Gamma_{\text{total}}] \times [B(f_0(1500) \rightarrow \pi \pi)] = (3.2 \pm 0.6 \pm 0.2) \times 10^{-5}$ which we divide by our best value $B(f_0(1500) \rightarrow \pi \pi) = (34.9 \pm 2.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using CLEO-c data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$ Γ_{149}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.3±0.8±0.1	136	^{1,2} DOBBS	15	$\psi(2S) \rightarrow \gamma K \bar{K}$

¹ DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f'_2(1525))/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K \bar{K})] = (2.9 \pm 0.6 \pm 0.3) \times 10^{-5}$ which we divide by our best value $B(f'_2(1525) \rightarrow K \bar{K}) = (88.7 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using CLEO-c data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{151}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.5 ±0.6 OUR AVERAGE				
3.6 ±0.4 ±0.5	290	¹ DOBBS	15	$\psi(2S) \rightarrow \gamma \pi \pi$
3.01±0.41±1.24	35.6 ± 4.8	² BAI	03C BES	$\psi(2S) \rightarrow \gamma \pi^+ \pi^-$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{152}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
6.6 ±0.7 OUR AVERAGE					

6.7 ±0.6 ±0.6		375	¹ DOBBS	15	$\psi(2S) \rightarrow \gamma K \bar{K}$
6.04±0.90±1.32	39.6 ± 5.9		^{2,3} BAI	03C BES	$\psi(2S) \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 15.6	90	6.8 ± 3.1	^{2,3} BAI	03C BES	$\psi(2S) \rightarrow \gamma K_S^0 K_S^0$
--------	----	-----------	--------------------	---------	---

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Includes unknown branching fractions to $K^+ K^-$ or $K_S^0 K_S^0$. We have multiplied the $K^+ K^-$ result by a factor of 2 and the $K_S^0 K_S^0$ result by a factor of 4 to obtain the $K \bar{K}$ result.

³ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

 $\Gamma(\gamma f_0(2100) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{153}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
4.8±0.5±0.9	373	¹ DOBBS	15	$\psi(2S) \rightarrow \gamma \pi \pi$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{154}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.2 \pm 0.6 \pm 0.8$	207	¹ DOBBS	15	$\psi(2S) \rightarrow \gamma K \bar{K}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{155}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5.8 \times 10^{-6}$	90	^{1,2} DOBBS	15	$\psi(2S) \rightarrow \gamma \pi \pi$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $\pi^+ \pi^-$ and $\pi^0 \pi^0$ are $3.2/4.3 \times 10^{-6}$ and $2.6/4.0 \times 10^{-6}$, respectively.

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{156}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 9.5 \times 10^{-6}$	90	^{1,2} DOBBS	15	$\psi(2S) \rightarrow \gamma K \bar{K}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $K^+ K^-$ and $K_S^0 K_S^0$ are $2.1/4.3 \times 10^{-6}$ and $3.7/5.5 \times 10^{-6}$, respectively.

 $\Gamma(\gamma \eta)/\Gamma_{\text{total}}$ Γ_{158}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.92 ± 0.18 OUR AVERAGE					
$0.85 \pm 0.18 \pm 0.04$		382	¹ ABLIKIM	17X BES3	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0$, $\gamma 3\pi^0$
$1.38 \pm 0.48 \pm 0.09$		13	¹ ABLIKIM	10F BES3	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0$, $\gamma 3\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2	90	PEDLAR	09	CLE3	$\psi(2S) \rightarrow \gamma X$
< 90	90	BAI	98F	BES	$\psi(2S) \rightarrow \pi^+ \pi^- 3\gamma$
< 200	90	YAMADA	77	DASP	$e^+ e^- \rightarrow 3\gamma$

¹ Combining the results from $\eta \rightarrow \pi^+ \pi^- \pi^0$ and $\eta \rightarrow 3\pi^0$ decay modes.

 $\Gamma(\gamma \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{159}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$8.71 \pm 1.25 \pm 1.64$	418	ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

 $\Gamma(\gamma \eta(1405) \rightarrow \gamma K \bar{K} \pi)/\Gamma_{\text{total}}$ Γ_{161}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.9	90	ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.3	90	ABLIKIM	06R	BES2	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$
< 1.2	90	¹ SCHARRE	80	MRK1	$e^+ e^-$

¹ Includes unknown branching fraction $\eta(1405) \rightarrow K \bar{K} \pi$.

 $\Gamma(\gamma \eta(1405) \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{162}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.36 \pm 0.25 \pm 0.05$	10	ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

$\Gamma(\gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{163}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.0 \times 10^{-7}$	90	ABLIKIM	17AJ BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$

 $\Gamma(\gamma\eta(1475) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}}$ Γ_{165}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.4	90	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.5	90	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$
--------	----	---------	----------	---

 $\Gamma(\gamma\eta(1475) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{166}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.88	90	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

 $\Gamma(\gamma 2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{167}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$39.6 \pm 2.8 \pm 5.0$	583	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma K^{*0} K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{168}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$37.0 \pm 6.1 \pm 7.2$	237	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}}$ Γ_{169}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$24.0 \pm 4.5 \pm 5.0$	41	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{170}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$25.6 \pm 3.6 \pm 3.6$	115	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{171}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$19.1 \pm 2.7 \pm 4.3$	132	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{172}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.9 ± 0.5 OUR AVERAGE				Error includes scale factor of 2.0.
$4.18 \pm 0.26 \pm 0.18$	348	¹ ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$
$2.9 \pm 0.4 \pm 0.4$	142	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

$\Gamma(\gamma f_2(1950) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{173}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.2 \pm 0.2 \pm 0.1$	111	¹ ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \gamma p \bar{p}$

¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

 $\Gamma(\gamma f_2(2150) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{174}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.72 \pm 0.18 \pm 0.03$	73	¹ ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \gamma p \bar{p}$

¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

 $\Gamma(\gamma X(1835) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{175}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$4.57 \pm 0.36 +1.77 -4.26$		ABLIKIM 12D	BES3	$J/\psi \rightarrow \gamma p \bar{p}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.6	90	ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \gamma p \bar{p}$
<5.4	90	ABLIKIM 07D	BES	$\psi(2S) \rightarrow \gamma p \bar{p}$

 $\Gamma(\gamma X \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{176}/Γ

For a narrow resonance in the range $2.2 < M(X) < 2.8$ GeV.

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2	90	ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \gamma p \bar{p}$

 $\Gamma(\gamma \pi^+ \pi^- p \bar{p})/\Gamma_{\text{total}}$ Γ_{177}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.8 \pm 1.2 \pm 0.7$	17	ABLIKIM 07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma 2(\pi^+ \pi^-) K^+ K^-)/\Gamma_{\text{total}}$ Γ_{178}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<22	90	ABLIKIM 07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma 3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{179}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<17	90	ABLIKIM 07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

 $\Gamma(\gamma K^+ K^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{180}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4	90	ABLIKIM 07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

$$\Gamma(\gamma\gamma J/\psi)/\Gamma_{\text{total}} \qquad \Gamma_{181}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.1 \pm 0.6^{+0.8}_{-1.0}$	1.1k	ABLIKIM	120	BES3 $e^+e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.2 ± 0.6	1.1k	¹ ABLIKIM	17N	BES3 $\psi(2S) \rightarrow \gamma\gamma J/\psi$
---------------	------	----------------------	-----	---

¹ Uses $B(J/\psi \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$ and $B(J/\psi \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033)\%$. No systematic error estimation.

$$\Gamma(e^+e^- \chi_{c0}(1P))/\Gamma_{\text{total}} \qquad \Gamma_{182}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$10.6 \pm 2.4 \pm 0.4$	48	¹ ABLIKIM	17I	BES3 $\psi(2S) \rightarrow e^+e^- \gamma J/\psi$

¹ ABLIKIM 17I reports $(11.7 \pm 2.5 \pm 1.0) \times 10^{-4}$ from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+e^- \chi_{c0}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (1.27 \pm 0.06) \times 10^{-2}$, which we rescale to our best value $B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (1.40 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(e^+e^- \chi_{c1}(1P))/\Gamma_{\text{total}} \qquad \Gamma_{183}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$8.5 \pm 0.6 \pm 0.2$	873	¹ ABLIKIM	17I	BES3 $\psi(2S) \rightarrow e^+e^- \gamma J/\psi$

¹ ABLIKIM 17I reports $(8.6 \pm 0.3 \pm 0.6) \times 10^{-4}$ from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+e^- \chi_{c1}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (33.9 \pm 1.2) \times 10^{-2}$, which we rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.3 \pm 1.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(e^+e^- \chi_{c2}(1P))/\Gamma_{\text{total}} \qquad \Gamma_{184}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$7.0 \pm 0.7 \pm 0.2$	227	¹ ABLIKIM	17I	BES3 $\psi(2S) \rightarrow e^+e^- \gamma J/\psi$

¹ ABLIKIM 17I reports $(6.9 \pm 0.5 \pm 0.6) \times 10^{-4}$ from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+e^- \chi_{c2}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (19.2 \pm 0.7) \times 10^{-2}$, which we rescale to our best value $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (19.0 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(e^+e^- \chi_{c0}(1P))/\Gamma(\gamma\chi_{c0}(1P)) \qquad \Gamma_{182}/\Gamma_{139}$$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$9.4 \pm 1.9 \pm 0.6$	48	¹ ABLIKIM	17I	BES3 $\psi(2S) \rightarrow e^+e^- \gamma J/\psi$

¹ Uses $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) \times B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (15.8 \pm 0.3 \pm 0.6) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.

$\Gamma(e^+e^-\chi_{c1}(1P))/\Gamma(\gamma\chi_{c1}(1P))$ $\Gamma_{183}/\Gamma_{140}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$8.3 \pm 0.3 \pm 0.4$	873	¹ ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+e^-\gamma J/\psi$

¹ Uses $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) \times B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (351.8 \pm 1.0 \pm 12.0) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.

$\Gamma(e^+e^-\chi_{c2}(1P))/\Gamma(\gamma\chi_{c2}(1P))$ $\Gamma_{184}/\Gamma_{141}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.6 \pm 0.5 \pm 0.4$	227	¹ ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+e^-\gamma J/\psi$

¹ Uses $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) \times B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (199.6 \pm 0.8 \pm 7.0) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.

————— WEAK DECAYS —————

$\Gamma(D^0 e^+e^- + c.c.)/\Gamma_{total}$ Γ_{185}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.4 \times 10^{-7}$	90	¹ ABLIKIM	17AF BES3	$e^+e^- \rightarrow \psi(2S)$

¹ Using D^0 decays to $K^-\pi^+$, $K^-\pi^+\pi^0$, and $K^-\pi^+\pi^+\pi^-$.

————— OTHER DECAYS —————

$\Gamma(\text{invisible})/\Gamma(e^+e^-)$ Γ_{186}/Γ_6

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 2.0	90	LEES	13I BABR	$B \rightarrow K^{(*)}\psi(2S)$

$\psi(2S)$ CROSS-PARTICLE BRANCHING RATIOS

For measurements involving $B(\psi(2S) \rightarrow \gamma\chi_{cJ}(1P)) \times B(\chi_{cJ}(1P) \rightarrow X)$ see the corresponding entries in the $\chi_{cJ}(1P)$ sections.

MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS

$\psi(2S) \rightarrow \gamma\chi_{cJ}(1P)$ and $\chi_{cJ} \rightarrow \gamma J/\psi(1S)$

$a_2(\chi_{c1})/a_2(\chi_{c2})$ Magnetic quadrupole transition amplitude ratio

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
63 ± 7 OUR AVERAGE				
61.7 ± 8.3	253k	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
67^{+19}_{-13}	59k	² ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

¹ Statistical and systematic errors combined.

² Statistical and systematic errors combined. Using values from fits with floating $M2$ amplitudes $a_2(\chi_{c1})$, $a_2(\chi_{c2})$, $b_2(\chi_{c1})$, $b_2(\chi_{c2})$ and fixed $E3$ amplitudes of $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$. Not independent of values for $a_2(\chi_{c1}(1P))$ and $a_2(\chi_{c2}(1P))$ from ARTUSO 09.

$b_2(\chi_{c2})/b_2(\chi_{c1})$ Magnetic quadrupole transition amplitude ratio

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
60±31 OUR AVERAGE				
74±40	253k	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
37 ⁺⁵³ ₋₄₇	59k	² ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

¹ Statistical and systematic errors combined. Derived from the reported measurement of $b_2(\chi_{c1})/b_2(\chi_{c2}) = 1.35 \pm 0.72$.

² Statistical and systematic errors combined. Using values from fits with floating $M2$ amplitudes $a_2(\chi_{c1})$, $a_2(\chi_{c2})$, $b_2(\chi_{c1})$, $b_2(\chi_{c2})$ and fixed $E3$ amplitudes of $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$. Not independent of values for $b_2(\chi_{c1}(1P))$ and $b_2(\chi_{c2}(1P))$ from ARTUSO 09.

 $\psi(2S)$ REFERENCES

ABLIKIM	17AF	PR D96 111101	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17AJ	PR D96 112008	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17AK	PR D96 112012	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17E	PL B770 217	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17I	PRL 118 221802	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17L	PR D95 052003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17U	PR D96 032001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17X	PR D96 052003	M. Ablikim <i>et al.</i>	(BES III Collab.)
DOBBS	17	PR D96 092004	S. Dobbs <i>et al.</i>	(NWES, WAYN)
LEES	17A	PR D95 052001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AAIJ	16Y	JHEP 1605 132	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	16L	PR D93 072003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15I	PR D91 092006	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15V	PL B749 414	M. Ablikim <i>et al.</i>	(BES III Collab.)
ANASHIN	15	PL B749 50	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
LEES	15J	PR D92 072008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	14G	PR D89 112006	M. Ablikim <i>et al.</i>	(BES III Collab.)
DOBBS	14	PL B739 90	S. Dobbs <i>et al.</i>	(NWES, WAYN)
ABLIKIM	13A	PRL 110 022001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13F	PR D87 052007	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13M	PR D87 092006	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13R	PR D88 032007	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13S	PR D88 032010	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13W	PR D88 112007	M. Ablikim <i>et al.</i>	(BES III Collab.)
LEES	13I	PR D87 112005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13O	PR D87 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Y	PR D88 072009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AAIJ	12H	EPJ C72 1972	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12G	PRL 109 042003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12H	PL B710 594	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12L	PR D86 072011	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12M	PR D86 092008	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12Q	CP C36 1040	M. Ablikim <i>et al.</i>	(BES II Collab.)
ANASHIN	12	PL B711 280	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
LEES	12E	PR D85 112009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
METREVELI	12	PR D85 092007	Z. Metreveli <i>et al.</i>	(NWES, FLOR, WAYN+)
GE	11	PR D84 032008	J.Y. Ge <i>et al.</i>	(CLEO Collab.)
ABLIKIM	10B	PRL 104 132002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	10F	PRL 105 261801	M. Ablikim <i>et al.</i>	(BES III Collab.)
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
CRONIN-HEN...	10	PR D81 052002	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)
ADAMS	09	PR D80 051106	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ARTUSO	09	PR D80 112003	M. Artuso <i>et al.</i>	(CLEO Collab.)

LIBBY	09	PR D80 072002	J. Libby <i>et al.</i>	(CLEO Collab.)
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
ABLIKIM	08B	PL B659 74	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)
DOBBS	08A	PRL 101 182003	S. Dobbs <i>et al.</i>	(CLEO Collab.)
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
ABLIKIM	07C	PL B648 149	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07D	PRL 99 011802	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ANASHIN	07	JETPL 85 347	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
		Translated from ZETFP 85 429.		
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Femilab E835 Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		PR D77 119902E (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)
PDG	07	Unofficial 2007 WWW edition		(PDG Collab.)
PEDLAR	07	PR D75 011102	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
ABLIKIM	06G	PR D73 052004	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06L	PRL 97 121801	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06W	PR D74 112003	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	06	PRL 96 082004	N.E. Adam <i>et al.</i>	(CLEO Collab.)
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT, BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)
DOBBS	06A	PR D74 011105	S. Dobbs <i>et al.</i>	(CLEO Collab.)
ABLIKIM	05E	PR D71 072006	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05I	PL B614 37	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05J	PL B619 247	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	05	PRL 94 012005	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ANDREOTTI	05	PR D71 032006	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)
BRIERE	05	PRL 95 062001	R.A. Briere <i>et al.</i>	(CLEO Collab.)
PEDLAR	05	PR D72 051108	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
ROSNER	05	PRL 95 102003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04K	PR D70 112003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04L	PR D70 112007	M. Ablikim <i>et al.</i>	(BES Collab.)
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
BAI	04B	PRL 92 052001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04C	PR D69 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
SETH	04	PR D69 097503	K.K. Seth	
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
BAI	03B	PR D67 052002	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)
AUBERT	02B	PR D65 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)
BAI	02	PR D65 052004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	02B	PL B550 24	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
PDG	02	PR D66 010001	K. Hagiwara <i>et al.</i>	(PDG Collab.)
BAI	01	PR D63 032002	J.Z. Bai <i>et al.</i>	(BES Collab.)
AMBROGIANI	00A	PR D62 032004	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98E	PR D57 3854	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98F	PR D58 097101	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98J	PRL 81 5080	J.Z. Bai <i>et al.</i>	(BES Collab.)
ARMSTRONG	97	PR D55 1153	T.A. Armstrong <i>et al.</i>	(E760 Collab.)
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 Collab., E706 Collab.)

ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
FRANKLIN	83	PRL 51 963	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
HIMEL	80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, SLAC)
OREGLIA	80	PRL 45 959	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)
BRANDELIK	79C	ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)
BRAUNSCH...	77	PL 67B 249	W. Braunschweig <i>et al.</i>	(DASP Collab.)
BURMESTER	77	PL 66B 395	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
YAMADA	77	Hamburg Conf. 69	S. Yamada	(DASP Collab.)
BARTEL	76	PL 64B 483	W. Bartel <i>et al.</i>	(DESY, HEIDP)
TANENBAUM	76	PRL 36 402	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL) IG
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)
ABRAMS	75	Stanford Symp. 25	G.S. Abrams	(LBL)
ABRAMS	75B	PRL 34 1181	G.S. Abrams <i>et al.</i>	(LBL, SLAC)
BOYARSKI	75C	Palermo Conf. 54	A.M. Boyarski <i>et al.</i>	(SLAC, LBL)
HILGER	75	PRL 35 625	E. Hilger <i>et al.</i>	(STAN, PENN)
LIBERMAN	75	Stanford Symp. 55	A.D. Liberman	(STAN)
LUTH	75	PRL 35 1124	V. Luth <i>et al.</i>	(SLAC, LBL) JPC
WIIK	75	Stanford Symp. 69	B.H. Wiik	(DESY)