

$$I(J^P) = 0(0^-)$$

I, J, P need confirmation. Quantum numbers shown are quark-model predictions.

B_s^0 MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
5366.88 ± 0.17 OUR FIT				
5366.84 ± 0.21 OUR AVERAGE				
5366.83 ± 0.25 ± 0.27		1 AAIJ	18AC LHCb	$p\bar{p}$ at 7, 8, 13 TeV
5367.08 ± 0.38 ± 0.15	128	2 AAIJ	16U LHCb	$p\bar{p}$ at 7, 8 TeV
5366.90 ± 0.28 ± 0.23		3 AAIJ	12E LHCb	$p\bar{p}$ at 7 TeV
5364.4 ± 1.3 ± 0.7		LOUVOT	09 BELL	$e^+e^- \rightarrow \gamma(5S)$
5366.01 ± 0.73 ± 0.33		4 ACOSTA	06 CDF	$p\bar{p}$ at 1.96 TeV
5369.9 ± 2.3 ± 1.3	32	5 ABE	96B CDF	$p\bar{p}$ at 1.8 TeV
5374 ± 16 ± 2	3	ABREU	94D DLPH	$e^+e^- \rightarrow Z$
5359 ± 19 ± 7	1	5 AKERS	94J OPAL	$e^+e^- \rightarrow Z$
5368.6 ± 5.6 ± 1.5	2	BUSKULIC	93G ALEP	$e^+e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5370 ± 1 ± 3		DRUTSKOY	07A BELL	Repl. by LOUVOT 09
5370 ± 40	6	6 AKERS	94J OPAL	$e^+e^- \rightarrow Z$
5383.3 ± 4.5 ± 5.0	14	ABE	93F CDF	Repl. by ABE 96B

¹ Uses $B_s \rightarrow \chi_{c1} K^+ K^-$ mode.

² Uses $J/\psi \rightarrow \mu^+ \mu^-$, $\phi \rightarrow K^+ K^-$ decays, and observes 128 ± 13 events of $B_s^0 \rightarrow J/\psi \phi \phi$.

³ Uses $B_s^0 \rightarrow J/\psi \phi$ fully reconstructed decays.

⁴ Uses exclusively reconstructed final states containing a $J/\psi \rightarrow \mu^+ \mu^-$ decays.

⁵ From the decay $B_s \rightarrow J/\psi(1S)\phi$.

⁶ From the decay $B_s \rightarrow D_s^- \pi^+$.

$m_{B_s^0} - m_B$

m_B is the average of our B masses ($m_{B^\pm} + m_{B^0}$)/2.

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
87.40 ± 0.18 OUR FIT				
87.42 ± 0.24 OUR AVERAGE				
87.60 ± 0.44 ± 0.09		1 AAIJ	15U LHCb	$p\bar{p}$ at 7, 8 TeV
87.42 ± 0.30 ± 0.09		2 AAIJ	12E LHCb	$p\bar{p}$ at 7 TeV
86.64 ± 0.80 ± 0.08		3 ACOSTA	06 CDF	$p\bar{p}$ at 1.96 TeV
• • • We use the following data for averages but not for fits. • • •				
89.7 ± 2.7 ± 1.2		ABE	96B CDF	$p\bar{p}$ at 1.8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
80 to 130	68	LEE-FRANZINI 90	CSB2	$e^+e^- \rightarrow \gamma(5S)$

¹ The reported result is $m_{B_s^0} - m_{B^0} = 87.45 \pm 0.44 \pm 0.09$ MeV. We convert it to the mass difference with respect to the average of $(m_{B^\pm} + m_{B^0})/2$. Uses the mode $B_s^0 \rightarrow \psi(2S) K^- \pi^+$.

² The reported result is $m_{B_s^0} - m_{B^+} = 87.52 \pm 0.30 \pm 0.12$ MeV. We convert it to the mass difference with respect to the average of $(m_{B^\pm} + m_{B^0})/2$.

³ The reported result is $m_{B_s^0} - m_{B^0} = 86.38 \pm 0.90 \pm 0.06$ MeV. We convert it to the mass difference with respect to the average of $(m_{B^\pm} + m_{B^0})/2$.

$m_{B_{sH}^0} - m_{B_{sL}^0}$

See the B_s^0 - \bar{B}_s^0 MIXING section near the end of these B_s^0 Listings.

B_s^0 MEAN LIFE

"OUR EVALUATION" is provided by the Heavy Flavor Averaging Group (HFLAV). It is derived from the average of $\Gamma_{B_s^0}$.

VALUE (10^{-12} s)	EVTS	DOCUMENT ID	TECN	COMMENT
1.510±0.004 OUR EVALUATION				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.518±0.041±0.027		¹ AALTONEN	11AP CDF	$p\bar{p}$ at 1.96 TeV
1.398±0.044 ^{+0.028} _{-0.025}		² ABAZOV	06V D0	$p\bar{p}$ at 1.96 TeV
1.42 ^{+0.14} _{-0.13} ±0.03		³ ABREU	00Y DLPH	$e^+ e^- \rightarrow Z$
1.53 ^{+0.16} _{-0.15} ±0.07		⁴ ABREU,P	00G DLPH	$e^+ e^- \rightarrow Z$
1.36 ±0.09 ^{+0.06} _{-0.05}		⁵ ABE	99D CDF	$p\bar{p}$ at 1.8 TeV
1.72 ^{+0.20} _{-0.19} ^{+0.18} _{-0.17}		⁶ ACKERSTAFF	98F OPAL	$e^+ e^- \rightarrow Z$
1.50 ^{+0.16} _{-0.15} ±0.04		⁵ ACKERSTAFF	98G OPAL	$e^+ e^- \rightarrow Z$
1.47 ±0.14 ±0.08		⁴ BARATE	98C ALEP	$e^+ e^- \rightarrow Z$
1.51 ±0.11		⁷ BARATE	98C ALEP	$e^+ e^- \rightarrow Z$
1.56 ^{+0.29} _{-0.26} ^{+0.08} _{-0.07}		⁵ ABREU	96F DLPH	Repl. by ABREU 00Y
1.65 ^{+0.34} _{-0.31} ±0.12		⁴ ABREU	96F DLPH	Repl. by ABREU 00Y
1.76 ±0.20 ^{+0.15} _{-0.10}		⁸ ABREU	96F DLPH	Repl. by ABREU 00Y
1.60 ±0.26 ^{+0.13} _{-0.15}		⁹ ABREU	96F DLPH	Repl. by ABREU,P 00G
1.67 ±0.14		¹⁰ ABREU	96F DLPH	$e^+ e^- \rightarrow Z$
1.61 ^{+0.30} _{-0.29} ^{+0.18} _{-0.16}	90	⁴ BUSKULIC	96E ALEP	Repl. by BARATE 98C
1.54 ^{+0.14} _{-0.13} ±0.04		⁵ BUSKULIC	96M ALEP	$e^+ e^- \rightarrow Z$
1.42 ^{+0.27} _{-0.23} ±0.11	76	⁵ ABE	95R CDF	Repl. by ABE 99D

1.74	$\begin{array}{c} +1.08 \\ -0.69 \end{array}$	± 0.07	8	11 ABE	95R	CDF	Sup. by ABE 96N
1.54	$\begin{array}{c} +0.25 \\ -0.21 \end{array}$	± 0.06	79	5 AKERS	95G	OPAL	Repl. by ACKER-STAFF 98G
1.59	$\begin{array}{c} +0.17 \\ -0.15 \end{array}$	± 0.03	134	5 BUSKULIC	95O	ALEP	Sup. by BUSKULIC 96M
0.96	± 0.37		41	12 ABREU	94E	DLPH	Sup. by ABREU 96F
1.92	$\begin{array}{c} +0.45 \\ -0.35 \end{array}$	± 0.04	31	5 BUSKULIC	94C	ALEP	Sup. by BUSKULIC 95O
1.13	$\begin{array}{c} +0.35 \\ -0.26 \end{array}$	± 0.09	22	5 ACTON	93H	OPAL	Sup. by AKERS 95G

¹ AALTONEN 11AP combines the fully reconstructed $B_s^0 \rightarrow D_s^- \pi^+$ decays and partially reconstructed $B_s^0 \rightarrow D_s^- X$ decays.

² Measured using $D_s \mu^+$ vertices.

³ Uses $D_s^- \ell^+$, and $\phi \ell^+$ vertices.

⁴ Measured using D_s hadron vertices.

⁵ Measured using $D_s^- \ell^+$ vertices.

⁶ ACKERSTAFF 98F use fully reconstructed $D_s^- \rightarrow \phi \pi^-$ and $D_s^- \rightarrow K^{*0} K^-$ in the inclusive B_s^0 decay.

⁷ Combined results from $D_s^- \ell^+$ and D_s hadron.

⁸ Measured using $\phi \ell$ vertices.

⁹ Measured using inclusive D_s vertices.

¹⁰ Combined result for the four ABREU 96F methods.

¹¹ Exclusive reconstruction of $B_s \rightarrow \psi \phi$.

¹² ABREU 94E uses the flight-distance distribution of D_s vertices, ϕ -lepton vertices, and $D_s \mu$ vertices.

$\Gamma_{B_s^0}$

"OUR EVALUATION" is an average performed by the Heavy Flavor Averaging Group (HFLAV) as described in our "Review on B - \bar{B} Mixing" in the B^0 section of these Listings. It includes the measurements of $\Gamma_{B_s^0}$ and $\Delta\Gamma_{B_s^0}$ listed in this section, as well as constraints from effective lifetimes with pure CP modes and flavor-specific modes.

VALUE (10^{12} s^{-1})	DOCUMENT ID	TECN	COMMENT
0.6624 \pm 0.0018 OUR EVALUATION			
0.664 \pm 0.004 OUR AVERAGE Error includes scale factor of 1.7. See the ideogram below.			
0.650 ± 0.006 ± 0.004	¹ AAIJ	17V LHCb	$p p$ at 7, 8 TeV
0.675 ± 0.003 ± 0.003	² AAD	16AP ATLS	$p p$ at 7, 8 TeV
0.668 ± 0.011 ± 0.006	³ AAIJ	16AK LHCb	$p p$ at 7, 8 TeV
0.6704 ± 0.0043 ± 0.0055	² KHACHATRYAN...16S	CMS	$p p$ at 8 TeV
0.6603 ± 0.0027 ± 0.0015	⁴ AAIJ	15I LHCb	$p p$ at 7, 8 TeV
0.654 ± 0.008 ± 0.004	² AALTENON	12AJ CDF	$p \bar{p}$ at 1.96 TeV
0.693 $\begin{array}{c} +0.018 \\ -0.017 \end{array}$	² ABAZOV	12D D0	$p \bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.677 ± 0.007	± 0.004	2 AAD	14U ATLS	Repl. by AAD 16AP
0.661 ± 0.004	± 0.006	5 AAIJ	13AR LHCb	Repl. by AAIJ 15I
0.677 ± 0.007	± 0.004	2 AAD	12CV ATLS	Repl. by AAD 14U
0.657 ± 0.009	± 0.008	2 AAIJ	12D LHCb	Repl. by AAIJ 13AR
0.654 ± 0.011	± 0.005	2,6 AALTONEN	12D CDF	Repl. by AALTONEN 12AJ
0.672 ± 0.027	± 0.013	2 ABAZOV	09E D0	Repl. by ABAZOV 08AM
0.658 ± 0.017	± 0.009	2,7 AALTONEN	08J CDF	Repl. by AALTONEN 12D
0.658 ± 0.022	± 0.004	2 ABAZOV	08AM D0	Repl. by ABAZOV 12D
0.658 ± 0.035	$+0.0130$ -0.004	2,7 ABAZOV	07 D0	Repl. by ABAZOV 09E
0.714 ± 0.007	± 0.010	2,7 ACOSTA	05 CDF	Repl. by AALTONEN 08J

1 Measured using time-dependent angular analysis of $B_s^0 \rightarrow J/\psi K^+ K^-$ in the region $m(KK) > 1.05$ GeV.

2 Measured using a time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays.


3 Measured using a time-dependent angular analysis of $B_s^0 \rightarrow \psi(2S)\phi$ decays.

4 Measured using a time-dependent angular analysis of $B_s^0 \rightarrow J/\psi K^+ K^-$ decays.

5 Measured using a combined time-dependent angular analysis of $B_s^0 \rightarrow J/\psi K^+ K^-$ and $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$ decays.

6 Assuming CPV phase $\phi_s = -0.04$.

7 Assuming CPV phase $\phi_s = 0$.



$\Delta\Gamma_{B_s^0}$

"OUR EVALUATION" is an average performed by the Heavy Flavor Averaging Group (HFLAV) as described in our "Review on B - \bar{B} Mixing" in the B^0 section of these Listings. It includes the measurements of $\Gamma_{B_s^0}$ and $\Delta\Gamma_{B_s^0}$ listed in this section, as well as constraints from effective lifetimes with pure CP modes and flavor-specific modes.

VALUE (10^{12} s^{-1})	DOCUMENT ID	TECN	COMMENT
0.090 ± 0.005 OUR EVALUATION			
0.083 ± 0.006 OUR AVERAGE			
0.066 ± 0.018 ± 0.010	1 AAIJ	17V LHCb	$p\bar{p}$ at 7, 8 TeV
0.085 ± 0.011 ± 0.007	2 AAD	16AP ATLAS	$p\bar{p}$ at 7, 8 TeV
0.066 +0.041 -0.044 ± 0.007	3 AAIJ	16AK LHCb	$p\bar{p}$ at 7, 8 TeV
0.095 ± 0.013 ± 0.007	2 KHACHATRYAN...16S CMS	CMS	$p\bar{p}$ at 8 TeV
0.0805 ± 0.0091 ± 0.0032	4 AAIJ	15I LHCb	$p\bar{p}$ at 7, 8 TeV
0.068 ± 0.026 ± 0.009	2 AALTONEN	12AJ CDF	$p\bar{p}$ at 1.96 TeV
0.163 +0.065 -0.064	2,5 ABAZOV	12D D0	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.053 ± 0.021 ± 0.010	2 AAD	14U ATLAS	Repl. by AAD 16AP
0.106 ± 0.011 ± 0.007	6 AAIJ	13AR LHCb	Repl. by AAIJ 15I
0.053 ± 0.021 ± 0.010	2 AAD	12CV ATLAS	Repl. by AAD 14U
0.123 ± 0.029 ± 0.011	2 AAIJ	12D LHCb	Repl. by AAIJ 13AR
0.075 ± 0.035 ± 0.006	7 AALTONEN	12D CDF	Repl. by AALTONEN 12AJ
0.085 +0.072 -0.078 ± 0.001	8 ABAZOV	09E D0	Repl. by ABAZOV 08AM
0.076 +0.059 -0.063 ± 0.006	9 AALTONEN	08J CDF	Repl. by AALTONEN 12D
0.19 ± 0.07 +0.02 -0.01	2,10 ABAZOV	08AM D0	Repl. by ABAZOV 12D
0.12 +0.08 -0.10 ± 0.02	9,11 ABAZOV	07 D0	Repl. by ABAZOV 07N
0.13 ± 0.09	12 ABAZOV	07N D0	Repl. by ABAZOV 09E
0.47 +0.19 -0.24 ± 0.01	9 ACOSTA	05 CDF	Repl. by AALTONEN 08J

¹ Measured using time-dependent angular analysis of $B_s^0 \rightarrow J/\psi K^+ K^-$ in the region $m(KK) > 1.05$ GeV.

² Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays.

³ Measured using time-dependent angular analysis of $B_s^0 \rightarrow \psi(2S) \phi$ decays.

⁴ Measured using time-dependent angular analysis of $B_s^0 \rightarrow J/\psi K^+ K^-$ decays.

⁵ The error includes both statistical and systematic uncertainties.

⁶ AAIJ 13AR result comes from a combined fit to $B_s^0 \rightarrow J/\psi K^+ K^-$ and $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$ data sets. Also reports $\Delta\Gamma_s = 0.100 \pm 0.016 \pm 0.003 \text{ ps}^{-1}$ from a fit to $B_s^0 \rightarrow J/\psi K^+ K^-$ decays.

⁷ Uses the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays and assuming CP -violating angle $\beta_s(B_s^0 \rightarrow J/\psi \phi) = 0.02$.

- ⁸ Measured the angular and lifetime parameters for the time-dependent angular untagged decays $B_d^0 \rightarrow J/\psi K^{*0}$ and $B_s^0 \rightarrow J/\psi \phi$.
- ⁹ Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays and assuming CP -violating phase $\phi_s = 0$.
- ¹⁰ Obtains 90% CL interval $-0.06 < \Delta\Gamma_s < 0.30$.
- ¹¹ ABAZOV 07 reports $0.17 \pm 0.09 \pm 0.02$ with CP -violating phase ϕ_s as a free parameter.
- ¹² Combines D^0 measurements of time-dependent angular distributions in $B_s^0 \rightarrow J/\psi \phi$ and charge asymmetry in semileptonic decays. There is a 4-fold ambiguity in the solution.

$\Delta\Gamma_{B_s^0}/\Gamma_{B_s^0}$

$\Gamma_{B_s^0}$ and $\Delta\Gamma_{B_s^0}$ are the decay rate average and difference between two B_s^0 CP eigenstates (light – heavy).

"OUR EVALUATION" is provided by the Heavy Flavor Averaging Group (HFLAV). It is derived from the averages of $\Gamma_{B_s^0}$ and $\Delta\Gamma_{B_s^0}$ (and their correlation).

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.135±0.008 OUR EVALUATION				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.090±0.009±0.023		¹ ESEN ² AAIJ ³ AALTONEN ⁴ ABAZOV	13 BELL 12D LHCb 12D CDF 12D D0	$e^+ e^- \rightarrow \gamma(5S)$ $p\bar{p}$ at 7 TeV $p\bar{p}$ at 1.96 TeV $p\bar{p}$ at 1.96 TeV
$0.147^{+0.036+0.042}_{-0.030-0.041}$		¹ ESEN	10 BELL	$e^+ e^- \rightarrow \gamma(5S)$
$0.072 \pm 0.021 \pm 0.022$		⁵ ABAZOV	09I D0	$p\bar{p}$ at 1.96 TeV
>0.012	95	⁵ AALTONEN	08F CDF	$p\bar{p}$ at 1.96 TeV
$0.116^{+0.09}_{-0.10} \pm 0.010$		⁶ AALTONEN	08J CDF	Repl. by AALTONEN 12D
$0.079^{+0.038+0.031}_{-0.035-0.030}$		⁵ ABAZOV	07Y D0	Repl. by ABAZOV 09I
$0.24^{+0.28+0.03}_{-0.38-0.04}$		^{6,7} ABAZOV	05W D0	Repl. by ABAZOV 08AM
$0.65^{+0.25}_{-0.33} \pm 0.01$		⁶ ACOSTA	05 CDF	Repl. by AALTONEN 08J
<0.46	95	⁸ ABREU	00Y DLPH	$e^+ e^- \rightarrow Z$
<0.69	95	⁹ ABREU,P	00G DLPH	$e^+ e^- \rightarrow Z$
$0.25^{+0.21}_{-0.14}$		¹⁰ BARATE	00K ALEP	$e^+ e^- \rightarrow Z$
<0.83	95	¹¹ ABE	99D CDF	$p\bar{p}$ at 1.8 TeV
<0.67	95	¹² ACCIARRI	98S L3	$e^+ e^- \rightarrow Z$

¹ Assumes CP violation is negligible.

² Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays.

³ Uses the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays and assuming CP -violating angle $\beta_s(B_s^0 \rightarrow J/\psi \phi) = 0.02$.

⁴ Measured using fully reconstructed $B_s \rightarrow J/\psi \phi$ decays.

⁵ Assumes $2 B(B_s^0 \rightarrow D_s^{(*)} D_s^{(*)}) \simeq \Delta\Gamma_s^{CP} / \Gamma_s$.

⁶ Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays.

⁷ Uses $|A_0|^2 - |A_{||}|^2 = 0.355 \pm 0.066$ from ACOSTA 05.

⁸ Uses $D_s^- \ell^+$, and $\phi \ell^+$ vertices.

⁹ Measured using D_s hadron vertices.

¹⁰ Uses $\phi\phi$ correlations from $B_s^0 \rightarrow D_s^{(*)+} D_s^{(*)-}$.

¹¹ ABE 99D assumes $\tau_{B_s^0} = 1.55 \pm 0.05$ ps.

¹² ACCIARRI 98S assumes $\tau_{B_s^0} = 1.49 \pm 0.06$ ps and PDG 98 values of b production fraction.

B_{sH}^0 MEAN LIFE

B_{sH}^0 is the heavy mass state of two B_s^0 CP eigenstates.

"OUR EVALUATION" is provided by the Heavy Flavor Averaging Group (HFLAV). It is derived from the averages of $\Gamma_{B_s^0}$ and $\Delta\Gamma_{B_s^0}$ (and their correlation).

VALUE (10^{-12} s)	DOCUMENT ID	TECN	COMMENT
1.619±0.009 OUR EVALUATION			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.677±0.034±0.011	1 SIRUNYAN	18BY CMS	$p\bar{p}$ at 8 TeV
2.04 ± 0.44 ± 0.05	2 AAIJ	17AI LHCb	$p\bar{p}$ at 7, 8, 13 TeV
1.70 ± 0.14 ± 0.05	3 ABAZOV	16C D0	$p\bar{p}$ at 1.96 TeV
1.75 ± 0.12 ± 0.07	4 AAIJ	13AB LHCb	$p\bar{p}$ at 7 TeV
1.652±0.024±0.024	5 AAIJ	13AR LHCb	$p\bar{p}$ at 7 TeV
1.700±0.040±0.026	6 AAIJ	12AN LHCb	$p\bar{p}$ at 7 TeV
	7 AALTONEN	12D CDF	$p\bar{p}$ at 1.96 TeV
1.70 $^{+0.12}_{-0.11}$ ± 0.03	6 AALTONEN	11AB CDF	$p\bar{p}$ at 1.96 TeV
1.613 $^{+0.123}_{-0.113}$	8,9 AALTONEN	08J CDF	Repl. by AALTONEN 12D
1.58 $^{+0.39}_{-0.42}$ $^{+0.01}_{-0.02}$	9 ABAZOV	05W D0	Repl. by ABAZOV 08AM
2.07 $^{+0.58}_{-0.46}$ ± 0.03	9 ACOSTA	05 CDF	Repl. by AALTONEN 08J

¹ Measured using $B_s^0 \rightarrow J/\psi\pi^+\pi^-$ decays with $0.9240 < m(\pi\pi) < 1.0204$ GeV, which is dominated by the $f_0(980)$ resonance, making it a CP -odd state.

² Measured using $B_s \rightarrow \mu^+\mu^-$ decays which, in the Standard Model, correspond to B_{sH}^0 decays. Assumes $-2 \operatorname{Re}(\lambda)/(1 + |\lambda|^2) = 1$.

³ Measured using $J/\psi\pi^+\pi^-$ mode with $0.880 < m(\pi\pi) < 1.080$ GeV/c², which is mostly $J/\psi f(0)(980)$ mode, a pure CP -odd final state.

⁴ Measured using a pure CP -odd final state $J/\psi K_S^0$ with the assumption that contributions from penguin diagrams are small.

⁵ Measured using $B_s \rightarrow J/\psi\pi^+\pi^-$ decays which, in the limit of $\phi_s = 0$ and $|\lambda| = 1$, correspond to B_{sH}^0 decays.

⁶ Measured using a pure CP -odd final state $J/\psi f_0(980)$.

⁷ Uses the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays assuming CP -violating angle $\beta_s(B^0 \rightarrow J/\psi\phi) = 0.02$.

⁸ Obtained from $\Delta\Gamma_s$ and Γ_s fit with a correlation of 0.6.

⁹ Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays.

B_{sL}^0 MEAN LIFE

B_{sL}^0 is the light mass state of two B_s^0 CP eigenstates.

"OUR EVALUATION" is provided by the Heavy Flavor Averaging Group (HFLAV). It is derived from the averages of $\Gamma_{B_s^0}$ and $\Delta\Gamma_{B_s^0}$ (and their correlation).

VALUE (10^{-12} s)	DOCUMENT ID	TECN	COMMENT
1.414 ± 0.006 OUR EVALUATION			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.40 ± 0.02	¹ SIRUNYAN	18BY CMS	$p\bar{p}$ at 8 TeV
$1.479 \pm 0.034 \pm 0.011$	² AAIJ	16AL LHCb	$p\bar{p}$ at 7, 8 TeV
$1.379 \pm 0.026 \pm 0.017$	³ AAIJ	14F LHCb	$p\bar{p}$ at 7, 8 TeV
$1.407 \pm 0.016 \pm 0.007$	⁴ AAIJ	14R LHCb	$p\bar{p}$ at 7 TeV
$1.440 \pm 0.096 \pm 0.009$	⁴ AAIJ	12 LHCb	Repl. by AAIJ 14R
$1.455 \pm 0.046 \pm 0.006$	⁴ AAIJ	12R LHCb	Repl. by AAIJ 14R
	⁵ AALTENEN	12D CDF	$p\bar{p}$ at 1.96 TeV
$1.437^{+0.054}_{-0.047}$	^{6,7} AALTENEN	08J CDF	Repl. by AALTENEN 12D
$1.24^{+0.14}_{-0.11}{}^{+0.01}_{-0.02}$	⁷ ABAZOV	05W D0	Repl. by ABAZOV 08AM
$1.05^{+0.16}_{-0.13}{}^{\pm 0.02}$	⁷ ACOSTA	05 CDF	Repl. by AALTENEN 08J
$1.27 \pm 0.33 \pm 0.08$	⁸ BARATE	00K ALEP	$e^+e^- \rightarrow Z$

¹ Measured using results in SIRUNYAN 18BY for the heavy B_s^0 lifetime obtained from $B_s^0 \rightarrow J/\psi\pi^+\pi^-$ decays and the average effective $B_s^0 \rightarrow J/\psi\phi$ lifetime, and magnitude squared of the CP -odd amplitude $|A_\perp|^2 = 0.250 \pm 0.006$. The uncertainty includes all statistical and systematic contributions.

² Measured using $B_s^0 \rightarrow J/\psi\eta$ decays.

³ Measured using $B_s^0 \rightarrow D_s^- D_s^+$. The effective lifetime is translated into a decay width of $\Gamma_L = 0.725 \pm 0.014 \pm 0.009 \text{ ps}^{-1}$.

⁴ Measured using $B_s^0 \rightarrow K^+ K^-$ decays. There may still be CPV in the decay.

⁵ Uses the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays and assuming CP -violating angle $\beta_s(B_s^0 \rightarrow J/\psi\phi) = 0.02$.


⁶ Obtained from $\Delta\Gamma_s$ and Γ_s fit with a correlation of 0.6.

⁷ Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays.

⁸ Uses $\phi\phi$ correlations from $B_s^0 \rightarrow D_s^{(*)+} D_s^{(*)-}$.

B_s^0 MEAN LIFE (Flavor specific)

VALUE (10^{-12} s)	DOCUMENT ID	TECN	COMMENT
1.527±0.011 OUR EVALUATION			
1.526±0.015 OUR AVERAGE			
1.547±0.013±0.011	¹ AAIJ	17AN LHCb	$p\bar{p}$ at 7, 8 TeV
1.479±0.010±0.021	² ABAZOV	15A D0	$p\bar{p}$ at 1.96 TeV
1.535±0.015±0.014	³ AAIJ	14AX LHCb	$p\bar{p}$ at 7 TeV
1.52 ± 0.15 ± 0.01	⁴ AAIJ	14F LHCb	$p\bar{p}$ at 7, 8 TeV
1.60 ± 0.06 ± 0.01	⁵ AAIJ	14R LHCb	$p\bar{p}$ at 7 TeV
1.518±0.041±0.027	⁶ AALTONEN	11AP CDF	$p\bar{p}$ at 1.96 TeV
1.42 $^{+0.14}_{-0.13}$ ± 0.03	⁷ ABREU	00Y DLPH	$e^+e^- \rightarrow Z$
1.36 ± 0.09 $^{+0.06}_{-0.05}$	⁸ ABE	99D CDF	$p\bar{p}$ at 1.8 TeV
1.50 $^{+0.16}_{-0.15}$ ± 0.04	⁸ ACKERSTAFF	98G OPAL	$e^+e^- \rightarrow Z$
1.54 $^{+0.14}_{-0.13}$ ± 0.04	⁸ BUSKULIC	96M ALEP	$e^+e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.398±0.044 $^{+0.028}_{-0.025}$	⁹ ABAZOV	06v D0	Repl. by ABAZOV 15A



B_s^0 MEAN LIFE (Flavor specific) (10^{-12} s)

¹ AAIJ 17AN value was measured using $B_s^0 \rightarrow D_s^{(*)-} \mu^+ \nu_\mu$ decays relative to $B^0 \rightarrow D^{(*)-} \mu^+ \nu_\mu$ decays.

² Measured using $B_s^0 \rightarrow D_s^- \mu^+ \nu_\mu X$ decays.

³ Measured using the $B_s^0 \rightarrow D_s^- \pi^+$ decays.

⁴ Measured using $B_s^0 \rightarrow D^+ D_s^-$.⁵ Measured using $B_s^0 \rightarrow \pi^+ K^-$ decays.⁶ AALTONEN 11AP combines the fully reconstructed $B_s^0 \rightarrow D_s^- \pi^+$ decays and partially reconstructed $B_s^0 \rightarrow D_s X$ decays.⁷ Uses $D_s^- \ell^+$, and $\phi \ell^+$ vertices.⁸ Measured using $D_s^- \ell^+$ vertices.⁹ Measured using $D_s^- \mu^+$ vertices.

B_s^0 MEAN LIFE ($B_s \rightarrow J/\psi \phi$)

VALUE (10^{-12} s)	DOCUMENT ID	TECN	COMMENT
1.480 ± 0.007 OUR EVALUATION			
1.480 ± 0.007 OUR AVERAGE			
$1.481 \pm 0.007 \pm 0.005$	¹ SIRUNYAN	18BY CMS	$p\bar{p}$ at 8 TeV
$1.480 \pm 0.011 \pm 0.005$	¹ AAIJ	14E LHCb	$p\bar{p}$ at 7 TeV
$1.444^{+0.098}_{-0.090} \pm 0.020$	¹ ABAZOV	05B D0	$p\bar{p}$ at 1.96 TeV
$1.34^{+0.23}_{-0.19} \pm 0.05$	² ABE	98B CDF	$p\bar{p}$ at 1.8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.39^{+0.13}_{-0.16}^{+0.01}_{-0.02}$	² ABAZOV	05W D0	$p\bar{p}$ at 1.96 TeV
$1.34^{+0.23}_{-0.19} \pm 0.05$	³ ABE	96N CDF	Repl. by ABE 98B

¹ Measured using fully reconstructed $B_s \rightarrow J/\psi \phi$ decays.² Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays.³ ABE 96N uses 58 ± 12 exclusive $B_s \rightarrow J/\psi \phi$ events.

$\tau_{B_s^0}/\tau_{B^0}$ MEAN LIFE RATIO

 $\tau_{B_s^0}/\tau_{B^0}$ (direct measurements)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.980 \pm 0.006 \pm 0.003$	¹ SIRUNYAN	18BY CMS	$p\bar{p}$ at 8 TeV
¹ Measured using $B_s^0 \rightarrow J/\psi \phi(1020)$ and $B^0 \rightarrow J/\psi K^*(892)^0$ decays.			

B_s^0 DECAY MODES

These branching fractions all scale with $B(\bar{b} \rightarrow B_s^0)$.

The branching fraction $B(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell \text{anything})$ is not a pure measurement since the measured product branching fraction $B(\bar{b} \rightarrow B_s^0) \times B(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell \text{anything})$ was used to determine $B(\bar{b} \rightarrow B_s^0)$, as described in the note on “ B^0 - \bar{B}^0 Mixing”

For inclusive branching fractions, e.g., $B \rightarrow D^\pm \text{anything}$, the values usually are multiplicities, not branching fractions. They can be greater than one.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 D_s^- \text{anything}$	(93 \pm 25) %	
$\Gamma_2 \ell\nu_\ell X$	(9.6 \pm 0.8) %	
$\Gamma_3 e^+\nu X^-$	(9.1 \pm 0.8) %	
$\Gamma_4 \mu^+\nu X^-$	(10.2 \pm 1.0) %	
$\Gamma_5 D_s^- \ell^+ \nu_\ell \text{anything}$	[a] (8.1 \pm 1.3) %	
$\Gamma_6 D_s^{*-} \ell^+ \nu_\ell \text{anything}$	(5.4 \pm 1.1) %	
$\Gamma_7 D_{s1}(2536)^- \mu^+ \nu_\mu, D_{s1}^- \rightarrow D^{*-} K_S^0$	(2.6 \pm 0.7) $\times 10^{-3}$	
$\Gamma_8 D_{s1}(2536)^- X \mu^+ \nu, D_{s1}^- \rightarrow \overline{D}^0 K^+$	(4.4 \pm 1.3) $\times 10^{-3}$	
$\Gamma_9 D_{s2}(2573)^- X \mu^+ \nu, D_{s2}^- \rightarrow \overline{D}^0 K^+$	(2.7 \pm 1.0) $\times 10^{-3}$	
$\Gamma_{10} D_s^- \pi^+$	(3.00 \pm 0.23) $\times 10^{-3}$	
$\Gamma_{11} D_s^- \rho^+$	(6.9 \pm 1.4) $\times 10^{-3}$	
$\Gamma_{12} D_s^- \pi^+ \pi^+ \pi^-$	(6.1 \pm 1.0) $\times 10^{-3}$	
$\Gamma_{13} D_{s1}(2536)^- \pi^+, D_{s1}^- \rightarrow D_s^- \pi^+ \pi^-$	(2.5 \pm 0.8) $\times 10^{-5}$	
$\Gamma_{14} D_s^\mp K^\pm$	(2.27 \pm 0.19) $\times 10^{-4}$	
$\Gamma_{15} D_s^- K^+ \pi^+ \pi^-$	(3.2 \pm 0.6) $\times 10^{-4}$	
$\Gamma_{16} D_s^+ D_s^-$	(4.4 \pm 0.5) $\times 10^{-3}$	
$\Gamma_{17} D_s^- D_s^+$	(2.8 \pm 0.5) $\times 10^{-4}$	
$\Gamma_{18} D^+ D^-$	(2.2 \pm 0.6) $\times 10^{-4}$	
$\Gamma_{19} D^0 \overline{D}^0$	(1.9 \pm 0.5) $\times 10^{-4}$	
$\Gamma_{20} D_s^{*-} \pi^+$	(2.0 \pm 0.5) $\times 10^{-3}$	
$\Gamma_{21} D_s^{*\mp} K^\pm$	(1.33 \pm 0.35) $\times 10^{-4}$	
$\Gamma_{22} D_s^{*-} \rho^+$	(9.6 \pm 2.1) $\times 10^{-3}$	
$\Gamma_{23} D_s^{*+} D_s^- + D_s^{*-} D_s^+$	(1.37 \pm 0.16) %	
$\Gamma_{24} D_s^{*+} D_s^{*-}$	(1.44 \pm 0.20) %	S=1.1
$\Gamma_{25} D_s^{(*)+} D_s^{(*)-}$	(4.5 \pm 1.4) %	
$\Gamma_{26} \overline{D}^{*0} \overline{K}^0$	(2.8 \pm 1.1) $\times 10^{-4}$	
$\Gamma_{27} \overline{D}^0 \overline{K}^0$	(4.3 \pm 0.9) $\times 10^{-4}$	
$\Gamma_{28} \overline{D}^0 K^- \pi^+$	(1.04 \pm 0.13) $\times 10^{-3}$	
$\Gamma_{29} \overline{D}^0 \overline{K}^*(892)^0$	(4.4 \pm 0.6) $\times 10^{-4}$	
$\Gamma_{30} \overline{D}^0 \overline{K}^*(1410)$	(3.9 \pm 3.5) $\times 10^{-4}$	
$\Gamma_{31} \overline{D}^0 \overline{K}_0^*(1430)$	(3.0 \pm 0.7) $\times 10^{-4}$	
$\Gamma_{32} \overline{D}^0 \overline{K}_2^*(1430)$	(1.1 \pm 0.4) $\times 10^{-4}$	
$\Gamma_{33} \overline{D}^0 \overline{K}^*(1680)$	< 7.8 $\times 10^{-5}$	CL=90%
$\Gamma_{34} \overline{D}^0 \overline{K}_0^*(1950)$	< 1.1 $\times 10^{-4}$	CL=90%

Γ_{35}	$\overline{D}^0 \overline{K}_3^*(1780)$	$< 2.6 \times 10^{-5}$	CL=90%
Γ_{36}	$\overline{D}^0 \overline{K}_4^*(2045)$	$< 3.1 \times 10^{-5}$	CL=90%
Γ_{37}	$\overline{D}^0 K^- \pi^+ (\text{non-resonant})$	$(2.1 \pm 0.8) \times 10^{-4}$	
Γ_{38}	$D_{s2}^*(2573)^- \pi^+, D_{s2}^* \rightarrow \overline{D}^0 K^-$	$(2.6 \pm 0.4) \times 10^{-4}$	
Γ_{39}	$D_{s1}^*(2700)^- \pi^+, D_{s1}^* \rightarrow \overline{D}^0 K^-$	$(1.6 \pm 0.8) \times 10^{-5}$	
Γ_{40}	$D_{s1}^*(2860)^- \pi^+, D_{s1}^* \rightarrow \overline{D}^0 K^-$	$(5 \pm 4) \times 10^{-5}$	
Γ_{41}	$D_{s3}^*(2860)^- \pi^+, D_{s3}^* \rightarrow \overline{D}^0 K^-$	$(2.2 \pm 0.6) \times 10^{-5}$	
Γ_{42}	$\overline{D}^0 K^+ K^-$	$(5.5 \pm 0.8) \times 10^{-5}$	
Γ_{43}	$\overline{D}^0 f_0(980)$	$< 3.1 \times 10^{-6}$	CL=90%
Γ_{44}	$\overline{D}^0 \phi$	$(3.0 \pm 0.5) \times 10^{-5}$	
Γ_{45}	$\overline{D}^{*0} \phi$	$(3.7 \pm 0.6) \times 10^{-5}$	
Γ_{46}	$D^{*\mp} \pi^\pm$	$< 6.1 \times 10^{-6}$	CL=90%
Γ_{47}	$\eta_c \phi$	$(5.0 \pm 0.9) \times 10^{-4}$	
Γ_{48}	$\eta_c \pi^+ \pi^-$	$(1.8 \pm 0.7) \times 10^{-4}$	
Γ_{49}	$J/\psi(1S) \phi$	$(1.08 \pm 0.08) \times 10^{-3}$	
Γ_{50}	$J/\psi(1S) \phi \phi$	$(1.24^{+0.17}_{-0.19}) \times 10^{-5}$	
Γ_{51}	$J/\psi(1S) \pi^0$	$< 1.2 \times 10^{-3}$	CL=90%
Γ_{52}	$J/\psi(1S) \eta$	$(4.0 \pm 0.7) \times 10^{-4}$	S=1.4
Γ_{53}	$J/\psi(1S) K_S^0$	$(1.88 \pm 0.15) \times 10^{-5}$	
Γ_{54}	$J/\psi(1S) \overline{K}^*(892)^0$	$(4.1 \pm 0.4) \times 10^{-5}$	
Γ_{55}	$J/\psi(1S) \eta'$	$(3.3 \pm 0.4) \times 10^{-4}$	
Γ_{56}	$J/\psi(1S) \pi^+ \pi^-$	$(2.09 \pm 0.23) \times 10^{-4}$	S=1.3
Γ_{57}	$J/\psi(1S) f_0(500), f_0 \rightarrow \pi^+ \pi^-$	$< 4 \times 10^{-6}$	CL=90%
Γ_{58}	$J/\psi(1S) \rho, \rho \rightarrow \pi^+ \pi^-$	$< 4 \times 10^{-6}$	CL=90%
Γ_{59}	$J/\psi(1S) f_0(980), f_0 \rightarrow \pi^+ \pi^-$	$(1.28 \pm 0.18) \times 10^{-4}$	S=1.7
Γ_{60}	$J/\psi(1S) f_2(1270), f_2 \rightarrow \pi^+ \pi^-$	$(1.1 \pm 0.4) \times 10^{-6}$	
Γ_{61}	$J/\psi(1S) f_2(1270)_0, f_2 \rightarrow \pi^+ \pi^-$	$(7.5 \pm 1.8) \times 10^{-7}$	
Γ_{62}	$J/\psi(1S) f_2(1270)_{ }, f_2 \rightarrow \pi^+ \pi^-$	$(1.09 \pm 0.34) \times 10^{-6}$	
Γ_{63}	$J/\psi(1S) f_2(1270)_\perp, f_2 \rightarrow \pi^+ \pi^-$	$(1.3 \pm 0.8) \times 10^{-6}$	
Γ_{64}	$J/\psi(1S) f_0(1370), f_0 \rightarrow \pi^+ \pi^-$	$(4.5^{+0.7}_{-4.0}) \times 10^{-5}$	
Γ_{65}	$J/\psi(1S) f_0(1500), f_0 \rightarrow \pi^+ \pi^-$	$(2.11^{+0.40}_{-0.29}) \times 10^{-5}$	
Γ_{66}	$J/\psi(1S) f'_2(1525)_0, f'_2 \rightarrow \pi^+ \pi^-$	$(1.07 \pm 0.24) \times 10^{-6}$	

Γ_{67}	$J/\psi(1S)f'_2(1525)_{ }, f'_2 \rightarrow \pi^+\pi^-$	$(1.3 \pm 2.7) \times 10^{-7}$
Γ_{68}	$J/\psi(1S)f'_2(1525)_{\perp}, f'_2 \rightarrow \pi^+\pi^-$	$(5 \pm 4) \times 10^{-7}$
Γ_{69}	$J/\psi(1S)f_0(1790), f_0 \rightarrow \pi^+\pi^-$	$(5.0 \pm 11.0) \times 10^{-6}$
Γ_{70}	$J/\psi(1S)\pi^+\pi^- (\text{nonresonant})$	$(1.8 \pm 1.1) \times 10^{-5}$
Γ_{71}	$J/\psi(1S)\bar{K}^0\pi^+\pi^-$	$< 4.4 \times 10^{-5}$
Γ_{72}	$J/\psi(1S)K^+K^-$	$(7.9 \pm 0.7) \times 10^{-4}$
Γ_{73}	$J/\psi(1S)K^0K^-\pi^+ + \text{c.c.}$	$(9.3 \pm 1.3) \times 10^{-4}$
Γ_{74}	$J/\psi(1S)\bar{K}^0K^+K^-$	$< 1.2 \times 10^{-5}$
Γ_{75}	$J/\psi(1S)f'_2(1525)$	$(2.6 \pm 0.6) \times 10^{-4}$
Γ_{76}	$J/\psi(1S)p\bar{p}$	$< 4.8 \times 10^{-6}$
Γ_{77}	$J/\psi(1S)\gamma$	$< 7.3 \times 10^{-6}$
Γ_{78}	$J/\psi(1S)\pi^+\pi^-\pi^+\pi^-$	$(7.8 \pm 1.0) \times 10^{-5}$
Γ_{79}	$J/\psi(1S)f_1(1285)$	$(7.0 \pm 1.4) \times 10^{-5}$
Γ_{80}	$\psi(2S)\eta$	$(3.3 \pm 0.9) \times 10^{-4}$
Γ_{81}	$\psi(2S)\eta'$	$(1.29 \pm 0.35) \times 10^{-4}$
Γ_{82}	$\psi(2S)\pi^+\pi^-$	$(7.1 \pm 1.3) \times 10^{-5}$
Γ_{83}	$\psi(2S)\phi$	$(5.4 \pm 0.6) \times 10^{-4}$
Γ_{84}	$\psi(2S)K^-\pi^+$	$(3.1 \pm 0.4) \times 10^{-5}$
Γ_{85}	$\psi(2S)\bar{K}^*(892)^0$	$(3.3 \pm 0.5) \times 10^{-5}$
Γ_{86}	$\chi_{c1}\phi$	$(2.04 \pm 0.30) \times 10^{-4}$
Γ_{87}	$\chi_{c1}K^+K^-$	
Γ_{88}	$\chi_{c2}K^+K^-$	
Γ_{89}	$\pi^+\pi^-$	$(6.8 \pm 0.8) \times 10^{-7}$
Γ_{90}	$\pi^0\pi^0$	$< 2.1 \times 10^{-4}$
Γ_{91}	$\eta\pi^0$	$< 1.0 \times 10^{-3}$
Γ_{92}	$\eta\eta$	$< 1.5 \times 10^{-3}$
Γ_{93}	$\rho^0\rho^0$	$< 3.20 \times 10^{-4}$
Γ_{94}	$\eta'\eta'$	$(3.3 \pm 0.7) \times 10^{-5}$
Γ_{95}	$\eta'\phi$	$< 8.2 \times 10^{-7}$
Γ_{96}	$\phi f_0(980), f_0(980) \rightarrow \pi^+\pi^-$	$(1.12 \pm 0.21) \times 10^{-6}$
Γ_{97}	$\phi f_2(1270), f_2(1270) \rightarrow \pi^+\pi^-$	$(6.1 \pm 1.8) \times 10^{-7}$
Γ_{98}	$\phi\rho^0$	$(2.7 \pm 0.8) \times 10^{-7}$
Γ_{99}	$\phi\pi^+\pi^-$	$(3.5 \pm 0.5) \times 10^{-6}$
Γ_{100}	$\phi\phi$	$(1.87 \pm 0.15) \times 10^{-5}$
Γ_{101}	$\phi\phi\phi$	$(2.2 \pm 0.7) \times 10^{-6}$
Γ_{102}	π^+K^-	$(5.6 \pm 0.6) \times 10^{-6}$
Γ_{103}	K^+K^-	$(2.54 \pm 0.17) \times 10^{-5}$
Γ_{104}	$K^0\bar{K}^0$	$(2.0 \pm 0.6) \times 10^{-5}$
Γ_{105}	$K^0\pi^+\pi^-$	$(9.4 \pm 2.1) \times 10^{-6}$

Γ_{106}	$K^0 K^\pm \pi^\mp$	$(8.4 \pm 0.9) \times 10^{-5}$	
Γ_{107}	$K^*(892)^- \pi^+$	$(2.9 \pm 1.1) \times 10^{-6}$	
Γ_{108}	$K^*(892)^\pm K^\mp$	$(1.12 \pm 0.22) \times 10^{-5}$	
Γ_{109}	$K_S^0 \bar{K}^*(892)^0 + \text{c.c.}$	$(1.6 \pm 0.4) \times 10^{-5}$	
Γ_{110}	$K^0 K^+ K^-$	$(1.3 \pm 0.6) \times 10^{-6}$	
Γ_{111}	$\bar{K}^*(892)^0 \rho^0$	$< 7.67 \times 10^{-4}$	CL=90%
Γ_{112}	$\bar{K}^*(892)^0 K^*(892)^0$	$(1.11 \pm 0.27) \times 10^{-5}$	
Γ_{113}	$K^*(892)^0 \bar{K}_2^*(1430)^0$		
Γ_{114}	$K_2^*(1430)^0 \bar{K}^*(892)^0$		
Γ_{115}	$K_2^*(1430)^0 \bar{K}_2^*(1430)^0$		
Γ_{116}	$\phi K^*(892)^0$	$(1.14 \pm 0.30) \times 10^{-6}$	
Γ_{117}	$p\bar{p}$	$< 1.5 \times 10^{-8}$	CL=90%
Γ_{118}	$p\bar{p} K^+ K^-$	$(4.5 \pm 0.5) \times 10^{-6}$	
Γ_{119}	$p\bar{p} K^+ \pi^-$	$(1.39 \pm 0.26) \times 10^{-6}$	
Γ_{120}	$p\bar{p} \pi^+ \pi^-$	$(4.3 \pm 2.0) \times 10^{-7}$	
Γ_{121}	$p\bar{\Lambda} K^- + \text{c.c.}$	$(5.5 \pm 1.0) \times 10^{-6}$	
Γ_{122}	$\Lambda_c^- \Lambda \pi^+$	$(3.6 \pm 1.6) \times 10^{-4}$	
Γ_{123}	$\Lambda_c^- \Lambda_c^+$	$< 8.0 \times 10^{-5}$	CL=95%

**Lepton Family number (*LF*) violating modes or
 $\Delta B = 1$ weak neutral current (*B1*) modes**

Γ_{124}	$\gamma\gamma$	<i>B1</i>	$< 3.1 \times 10^{-6}$	CL=90%
Γ_{125}	$\phi\gamma$	<i>B1</i>	$(3.4 \pm 0.4) \times 10^{-5}$	
Γ_{126}	$\mu^+ \mu^-$	<i>B1</i>	$(3.0 \pm 0.4) \times 10^{-9}$	
Γ_{127}	$e^+ e^-$	<i>B1</i>	$< 2.8 \times 10^{-7}$	CL=90%
Γ_{128}	$\tau^+ \tau^-$	<i>B1</i>	$< 6.8 \times 10^{-3}$	CL=95%
Γ_{129}	$\mu^+ \mu^- \mu^+ \mu^-$	<i>B1</i>	$< 2.5 \times 10^{-9}$	CL=95%
Γ_{130}	$S P, S \rightarrow \mu^+ \mu^-, P \rightarrow \mu^+ \mu^-$	<i>B1</i>	[b] $< 2.2 \times 10^{-9}$	CL=95%
Γ_{131}	$\phi(1020) \mu^+ \mu^-$	<i>B1</i>	$(8.2 \pm 1.2) \times 10^{-7}$	
Γ_{132}	$\bar{K}^*(892)^0 \mu^+ \mu^-$		$(2.9 \pm 1.1) \times 10^{-8}$	
Γ_{133}	$\pi^+ \pi^- \mu^+ \mu^-$	<i>B1</i>	$(8.4 \pm 1.7) \times 10^{-8}$	
Γ_{134}	$\phi \nu \bar{\nu}$	<i>B1</i>	$< 5.4 \times 10^{-3}$	CL=90%
Γ_{135}	$e^\pm \mu^\mp$	<i>LF</i>	[c] $< 5.4 \times 10^{-9}$	CL=90%

[a] Not a pure measurement. See note at head of B_s^0 Decay Modes.

[b] Here *S* and *P* are the hypothetical scalar and pseudoscalar particles with masses of 2.5 GeV/c² and 214.3 MeV/c², respectively.

[c] The value is for the sum of the charge states or particle/antiparticle states indicated.

CONSTRAINED FIT INFORMATION

An overall fit to 12 branching ratios uses 20 measurements and one constraint to determine 8 parameters. The overall fit has a $\chi^2 = 26.6$ for 13 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_{12}	28					
x_{14}	92	26				
x_{49}	0	0	0			
x_{56}	0	0	0	72		
x_{59}	0	0	0	57	67	
x_{100}	0	0	0	29	21	17
	x_{10}	x_{12}	x_{14}	x_{49}	x_{56}	x_{59}

B_s^0 BRANCHING RATIOS

$$\Gamma(D_s^- \text{anything}) / \Gamma_{\text{total}} \quad \Gamma_1 / \Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.93 ± 0.25 OUR AVERAGE				
0.91 ± 0.18 ± 0.41		¹ DRUTSKOY 07	BELL	$e^+ e^- \rightarrow \gamma(4S)$
0.81 ± 0.24 ± 0.22	90	² BUSKULIC 96E	ALEP	$e^+ e^- \rightarrow Z$
1.56 ± 0.58 ± 0.44	147	³ ACTON 92N	OPAL	$e^+ e^- \rightarrow Z$

¹ The extraction of this result takes into account the correlation between the measurements of $B(\gamma(5S) \rightarrow D_s^- X)$ and $B(\gamma(5S) \rightarrow D^0 X)$.

² BUSKULIC 96E separate $c\bar{c}$ and $b\bar{b}$ sources of D_s^+ mesons using a lifetime tag, subtract generic $\bar{b} \rightarrow W^+ \rightarrow D_s^+$ events, and obtain $B(\bar{b} \rightarrow B_s^0) \times B(B_s^0 \rightarrow D_s^- \text{anything}) = 0.088 \pm 0.020 \pm 0.020$ assuming $B(D_s \rightarrow \phi\pi) = (3.5 \pm 0.4) \times 10^{-2}$ and PDG 1994 values for the relative partial widths to other D_s channels. We evaluate using our current values $B(\bar{b} \rightarrow B_s^0) = 0.107 \pm 0.014$ and $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$. Our first error is their experiment's and our second error is that due to $B(\bar{b} \rightarrow B_s^0)$ and $B(D_s \rightarrow \phi\pi)$.

³ ACTON 92N assume that excess of 147 ± 48 D_s^0 events over that expected from B^0 , B^+ , and $c\bar{c}$ is all from B_s^0 decay. The product branching fraction is measured to be $B(\bar{b} \rightarrow B_s^0)B(B_s^0 \rightarrow D_s^- \text{anything}) \times B(D_s^- \rightarrow \phi\pi^-) = (5.9 \pm 1.9 \pm 1.1) \times 10^{-3}$. We evaluate using our current values $B(\bar{b} \rightarrow B_s^0) = 0.107 \pm 0.014$ and $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$. Our first error is their experiment's and our second error is that due to $B(\bar{b} \rightarrow B_s^0)$ and $B(D_s \rightarrow \phi\pi)$.

$\Gamma(\ell\nu_\ell X)/\Gamma_{\text{total}}$ VALUE (units 10^{-2}) **9.6 ± 0.8 OUR AVERAGE** $9.6 \pm 0.4 \pm 0.7$ $9.5^{+2.5}_{-2.0} {}^{+1.1}_{-1.9}$ DOCUMENT IDTECNCOMMENT¹ OSWALD 13 BELL $e^+ e^- \rightarrow \gamma(5S)$ ² LEES 12A BABR $e^+ e^-$ Γ_2/Γ

- ¹ The measurement corresponds to the average of the electron and muon branching fractions.
² The measurement corresponds to a branching fraction where the lepton originates from bottom decay and is the average between the electron and muon branching fractions. LEES 12A uses the correlation of the production of ϕ mesons in association with a lepton in $e^+ e^-$ data taken at center-of-mass energies between 10.54 and 11.2 GeV.

 $\Gamma(e^+\nu X^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-2}) **$9.1 \pm 0.5 \pm 0.6$** DOCUMENT IDTECNCOMMENTOSWALD 13 BELL $e^+ e^- \rightarrow \gamma(5S)$ Γ_3/Γ $\Gamma(\mu^+\nu X^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-2}) **$10.2 \pm 0.6 \pm 0.8$** DOCUMENT IDTECNCOMMENTOSWALD 13 BELL $e^+ e^- \rightarrow \gamma(5S)$ Γ_4/Γ $\Gamma(D_s^- \ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$ Γ_5/Γ

The values and averages in this section serve only to show what values result if one assumes our $B(\bar{b} \rightarrow B_s^0)$. They cannot be thought of as measurements since the underlying product branching fractions were also used to determine $B(\bar{b} \rightarrow B_s^0)$ as described in the note on "Production and Decay of b -Flavored Hadrons."

VALUE (units 10^{-2}) **8.1 ± 1.3 OUR AVERAGE** $8.2 \pm 0.2 \pm 1.5$ $7.6 \pm 1.2 \pm 2.1$ $10.7 \pm 4.3 \pm 2.9$ $10.3 \pm 3.6 \pm 2.8$ $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ 13 ± 4 ± 4

27

DOCUMENT IDTECNCOMMENT¹ OSWALD 15 BELL $e^+ e^- \rightarrow \gamma(5S)$ ² BUSKULIC 950 ALEP $e^+ e^- \rightarrow Z$ ³ ABREU 92M DLPH $e^+ e^- \rightarrow Z$ ⁴ ACTON 92N OPAL $e^+ e^- \rightarrow Z$ 5 BUSKULIC 92E ALEP $e^+ e^- \rightarrow Z$

¹ Obtains $B_s \rightarrow D_s X e\nu$, and $D_s X \mu\nu$ separately, then combines them by assuming systematic uncertainties are fully correlated, except for the one on lepton identification. The third uncertainty adds in quadrature systematic uncertainties from external sources (number of B_s events, and $D_s^{(*)}$ branching fractions). OSWALD 15 also measures the cross-section $\sigma(e^+ e^- \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = 53.8 \pm 1.4 \pm 5.3$ pb at $\sqrt{s} = 10.86$ GeV.

² BUSKULIC 950 use $D_s \ell$ correlations. The measured product branching ratio is $B(\bar{b} \rightarrow B_s) \times B(B_s \rightarrow D_s^- \ell^+ \nu_\ell \text{anything}) = (0.82 \pm 0.09^{+0.13}_{-0.14})\%$ assuming $B(D_s \rightarrow \phi\pi) = (3.5 \pm 0.4) \times 10^{-2}$ and PDG 1994 values for the relative partial widths to the six other D_s channels used in this analysis. Combined with results from $\gamma(4S)$ experiments this can be used to extract $B(\bar{b} \rightarrow B_s) = (11.0 \pm 1.2^{+2.5}_{-2.6})\%$. We evaluate using our current values $B(\bar{b} \rightarrow B_s^0) = 0.107 \pm 0.014$ and $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$. Our first error is their experiment's and our second error is that due to $B(\bar{b} \rightarrow B_s^0)$ and $B(D_s \rightarrow \phi\pi)$.

³ ABREU 92M measured muons only and obtained product branching ratio $B(Z \rightarrow b\bar{b}) \times B(\bar{b} \rightarrow B_s) \times B(B_s \rightarrow D_s \mu^+ \nu_\mu \text{anything}) \times B(D_s \rightarrow \phi\pi) = (18 \pm 8) \times 10^{-5}$. We evaluate using our current values $B(\bar{b} \rightarrow B_s^0) = 0.107 \pm 0.014$ and $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$. Our first error is their experiment's and our second error is that due to $B(\bar{b} \rightarrow B_s^0)$ and $B(D_s \rightarrow \phi\pi)$. We use $B(Z \rightarrow b\bar{b}) = 2B(Z \rightarrow b\bar{b}) = 2 \times (0.2212 \pm 0.0019)$.

⁴ ACTON 92N is measured using $D_s \rightarrow \phi\pi^+$ and $K^*(892)^0 K^+$ events. The product branching fraction measured is measured to be $B(\bar{b} \rightarrow B_s^0)B(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell \text{anything}) \times B(D_s^- \rightarrow \phi\pi^-) = (3.9 \pm 1.1 \pm 0.8) \times 10^{-4}$. We evaluate using our current values $B(\bar{b} \rightarrow B_s^0) = 0.107 \pm 0.014$ and $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$. Our first error is their experiment's and our second error is that due to $B(\bar{b} \rightarrow B_s^0)$ and $B(D_s \rightarrow \phi\pi)$.

⁵ BUSKULIC 92E is measured using $D_s \rightarrow \phi\pi^+$ and $K^*(892)^0 K^+$ events. They use $2.7 \pm 0.7\%$ for the $\phi\pi^+$ branching fraction. The average product branching fraction is measured to be $B(\bar{b} \rightarrow B_s^0)B(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell \text{anything}) = 0.020 \pm 0.0055^{+0.005}_{-0.006}$. We evaluate using our current values $B(\bar{b} \rightarrow B_s^0) = 0.107 \pm 0.014$ and $B(D_s \rightarrow \phi\pi) = 0.036 \pm 0.009$. Our first error is their experiment's and our second error is that due to $B(\bar{b} \rightarrow B_s^0)$ and $B(D_s \rightarrow \phi\pi)$. Superseded by BUSKULIC 95O.

$\Gamma(D_s^{*-} \ell^+ \nu_\ell \text{anything})/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
5.4±0.4±1.0	¹ OSWALD	15	$e^+ e^- \rightarrow \gamma(5S)$

¹ Obtains $B_s \rightarrow D_s^* X e\nu$, and $D_s^* X \mu\nu$ separately, then combines them by assuming systematic uncertainties are fully correlated, except for the one on lepton identification. The third uncertainty adds in quadrature systematic uncertainties from external sources (number of B_s events, and $D_s^{(*)}$ branching fractions). OSWALD 15 also measures the cross-section $\sigma(e^+ e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)}) = 53.8 \pm 1.4 \pm 5.3 \text{ pb}$ at $\sqrt{s} = 10.86 \text{ GeV}$.

$\Gamma(D_{s1}(2536)^- \mu^+ \nu_\mu, D_{s1}^- \rightarrow D^{*-} K_S^0)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.6±0.7±0.1	¹ ABAZOV	09G	$p\bar{p}$ at 1.96 TeV

¹ ABAZOV 09G reports $[\Gamma(B_s^0 \rightarrow D_{s1}(2536)^- \mu^+ \nu_\mu, D_{s1}^- \rightarrow D^{*-} K_S^0)/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow B_s^0)] = (2.66 \pm 0.52 \pm 0.45) \times 10^{-4}$ which we divide by our best value $B(\bar{b} \rightarrow B_s^0) = (10.3 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{s1}(2536)^- X \mu^+ \nu, D_{s1}^- \rightarrow \bar{D}^0 K^+)/\Gamma(D_s^- \ell^+ \nu_\ell \text{anything})$ Γ_8/Γ_5

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
5.4±1.2±0.5	AAIJ	11A	$p p$ at 7 TeV

$\Gamma(D_{s2}(2573)^- X \mu^+ \nu, D_{s2}^- \rightarrow \bar{D}^0 K^+)/\Gamma(D_s^- \ell^+ \nu_\ell \text{anything})$ Γ_9/Γ_5

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
3.3±1.0±0.4	AAIJ	11A	$p p$ at 7 TeV

$$\frac{\Gamma(D_{s1}(2536)^- X \mu^+ \nu, D_{s1}^- \rightarrow \bar{D}^0 K^+)/\Gamma(D_{s2}(2573)^- X \mu^+ \nu, D_{s2}^- \rightarrow \bar{D}^0 K^+)}{\Gamma_8/\Gamma_9}$$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.61 \pm 0.14 \pm 0.05	¹ AAIJ	11A	LHCb $p p$ at 7 TeV

¹ Not independent of other AAIJ 11A measurements.

$$\frac{\Gamma(D_s^- \pi^+)/\Gamma_{\text{total}}}{\Gamma_{10}/\Gamma}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.00 \pm 0.23 OUR FIT				
2.99 \pm 0.24 OUR AVERAGE				

2.95 \pm 0.05 $^{+0.25}_{-0.28}$	¹ AAIJ	12AG	LHCb	$p p$ at 7 TeV
3.6 \pm 0.5 \pm 0.5	² LOUVOT	09	BELL	$e^+ e^- \rightarrow \gamma(5S)$
2.8 \pm 0.6 \pm 0.1	³ ABULENCIA	07C	CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
6.8 \pm 2.2 \pm 1.6	DRUTSKOY	07A	BELL	Repl. by LOUVOT 09
3.3 \pm 1.1 \pm 0.2	⁴ ABULENCIA	06J	CDF	Repl. by ABULENCIA 07C
<130 seen	⁶ AKERS	94J	OPAL	$e^+ e^- \rightarrow Z$
	1	93G	ALEP	$e^+ e^- \rightarrow Z$

¹ AAIJ 12AG reports $(2.95 \pm 0.05 \pm 0.17)^{+0.18}_{-0.22} \times 10^{-3}$ where the last uncertainty comes from the semileptonic f_s/f_d measurement. We combined the systematics in quadrature.

² LOUVOT 09 reports $(3.67^{+0.35+0.65}_{-0.33-0.645}) \times 10^{-3}$ from a measurement of $[\Gamma(B_s^0 \rightarrow D_s^- \pi^+)/\Gamma_{\text{total}}] \times [B(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)})]$ assuming $B(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.5 \pm 2.6) \times 10^{-2}$, which we rescale to our best value $B(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (20.1 \pm 3.1) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ABULENCIA 07C reports $[\Gamma(B_s^0 \rightarrow D_s^- \pi^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- \pi^+)] = 1.13 \pm 0.08 \pm 0.23$ which we multiply by our best value $B(B^0 \rightarrow D^- \pi^+) = (2.52 \pm 0.13) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ ABULENCIA 06J reports $[\Gamma(B_s^0 \rightarrow D_s^- \pi^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- \pi^+)] = 1.32 \pm 0.18 \pm 0.38$ which we multiply by our best value $B(B^0 \rightarrow D^- \pi^+) = (2.52 \pm 0.13) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ AKERS 94J sees ≤ 6 events and measures the limit on the product branching fraction $f(\bar{b} \rightarrow B_s^0) \cdot B(B_s^0 \rightarrow D_s^- \pi^+) < 1.3\%$ at CL = 90%. We divide by our current value $B(\bar{b} \rightarrow B_s^0) = 0.105$.

$$\frac{\Gamma(D_s^- \rho^+)/\Gamma(D_s^- \pi^+)}{\Gamma_{11}/\Gamma_{10}}$$

VALUE	DOCUMENT ID	TECN	COMMENT
2.3 \pm 0.4 \pm 0.2	LOUVOT	10	BELL $e^+ e^- \rightarrow \gamma(5S)$



$$\frac{\Gamma(D_s^- \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}}{\Gamma_{12}/\Gamma}$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
6.1 \pm 1.0 OUR FIT			
6.3 \pm 1.5 \pm 0.7			

¹ ABULENCIA 07C CDF $p\bar{p}$ at 1.96 TeV

¹ ABULENCIA 07C reports $[\Gamma(B_s^0 \rightarrow D_s^- \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}] / [B(B_s^0 \rightarrow D^- \pi^+ \pi^+ \pi^-)] = 1.05 \pm 0.10 \pm 0.22$ which we multiply by our best value $B(B_s^0 \rightarrow D^- \pi^+ \pi^+ \pi^-) = (6.0 \pm 0.7) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_s^- \pi^+ \pi^+ \pi^-)/\Gamma(D_s^- \pi^+)$	Γ_{12}/Γ_{10}		
VALUE	DOCUMENT ID	TECN	COMMENT
2.05 ± 0.34 OUR FIT			
2.01 ± 0.37 ± 0.20	AAIJ	11E LHCb	$p p$ at 7 TeV

$\Gamma(D_{s1}(2536)^- \pi^+, D_{s1}^- \rightarrow D_s^- \pi^+ \pi^-)/\Gamma(D_s^- \pi^+ \pi^+ \pi^-)$	Γ_{13}/Γ_{12}		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
4.0 ± 1.0 ± 0.4	AAIJ	12AX LHCb	$p p$ at 7 TeV

$\Gamma(D_s^\mp K^\pm)/\Gamma_{\text{total}}$	Γ_{14}/Γ		
VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
2.27 ± 0.19 OUR FIT			

2.3	+1.2	+0.4	¹ LOUVOT 09 BELL $e^+ e^- \rightarrow \gamma(5S)$
------------	-------------	-------------	--

¹ LOUVOT 09 reports $(2.4^{+1.2}_{-1.0} \pm 0.42) \times 10^{-4}$ from a measurement of $[\Gamma(B_s^0 \rightarrow D_s^\mp K^\pm)/\Gamma_{\text{total}}] \times [B(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)})]$ assuming $B(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.5 \pm 2.6) \times 10^{-2}$, which we rescale to our best value $B(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (20.1 \pm 3.1) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_s^\mp K^\pm)/\Gamma(D_s^- \pi^+)$	Γ_{14}/Γ_{10}		
VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
7.55 ± 0.24 OUR FIT			
7.55 ± 0.24 OUR AVERAGE			

7.52 ± 0.15 ± 0.19	AAIJ	15AC LHCb	$p p$ at 7, 8 TeV
9.7 ± 1.8 ± 0.9	AALTONEN	09AQ CDF	$p \bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.46 ± 0.43 ± 0.25	AAIJ	12AG LHCb	Repl. by AAIJ 15AC
--------------------	------	-----------	--------------------

$\Gamma(D_s^- K^+ \pi^+ \pi^-)/\Gamma(D_s^- \pi^+ \pi^+ \pi^-)$	Γ_{15}/Γ_{12}		
VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
5.2 ± 0.5 ± 0.3	AAIJ	12AX LHCb	$p p$ at 7 TeV

$\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$	Γ_{16}/Γ		
VALUE (units 10^{-3}) CL%	DOCUMENT ID	TECN	COMMENT
4.4 ± 0.5 OUR AVERAGE			

4.0 ± 0.2 ± 0.5	¹ AAIJ	13AP LHCb	$p p$ at 7 TeV
5.8 ^{+1.1} _{-0.9} ± 1.3	² ESEN	13 BELL	$e^+ e^- \rightarrow \gamma(5S)$
5.2 ± 0.8 ± 0.6	³ AALTONEN	12C CDF	$p \bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$10.3^{+3.9+2.6}_{-3.2-2.5}$	⁴ ESEN	10	BELL	Repl. by ESEN 13
$10.4^{+3.5+1.1}_{-3.2}$	⁵ AALTONEN	08F	CDF	Repl. by AALTONEN 12C
<67	90	DRUTSKOY	07A	BELL
				Repl. by ESEN 10
¹ Uses $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$.				
² Use $\Gamma(5S) \rightarrow B_s^* \bar{B}_s^*$ decays assuming $B(\Gamma(5S) \rightarrow B_s^* \bar{B}_s^*) = (17.1 \pm 3.0)\%$ and $\Gamma(\Gamma(5S) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Gamma(5S) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (87.0 \pm 1.7)\%$.				
³ AALTONEN 12C reports $(f_s/f_d) (B(B_s^0 \rightarrow D_s^+ D_s^-) / B(B^0 \rightarrow D^- D_s^+)) = 0.183 \pm 0.021 \pm 0.017$. We multiply this result by our best value of $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$ and divide by our best value of f_s/f_d , where $1/2 f_s/f_d = 0.1275 \pm 0.0065$. Our first quoted uncertainty is the combined experiment's uncertainty and our second is the systematic uncertainty from using out best values.				
⁴ Uses $\Gamma(10860) \rightarrow B_s^* \bar{B}_s^*$ assuming $B(\Gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.3 \pm 2.9)\%$ and $\Gamma(\Gamma(10860) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (90.1^{+3.8}_{-4.0})\%$.				
⁵ AALTONEN 08F reports $[\Gamma(B_s^0 \rightarrow D_s^+ D_s^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- D_s^+)] = 1.44^{+0.48}_{-0.44}$ which we multiply by our best value $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(D_s^- D^+)/\Gamma_{\text{total}}$

Γ_{17}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$2.8 \pm 0.4 \pm 0.3$	¹ AAIJ	14AA LHCb	$p p$ at 7 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$3.6 \pm 0.6 \pm 0.5$	² AAIJ	13AP LHCb	Repl. by AAIJ 14AA
¹ AAIJ 14AA reports $[\Gamma(B_s^0 \rightarrow D_s^- D^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- D_s^+)] = 0.038 \pm 0.004 \pm 0.003$ which we multiply by our best value $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value..			
² Uses $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$.			

$\Gamma(D^+ D^-)/\Gamma_{\text{total}}$

Γ_{18}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$2.2 \pm 0.4 \pm 0.4$	¹ AAIJ	13AP LHCb	$p p$ at 7 TeV
¹ Uses $B(B^0 \rightarrow D^- D^+) = (2.11 \pm 0.31) \times 10^{-4}$ and $B(B^+ \rightarrow \bar{D}^0 D_s^+) = (10.1 \pm 1.7) \times 10^{-3}$.			

$\Gamma(D^0 \bar{D}^0)/\Gamma_{\text{total}}$

Γ_{19}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$1.9 \pm 0.3 \pm 0.4$	¹ AAIJ	13AP LHCb	$p p$ at 7 TeV
¹ Uses $B(B^0 \rightarrow D^- D^+) = (2.11 \pm 0.31) \times 10^{-4}$ and $B(B^+ \rightarrow \bar{D}^0 D_s^+) = (10.1 \pm 1.7) \times 10^{-3}$.			

$\Gamma(D_s^{*-} \pi^+)/\Gamma(D_s^- \pi^+)$	VALUE
$0.65^{+0.15}_{-0.13} \pm 0.07$	

Γ_{20}/Γ_{10}	DOCUMENT ID	TECN	COMMENT
	LOUVOT	10	BELL $e^+ e^- \rightarrow \gamma(5S)$

$\Gamma(D_s^{*\mp} K^\pm)/\Gamma(D_s^{*-} \pi^+)$	VALUE
$0.068 \pm 0.005^{+0.003}_{-0.002}$	

Γ_{21}/Γ_{20}	DOCUMENT ID	TECN	COMMENT
	AAIJ	15AD LHCb	$p p$ at 7, 8 TeV

$\Gamma(D_s^{*-} \rho^+)/\Gamma(D_s^- \pi^+)$	VALUE
$3.2 \pm 0.6 \pm 0.3$	

Γ_{22}/Γ_{10}	DOCUMENT ID	TECN	COMMENT
	LOUVOT	10	BELL $e^+ e^- \rightarrow \gamma(5S)$

$\Gamma(D_s^{*-} \rho^+)/\Gamma(D_s^- \rho^+)$	VALUE
$1.4 \pm 0.3 \pm 0.1$	

Γ_{22}/Γ_{11}	DOCUMENT ID	TECN	COMMENT
	¹ LOUVOT	10	BELL $e^+ e^- \rightarrow \gamma(5S)$

¹ Not independent of other LOUVOT 10 measurements.

$[\Gamma(D_s^{*+} D_s^-) + \Gamma(D_s^{*-} D_s^+)]/\Gamma_{\text{total}}$	VALUE (units 10^{-3})	CL%
13.7 ± 1.6 OUR AVERAGE		

Γ_{23}/Γ	DOCUMENT ID	TECN	COMMENT
	1 AAIJ	16P LHCb	$p p$ at 7 TeV

$13.6 \pm 1.0 \pm 1.4$

$17.6^{+2.3}_{-2.2} \pm 4.0$

$12.0 \pm 1.6 \pm 1.4$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$27.5^{+8.3}_{-7.1} \pm 6.9$

<121

90

⁴ ESEN 10 BELL Repl. by ESEN 13

DRUTSKOY 07A BELL Repl. by ESEN 10

¹ AAIJ 16P reports $[\Gamma(B_s^0 \rightarrow D_s^{*+} D_s^-) + \Gamma(D_s^{*-} D_s^+)]/\Gamma_{\text{total}} / [B(B^0 \rightarrow D^- D_s^+)] = 1.88 \pm 0.08 \pm 0.12$ which we multiply by our best value $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Use $\gamma(5S) \rightarrow B_s^* \bar{B}_s^*$ decays assuming $B(\gamma(5S) \rightarrow B_s^* \bar{B}_s^*) = (17.1 \pm 3.0)\%$ and $\Gamma(\gamma(5S) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\gamma(5S) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (87.0 \pm 1.7)\%$.

³ AALTONEN 12C reports $(f_s/f_d) (B(B_s^0 \rightarrow D_s^{*+} D_s^- + D_s^{*-} D_s^+)) / B(B^0 \rightarrow D^- D_s^+) = 0.424 \pm 0.046 \pm 0.035$. We multiply this result by our best value of $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$ and divide by our best value of f_s/f_d , where $1/2 f_s/f_d = 0.1275 \pm 0.0065$. Our first quoted uncertainty is the combined experiment's uncertainty and our second is the systematic uncertainty from using out best values.

⁴ Uses $\gamma(10860) \rightarrow B_s^* \bar{B}_s^*$ assuming $B(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.3 \pm 2.9)\%$ and $\Gamma(\gamma(10860) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (90.1^{+3.8}_{-4.0})\%$.

$\Gamma(D_s^{*+} D_s^{*-})/\Gamma_{\text{total}}$	Γ_{24}/Γ				
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
14.4 ± 2.0 OUR AVERAGE		Error includes scale factor of 1.1.			
12.7 ± 1.3 ± 1.4		¹ AAIJ	16P	LHCb	$p\bar{p}$ at 7 TeV
$19.8^{+3.3+5.2}_{-3.1-5.0}$		² ESEN	13	BELL	$e^{+}e^{-} \rightarrow \gamma(5S)$
18.6 ± 2.8 ± 2.2		³ AALTONEN	12C	CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$30.8^{+12.2+8.5}_{-10.4-8.6}$		⁴ ESEN	10	BELL	Repl. by ESEN 13
<257	90	DRUTSKOY	07A	BELL	Repl. by ESEN 10
¹ AAIJ 16P reports $[\Gamma(B_s^0 \rightarrow D_s^{*+} D_s^{*-})/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- D_s^+)] = 1.76 \pm 0.11 \pm 0.14$ which we multiply by our best value $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
² Use $\gamma(5S) \rightarrow B_s^* \bar{B}_s^*$ decays assuming $B(\gamma(5S) \rightarrow B_s^* \bar{B}_s^*) = (17.1 \pm 3.0)\%$ and $\Gamma(\gamma(5S) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\gamma(5S) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (87.0 \pm 1.7)\%$.					
³ AALTONEN 12C reports $(f_s/f_d) (B(B_s^0 \rightarrow D_s^{*+} D_s^{*-}) / B(B^0 \rightarrow D^- D_s^+)) = 0.654 \pm 0.072 \pm 0.065$. We multiply this result by our best value of $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$ and divide by our best value of f_s/f_d , where $1/2 f_s/f_d = 0.1275 \pm 0.0065$. Our first quoted uncertainty is the combined experiment's uncertainty and our second is the systematic uncertainty from using out best values.					
⁴ Uses $\gamma(10860) \rightarrow B_s^* \bar{B}_s^*$ assuming $B(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.3 \pm 2.9)\%$ and $\Gamma(\gamma(10860) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (90.1^{+3.8}_{-4.0})\%$.					

$\Gamma(D_s^{(*)+} D_s^{*-})/\Gamma_{\text{total}}$	Γ_{25}/Γ				
VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT	
4.5 ± 1.4 OUR EVALUATION		“OUR EVALUATION” is an average using rescaled values of the data listed below.			
3.4 ± 0.4 OUR AVERAGE		The average and rescaling were performed by the Heavy Flavor Averaging Group (HFLAV) and are described at http://www.slac.stanford.edu/xorg/hflav/ . The averaging/rescaling procedure takes into account correlations between the measurements.			
3.07 ± 0.22 ± 0.33		¹ AAIJ	16P	LHCb	$p\bar{p}$ at 7 TeV
$4.32^{+0.42+1.04}_{-0.39-1.03}$		² ESEN	13	BELL	$e^{+}e^{-} \rightarrow \gamma(5S)$
3.6 ± 0.4 ± 0.4		³ AALTONEN	12C	CDF	$p\bar{p}$ at 1.96 TeV
3.5 ± 1.0 ± 1.1		⁴ ABAZOV	09I	D0	$p\bar{p}$ at 1.96 TeV
14 ± 6 ± 3		^{5,6} BARATE	00K	ALEP	$e^{+}e^{-} \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$6.85^{+1.53+1.79}_{-1.30-1.80}$		^{7,8} ESEN	10	BELL	Repl. by ESEN 13
$3.9^{+1.9+1.6}_{-1.7-1.5}$		⁴ ABAZOV	07Y	D0	Repl. by ABAZOV 09I
<0.218	90	BARATE	98Q	ALEP	$e^{+}e^{-} \rightarrow Z$
¹ AAIJ 16P reports $[\Gamma(B_s^0 \rightarrow D_s^{(*)+} D_s^{*-})/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- D_s^+)] = 4.24 \pm 0.14 \pm 0.27$ which we multiply by our best value $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$.					

Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Use $\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*$ decays assuming $B(\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*) = (17.1 \pm 3.0)\%$ and $\Gamma(\Upsilon(5S) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(5S) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (87.0 \pm 1.7)\%$.

³ AALTONEN 12C reports (f_s/f_d) ($B(B_s^0 \rightarrow D_s^{(*)+} D_s^{(*)-}) / B(B^0 \rightarrow D^- D_s^+) = 1.261 \pm 0.095 \pm 0.112$). We multiply this result by our best value of $B(B^0 \rightarrow D^- D_s^+) = (7.2 \pm 0.8) \times 10^{-3}$ and divide by our best value of f_s/f_d , where $1/2 f_s/f_d = 0.1275 \pm 0.0065$. Our first quoted uncertainty is the combined experiment's uncertainty and our second is the systematic uncertainty from using our best values.

⁴ Uses the final states where $D_s^+ \rightarrow \phi \pi^+$ and $D_s^- \rightarrow \phi \mu^- \bar{\nu}_\mu$.

⁵ Reports $B(B_s^0 \text{ (short)} \rightarrow D_s^{(*)} D_s^{(*)}) = (0.23 \pm 0.10 \pm 0.05) \cdot [0.17/B(D_s \rightarrow \phi \chi)]^2$ assuming $B(B_s^0 \rightarrow B_s^0 \text{ (short)}) = 50\%$. We use our best value of $B(D_s \rightarrow \phi \chi) = 15.7 \pm 1.0\%$ to obtain the quoted result.

⁶ Uses $\phi \phi$ correlations from $B_s^0 \text{ (short)} \rightarrow D_s^{(*)+} D_s^{(*)-}$.

⁷ Sum of exclusive $B_s \rightarrow D_s^+ D_s^-$, $B_s \rightarrow D_s^{*\pm} D_s^{\mp}$ and $B_s \rightarrow D_s^{*+} D_s^{*-}$.

⁸ Uses $\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*$ assuming $B(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.3 \pm 2.9)\%$ and $\Gamma(\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (90.1^{+3.8}_{-4.0})\%$.

$\Gamma(\bar{D}^{*0} K^0)/\Gamma_{\text{total}}$

Γ_{26}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$2.8 \pm 1.0 \pm 0.5$	1 AAIJ	16C LHCb	$p p$ at 7, 8 TeV

¹ Measured and normalized to the $B_s^0 \rightarrow \bar{D}^{*0} K_S^0$ decay with $f_s/f_d = 0.259 \pm 0.015$. Signal significance is 4.4 standard deviations.

$\Gamma(\bar{D}^0 K^0)/\Gamma_{\text{total}}$

Γ_{27}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$4.3 \pm 0.5 \pm 0.7$	1 AAIJ	16C LHCb	$p p$ at 7, 8 TeV

¹ Measured and normalized to the $B^0 \rightarrow \bar{D}^0 K_S^0$ decay with $f_s/f_d = 0.259 \pm 0.015$.

$\Gamma(\bar{D}^0 K^- \pi^+)/\Gamma_{\text{total}}$

Γ_{28}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$10.4 \pm 1.1 \pm 0.5$	1 AAIJ	13AQ LHCb	$p p$ at 7 TeV

¹ AAIJ 13AQ reports $[\Gamma(B_s^0 \rightarrow \bar{D}^0 K^- \pi^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-)] = 1.18 \pm 0.05 \pm 0.12$ which we multiply by our best value $B(B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-) = (8.8 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{D}^0 K^*(892)^0)/\Gamma_{\text{total}}$

Γ_{29}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
4.4 ± 0.6 OUR AVERAGE	1 AAIJ	14BH LHCb	$p p$ at 7, 8 TeV

4.29 $\pm 0.09 \pm 0.65$

4.7 $\pm 1.2 \pm 0.3$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.5 $\pm 0.4 \pm 0.4$ 3 AAIJ 13BX LHCb Repl. by AAIJ 14BH

¹ Uses Dalitz plot analysis of $B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$ decays.

² AAIJ 11D reports $[\Gamma(B_s^0 \rightarrow \bar{D}^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow \bar{D}^0 \rho^0)] = 1.48 \pm 0.34 \pm 0.19$ which we multiply by our best value $B(B^0 \rightarrow \bar{D}^0 \rho^0) = (3.21 \pm 0.21) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ AAIJ 13BX reports $[\Gamma(B_s^0 \rightarrow \bar{D}^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow \bar{D}^0 K^*(892)^0)] = 7.8 \pm 0.7 \pm 0.3 \pm 0.6$ which we multiply by our best value $B(B^0 \rightarrow \bar{D}^0 K^*(892)^0) = (4.5 \pm 0.6) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{D}^0 \bar{K}^*(1410))/\Gamma_{\text{total}}$

Γ_{30}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
38.6 ± 11.4 ± 33.3	¹ AAIJ	14BH LHCb	$p p$ at 7, 8 TeV

¹ Uses Dalitz plot analysis of $B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$ decays.

$\Gamma(\bar{D}^0 \bar{K}_0^*(1430))/\Gamma_{\text{total}}$

Γ_{31}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
30.0 ± 2.4 ± 6.8	¹ AAIJ	14BH LHCb	$p p$ at 7, 8 TeV

¹ Uses Dalitz plot analysis of $B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$ decays. Corresponds to the resonant $K_0^*(1430)$ part of LASS parametrization.

$\Gamma(\bar{D}^0 \bar{K}_2^*(1430))/\Gamma_{\text{total}}$

Γ_{32}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
11.1 ± 1.8 ± 3.8	¹ AAIJ	14BH LHCb	$p p$ at 7, 8 TeV

¹ Uses Dalitz plot analysis of $B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$ decays.

$\Gamma(\bar{D}^0 \bar{K}^*(1680))/\Gamma_{\text{total}}$

Γ_{33}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<7.8	90	¹ AAIJ	14BH LHCb	$p p$ at 7, 8 TeV

¹ Uses Dalitz plot analysis of $B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$ decays.

$\Gamma(\bar{D}^0 \bar{K}_0^*(1950))/\Gamma_{\text{total}}$

Γ_{34}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<11	90	¹ AAIJ	14BH LHCb	$p p$ at 7, 8 TeV

¹ Uses Dalitz plot analysis of $B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$ decays.

$\Gamma(\bar{D}^0 \bar{K}_3^*(1780))/\Gamma_{\text{total}}$

Γ_{35}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<2.6	90	¹ AAIJ	14BH LHCb	$p p$ at 7, 8 TeV

¹ Uses Dalitz plot analysis of $B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$ decays.

$\Gamma(\bar{D}^0 \bar{K}_4^*(2045))/\Gamma_{\text{total}}$

Γ_{36}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<3.1	90	¹ AAIJ	14BH LHCb	$p p$ at 7, 8 TeV

¹ Uses Dalitz plot analysis of $B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$ decays.

$\Gamma(\bar{D}^0 K^- \pi^+ (\text{non-resonant})) / \Gamma_{\text{total}}$ Γ_{37}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$20.6 \pm 3.8 \pm 7.3$	¹ AAIJ	14BH LHCb	$p p$ at 7, 8 TeV

¹ Uses Dalitz plot analysis of $B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$ decays. Corresponds to the non-resonant part of the LASS parametrization.

 $\Gamma(D_{s2}^*(2573)^- \pi^+, D_{s2}^* \rightarrow \bar{D}^0 K^-) / \Gamma_{\text{total}}$ Γ_{38}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$25.7 \pm 0.7 \pm 4.0$	¹ AAIJ	14BH LHCb	$p p$ at 7, 8 TeV

¹ Uses Dalitz plot analysis of $B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$ decays.

 $\Gamma(D_{s1}^*(2700)^- \pi^+, D_{s1}^* \rightarrow \bar{D}^0 K^-) / \Gamma_{\text{total}}$ Γ_{39}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.6 \pm 0.4 \pm 0.7$	¹ AAIJ	14BH LHCb	$p p$ at 7, 8 TeV

¹ Uses Dalitz plot analysis of $B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$ decays.

 $\Gamma(D_{s1}^*(2860)^- \pi^+, D_{s1}^* \rightarrow \bar{D}^0 K^-) / \Gamma_{\text{total}}$ Γ_{40}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$5.0 \pm 1.2 \pm 3.4$	¹ AAIJ	14BH LHCb	$p p$ at 7, 8 TeV

¹ Uses Dalitz plot analysis of $B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$ decays.

 $\Gamma(D_{s3}^*(2860)^- \pi^+, D_{s3}^* \rightarrow \bar{D}^0 K^-) / \Gamma_{\text{total}}$ Γ_{41}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.2 \pm 0.1 \pm 0.6$	¹ AAIJ	14BH LHCb	$p p$ at 7, 8 TeV

¹ Uses Dalitz plot analysis of $B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$ decays.

 $\Gamma(\bar{D}^0 K^+ K^-) / \Gamma_{\text{total}}$ Γ_{42}/Γ

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$5.5 \pm 0.7 \pm 0.5$	¹ AAIJ	18AZ LHCb	$p p$ at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.3 \pm 2.0 \pm 0.5$ ^{2,3} AAIJ 12AM LHCb Repl. by AAIJ 18AZ

¹ AAIJ 18AZ reports $[\Gamma(B_s^0 \rightarrow \bar{D}^0 K^+ K^-) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow \bar{D}^0 K^+ K^-)] = 0.930 \pm 0.089 \pm 0.069$ which we multiply by our best value $B(B^0 \rightarrow \bar{D}^0 K^+ K^-) = (5.9 \pm 0.5) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² AAIJ 12AM reports $[\Gamma(B_s^0 \rightarrow \bar{D}^0 K^+ K^-) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow \bar{D}^0 K^+ K^-)] = 0.90 \pm 0.27 \pm 0.20$ which we multiply by our best value $B(B^0 \rightarrow \bar{D}^0 K^+ K^-) = (5.9 \pm 0.5) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Uses $B(b \rightarrow B_s^0)/B(b \rightarrow B^0) = 0.267^{+0.023}_{-0.020}$ measured by the same authors.

 $\Gamma(\bar{D}^0 f_0(980)) / \Gamma_{\text{total}}$ Γ_{43}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.1 \times 10^{-6}$	90	AAIJ	15AG LHCb	$p p$ at 7, 8 TeV

$\Gamma(\bar{D}^0\phi)/\Gamma(\bar{D}^0\bar{K}^*(892)^0)$ Γ_{44}/Γ_{29}

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.069 \pm 0.013 \pm 0.007$	AAIJ	13BX LHCb	Repl. by AAIJ 18AY

 $\Gamma(\bar{D}^0\phi)/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$3.0 \pm 0.4 \pm 0.2$	¹ AAIJ	18AY LHCb	$p\bar{p}$ at 7 and 8 TeV
¹ AAIJ 18AY reports $[\Gamma(B_s^0 \rightarrow \bar{D}^0\phi)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow \bar{D}^0\pi^+\pi^-)] = (3.4 \pm 0.4 \pm 0.3) \times 10^{-2}$ which we multiply by our best value $B(B^0 \rightarrow \bar{D}^0\pi^+\pi^-) = (8.8 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			

 $\Gamma(\bar{D}^{*0}\phi)/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$3.7 \pm 0.6 \pm 0.2$	¹ AAIJ	18AY LHCb	$p\bar{p}$ at 7 and 8 TeV
¹ AAIJ 18AY reports $[\Gamma(B_s^0 \rightarrow \bar{D}^{*0}\phi)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow \bar{D}^0\pi^+\pi^-)] = (4.2 \pm 0.5 \pm 0.4) \times 10^{-2}$ which we multiply by our best value $B(B^0 \rightarrow \bar{D}^0\pi^+\pi^-) = (8.8 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			

 $\Gamma(D^{*\mp}\pi^\pm)/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.1 \times 10^{-6}$	90	¹ AAIJ	13AL LHCb	$p\bar{p}$ at 7 TeV
¹ Uses $f_s/f_d = 0.256 \pm 0.020$ and $B(B^0 \rightarrow D^{*-}\pi^+) = (2.76 \pm 0.13) \times 10^{-3}$.				

 $\Gamma(\eta_c\phi)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$5.01 \pm 0.53 \pm 0.68$	¹ AAIJ	17U LHCb	$p\bar{p}$ at 7, 8 TeV

¹ The last uncertainty includes the limited knowledge of the external branching fractions where the η_c is reconstructed in the $p\bar{p}, K^+K^-\pi^+\pi^-$, $\pi^+\pi^-\pi^+\pi^-$, and $K^+K^-K^+K^-$ decays and $\phi(1020) \rightarrow K^+K^-$.

 $\Gamma(\eta_c\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$1.76 \pm 0.59 \pm 0.31$	¹ AAIJ	17U LHCb	$p\bar{p}$ at 7, 8 TeV
¹ The last uncertainty includes the limited knowledge of the external branching fractions where the η_c is reconstructed in the $p\bar{p}, K^+K^-\pi^+\pi^-$, $\pi^+\pi^-\pi^+\pi^-$, and $K^+K^-K^+K^-$ decays. The significance of the signal, including systematic uncertainties, is 4.6 standard deviations.			

 $\Gamma(J/\psi(1S)\phi)/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.08 ± 0.08 OUR FIT		¹ AAIJ	13AN LHCb	$p\bar{p}$ at 7 TeV
1.10 ± 0.09 OUR AVERAGE		² THORNE	13 BELL	$e^+e^- \rightarrow \gamma(5S)$
$1.050 \pm 0.013 \pm 0.104$		³ ABE	96Q CDF	$p\bar{p}$
$1.25 \pm 0.07 \pm 0.23$				
$1.5 \pm 0.5 \pm 0.1$				

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6	1	⁴ AKERS	94J	OPAL	$e^+ e^- \rightarrow Z$
seen	14	⁵ ABE	93F	CDF	$p\bar{p}$ at 1.8 TeV
seen	1	⁶ ACTON	92N	OPAL	Sup. by AKERS 94J

¹ Uses $f_s/f_d = 0.256 \pm 0.020$ and $B(B^+ \rightarrow J/\psi K^+) = (10.18 \pm 0.42) \times 10^{-4}$.

² Uses $f_s = (17.2 \pm 3.0)\%$ as the fraction of $\Upsilon(5S)$ decaying to $B_s^{(*)}\bar{B}_s^{(*)}$.

³ ABE 96Q reports $[\Gamma(B_s^0 \rightarrow J/\psi(1S)\phi)/\Gamma_{\text{total}}] \times [\Gamma(\bar{b} \rightarrow B_s^0)/[\Gamma(\bar{b} \rightarrow B^+) + \Gamma(\bar{b} \rightarrow B^0)] = (0.185 \pm 0.055 \pm 0.020) \times 10^{-3}$ which we divide by our best value $\Gamma(\bar{b} \rightarrow B_s^0)/[\Gamma(\bar{b} \rightarrow B^+) + \Gamma(\bar{b} \rightarrow B^0)] = 0.1275 \pm 0.0065$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ AKERS 94J sees one event and measures the limit on the product branching fraction $f(\bar{b} \rightarrow B_s^0) \cdot B(B_s^0 \rightarrow J/\psi(1S)\phi) < 7 \times 10^{-4}$ at CL = 90%. We divide by $B(\bar{b} \rightarrow B_s^0) = 0.112$.

⁵ ABE 93F measured using $J/\psi(1S) \rightarrow \mu^+ \mu^-$ and $\phi \rightarrow K^+ K^-$.

⁶ In ACTON 92N a limit on the product branching fraction is measured to be

$$f(\bar{b} \rightarrow B_s^0) \cdot B(B_s^0 \rightarrow J/\psi(1S)\phi) \leq 0.22 \times 10^{-2}.$$

$\Gamma(J/\psi(1S)\phi\phi)/\Gamma(J/\psi(1S)\phi)$

Γ_{50}/Γ_{49}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.15 \pm 0.12^{+0.05}_{-0.09}$	128	¹ AAIJ	16U	LHCb $p\bar{p}$ at 7, 8 TeV

¹ Uses $J/\psi \rightarrow \mu^+ \mu^-$, $\phi \rightarrow K^+ K^-$ decays, and observes 128 ± 13 events of $B_s^0 \rightarrow J/\psi\phi\phi$.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma_{\text{total}}$

Γ_{51}/Γ

VALUE	CL%	DOCUMENT ID	TECN
$<1.2 \times 10^{-3}$	90	¹ ACCIARRI	97C L3

¹ ACCIARRI 97C assumes B^0 production fraction ($39.5 \pm 4.0\%$) and B_s ($12.0 \pm 3.0\%$).

$\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$

Γ_{52}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
4.0 ± 0.7 OUR AVERAGE				Error includes scale factor of 1.4.
$3.6^{+0.5}_{-0.6}{}^{+0.3}_{-0.2}$		¹ AAIJ	13A	LHCb $p\bar{p}$ at 7 TeV
$5.10 \pm 0.50^{+1.17}_{-0.83}$		² LI	12	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<38	90	³ ACCIARRI	97C L3
-----	----	-----------------------	--------

¹ AAIJ 13A reports $[\Gamma(B_s^0 \rightarrow J/\psi(1S)\eta)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)\rho^0)] = 14.0 \pm 1.2^{+1.1}_{-1.5}{}^{+1.1}_{-1.0}$ which we multiply by our best value $B(B^0 \rightarrow J/\psi(1S)\rho^0) = (2.55^{+0.18}_{-0.16}) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Observed for the first time with significances over 10σ . The second error are total systematic uncertainties including the error on $N(B_s^{(*)}\bar{B}_s^{(*)})$.

³ ACCIARRI 97C assumes B^0 production fraction ($39.5 \pm 4.0\%$) and B_s ($12.0 \pm 3.0\%$).

$\Gamma(J/\psi(1S)K_S^0)/\Gamma_{\text{total}}$	Γ_{53}/Γ		
VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
1.88 ± 0.15 OUR AVERAGE			
$1.88 \pm 0.14 \pm 0.07$	¹ AAIJ	15AL LHCb	$p p$ at 7, 8 TeV
$1.9 \pm 0.4 \pm 0.1$	² AALTONEN	11A CDF	$p\bar{p}$ at 1.96 TeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$1.92 \pm 0.15 \pm 0.12$	³ AAIJ	13AB LHCb	Repl. by AAIJ 15AL
$1.92^{+0.25}_{-0.24} \pm 0.12$	⁴ AAIJ	120 LHCb	Repl. by AAIJ 13AB
1 AAIJ 15AL reports $[\Gamma(B_s^0 \rightarrow J/\psi(1S)K_S^0)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)K_S^0)] = (4.31 \pm 0.17 \pm 0.12 \pm 0.25) \times 10^{-2}$ which we multiply by our best value $B(B^0 \rightarrow J/\psi(1S)K_S^0) = (4.36 \pm 0.16) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. 2 AALTONEN 11A reports $[\Gamma(B_s^0 \rightarrow J/\psi(1S)K_S^0)/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow B_s^0)] / [B(\bar{b} \rightarrow B^0)] / [B(B^0 \rightarrow J/\psi(1S)K_S^0)] = (1.09 \pm 0.19 \pm 0.11) \times 10^{-2}$ which we multiply or divide by our best values $B(\bar{b} \rightarrow B_s^0) = (10.3 \pm 0.5) \times 10^{-2}$, $B(\bar{b} \rightarrow B^0) = (40.5 \pm 0.6) \times 10^{-2}$, $B(B^0 \rightarrow J/\psi(1S)K_S^0) = 1/2 \times B(B^0 \rightarrow J/\psi(1S)K^0) = 1/2 \times (8.73 \pm 0.32) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best values. 3 AAIJ 13AB reports $(1.97 \pm 0.14 \pm 0.07 \pm 0.15 \pm 0.08) \times 10^{-5}$ from a measurement of $[\Gamma(B_s^0 \rightarrow J/\psi(1S)K_S^0)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)K^0)] \times [\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0)]$ assuming $B(B^0 \rightarrow J/\psi(1S)K^0) = (8.98 \pm 0.35) \times 10^{-4}$, $\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0) = 0.256 \pm 0.020$, which we rescale to our best values $B(B^0 \rightarrow J/\psi(1S)K^0) = (8.73 \pm 0.32) \times 10^{-4}$, $\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0) = 0.255 \pm 0.013$. Our first error is their experiment's error and our second error is the systematic error from using our best values. 4 AAIJ 120 reports $(1.83 \pm 0.21 \pm 0.10 \pm 0.14 \pm 0.07) \times 10^{-5}$ from a measurement of $[\Gamma(B_s^0 \rightarrow J/\psi(1S)K_S^0)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)K^0)] \times [\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0)]$ assuming $B(B^0 \rightarrow J/\psi(1S)K^0) = (8.71 \pm 0.32) \times 10^{-4}$, $\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0) = 0.267^{+0.021}_{-0.02}$, which we rescale to our best values $B(B^0 \rightarrow J/\psi(1S)K^0) = (8.73 \pm 0.32) \times 10^{-4}$, $\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0) = 0.255 \pm 0.013$. Our first error is their experiment's error and our second error is the systematic error from using our best values.			

$\Gamma(J/\psi(1S)\bar{K}^*(892)^0)/\Gamma_{\text{total}}$	Γ_{54}/Γ		
VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$4.14 \pm 0.18 \pm 0.35$			
$4.14 \pm 0.18 \pm 0.35$	¹ AAIJ	15AV LHCb	$p p$ at 7, 8 TeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$4.4^{+0.5}_{-0.4} \pm 0.8$	² AAIJ	12AP LHCb	Repl. by AAIJ 15AV
$8 \pm 4 \pm 1$	³ AALTONEN	11A CDF	$p\bar{p}$ at 1.96 TeV
1 AAIJ 15AV result combines two measurements with different normalizing modes of $B^0 \rightarrow J/\psi K^*(892)^0$ and $B_s^0 \rightarrow J/\psi \phi$. 2 AAIJ 12AP reports $B(B_s^0 \rightarrow J/\psi(1S)\bar{K}^*(892)^0)/B(B^0 \rightarrow J/\psi(1S)K^*(892)^0) = (3.43^{+0.34}_{-0.36} \pm 0.50) \times 10^{-2}$ and $B(B^0 \rightarrow J/\psi(1S)K^*(892)^0) = (1.29 \pm 0.05 \pm 0.13) \times 10^{-3}$ after correcting for the contribution from $K\pi$ S-wave beneath the K^* peak.			

³AALTONEN 11A reports $[\Gamma(B_s^0 \rightarrow J/\psi(1S)\bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow B_s^0)] / [B(\bar{b} \rightarrow B^0)] / [B(B^0 \rightarrow J/\psi(1S)K^*(892)^0)] = 0.0168 \pm 0.0024 \pm 0.0068$ which we multiply or divide by our best values $B(\bar{b} \rightarrow B_s^0) = (10.3 \pm 0.5) \times 10^{-2}$, $B(\bar{b} \rightarrow B^0) = (40.5 \pm 0.6) \times 10^{-2}$, $B(B^0 \rightarrow J/\psi(1S)K^*(892)^0) = (1.27 \pm 0.05) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(J/\psi(1S)\eta')/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
3.3 ± 0.4 OUR AVERAGE			
3.2 $^{+0.4}_{-0.5}$ ± 0.2	¹ AAIJ	13A	LHCb $p p$ at 7 TeV
$3.71 \pm 0.61^{+0.85}_{-0.60}$	² LI	12	BELL $e^+ e^- \rightarrow \gamma(4S)$

¹AAIJ 13A reports $[\Gamma(B_s^0 \rightarrow J/\psi(1S)\eta')/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)\rho^0)] = 12.7 \pm 1.1^{+0.5+1.0}_{-1.3-0.9}$ which we multiply by our best value $B(B^0 \rightarrow J/\psi(1S)\rho^0) = (2.55^{+0.18}_{-0.16}) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²Observed for the first time with significances over 10σ . The second error are total systematic uncertainties including the error on $N(B_s^{(*)}\bar{B}_s^{(*)})$.

$\Gamma(J/\psi(1S)\eta')/\Gamma(J/\psi(1S)\eta)$ Γ_{55}/Γ_{52}

VALUE	DOCUMENT ID	TECN	COMMENT
0.87 ± 0.06 OUR AVERAGE			
0.902 $\pm 0.072 \pm 0.045$	¹ AAIJ	15D	LHCb $p p$ at 7, 8 TeV
0.90 $\pm 0.09^{+0.06}_{-0.02}$	² AAIJ	13A	LHCb $p p$ at 7 TeV
0.73 $\pm 0.14 \pm 0.02$	² LI	12	BELL $e^+ e^- \rightarrow \gamma(4S)$

¹Uses $J/\psi \rightarrow \mu^+ \mu^-$, $\eta' \rightarrow \rho^0 \gamma$, and $\eta' \rightarrow \eta \pi^+ \pi^-$ decays.

²Strongly correlated with measurements of $\Gamma(J/\psi(1S)\eta)/\Gamma$ and $\Gamma(J/\psi(1S)\eta')/\Gamma$ reported in the same reference.

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\phi)$ Γ_{56}/Γ_{49}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
19.4 ± 1.5 OUR FIT Error includes scale factor of 2.2.			
19.9 ± 0.7 ± 0.2	¹ AAIJ	12AO	LHCb $p p$ at 7 TeV

¹AAIJ 12AO reports $(19.79 \pm 0.47 \pm 0.52) \times 10^{-2}$ from a measurement of $[\Gamma(B_s^0 \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma(B_s^0 \rightarrow J/\psi(1S)\phi)] / [B(\phi(1020) \rightarrow K^+ K^-)]$ assuming $B(\phi(1020) \rightarrow K^+ K^-) = (48.9 \pm 0.5) \times 10^{-2}$, which we rescale to our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(J/\psi(1S)f_0(980), f_0 \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{59}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
1.28 ± 0.18 OUR FIT Error includes scale factor of 1.7.			
1.16 $^{+0.31+0.30}_{-0.19-0.25}$	¹ LI	11	BELL $e^+ e^- \rightarrow \gamma(5S)$

¹The second error includes both the detector systematic and the uncertainty in the number of produced $\gamma(5S) \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$ pairs.

$$\Gamma(J/\psi(1S)f_0(500), f_0 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)f_0(980), f_0 \rightarrow \pi^+\pi^-) \quad \Gamma_{57}/\Gamma_{59}$$

VALUE	CL %	DOCUMENT ID	TECN	COMMENT
<0.034	90	1 AAIJ	14BR LHCb	$p p$ at 7, 8 TeV

¹ Reported first of two solutions using the full Dalitz analysis.

$$\Gamma(J/\psi(1S)\rho, \rho \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{58}/\Gamma_{56}$$

VALUE	CL %	DOCUMENT ID	TECN	COMMENT
<0.017	90	1 AAIJ	14BR LHCb	$p p$ at 7, 8 TeV

¹ Reported first of two solutions using the full Dalitz analysis.

$$\Gamma(J/\psi(1S)f_0(980), f_0 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{59}/\Gamma_{56}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.61 $^{+0.06}_{-0.07}$ OUR FIT			Error includes scale factor of 2.1.

0.703 ± 0.015 $^{+0.004}_{-0.051}$	1 AAIJ	14BR LHCb	$p p$ at 7, 8 TeV
---	--------	-----------	-------------------

¹ Reported first of two solutions using the full Dalitz analysis.

$$\Gamma(J/\psi(1S)f_2(1270)_0, f_2 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{61}/\Gamma_{56}$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0.36 $\pm 0.07 \pm 0.03$	1 AAIJ	14BR LHCb	$p p$ at 7, 8 TeV

¹ Reported first of two solutions using the full Dalitz analysis.

$$\Gamma(J/\psi(1S)f_2(1270)_{||}, f_2 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{62}/\Gamma_{56}$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0.52 ± 0.15 $^{+0.05}_{-0.02}$	1 AAIJ	14BR LHCb	$p p$ at 7, 8 TeV

¹ Reported first of two solutions using the full Dalitz analysis.

$$\Gamma(J/\psi(1S)f_2(1270)_{\perp}, f_2 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{63}/\Gamma_{56}$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0.63 ± 0.34 $^{+0.16}_{-0.08}$	1 AAIJ	14BR LHCb	$p p$ at 7, 8 TeV

¹ Reported first of two solutions using the full Dalitz analysis.

$$\Gamma(J/\psi(1S)f_0(1500), f_0 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{65}/\Gamma_{56}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.101 ± 0.008 $^{+0.011}_{-0.003}$	1 AAIJ	14BR LHCb	$p p$ at 7, 8 TeV

¹ Reported first of two solutions using the full Dalitz analysis.

$$\Gamma(J/\psi(1S)f'_2(1525)_0, f'_2 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{66}/\Gamma_{56}$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0.51 ± 0.09 $^{+0.05}_{-0.04}$	1 AAIJ	14BR LHCb	$p p$ at 7, 8 TeV

¹ Reported first of two solutions using the full Dalitz analysis.

$\Gamma(J/\psi(1S)f'_2(1525)_{\parallel}, f'_2 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_{67}/Γ_{56}

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0.06^{+0.13}_{-0.04}^{±0.01}	¹ AAIJ	14BR LHCb	$p\bar{p}$ at 7, 8 TeV

¹ Reported first of two solutions using the full Dalitz analysis.

 $\Gamma(J/\psi(1S)f'_2(1525)_{\perp}, f'_2 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_{68}/Γ_{56}

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0.26^{+0.06}_{-0.04}	¹ AAIJ	14BR LHCb	$p\bar{p}$ at 7, 8 TeV

¹ Reported first of two solutions using the full Dalitz analysis.

 $\Gamma(J/\psi(1S)f_0(1790), f_0 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_{69}/Γ_{56}

VALUE	DOCUMENT ID	TECN	COMMENT
0.024^{+0.050}_{-0.002}	¹ AAIJ	14BR LHCb	$p\bar{p}$ at 7, 8 TeV

¹ Reported first of two solutions using the full Dalitz analysis.

 $\Gamma(J/\psi(1S)f_0(980), f_0 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\phi)$ Γ_{59}/Γ_{49}

VALUE	DOCUMENT ID	TECN	COMMENT
0.119^{+0.013}_{-0.014} OUR FIT			Error includes scale factor of 2.4.

0.111^{+0.020}_{-0.018} OUR AVERAGE Error includes scale factor of 2.5. See the ideogram below.

0.069 $\pm 0.012 \pm 0.001$ ¹ KHACHATRYAN 16Q CMS $p\bar{p}$ at 7 TeV

0.140 $\pm 0.026 \pm 0.002$ ^{2,3} AAIJ 12AO LHCb $p\bar{p}$ at 7 TeV

0.135 $\pm 0.036 \pm 0.001$ ⁴ ABAZOV 12C D0 $p\bar{p}$ at 1.96 TeV

0.126 $\pm 0.012 \pm 0.001$ ⁵ AALTONEN 11AB CDF $p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.124 $\pm 0.026 \pm 0.001$ ⁶ AAIJ 11 LHCb Repl. by AAIJ 12AO

¹ KHACHATRYAN 16Q reports $[\Gamma(B_s^0 \rightarrow J/\psi(1S)f_0(980), f_0 \rightarrow \pi^+\pi^-)/\Gamma(B_s^0 \rightarrow J/\psi(1S)\phi)] / [B(\phi(1020) \rightarrow K^+K^-)] = 0.140 \pm 0.008 \pm 0.023$ which we multiply by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.


² AAIJ 12AO reports $(13.9 \pm 0.6^{+2.5}_{-1.2}) \times 10^{-2}$ from a measurement of $[\Gamma(B_s^0 \rightarrow J/\psi(1S)f_0(980), f_0 \rightarrow \pi^+\pi^-)/\Gamma(B_s^0 \rightarrow J/\psi(1S)\phi)] / [B(\phi(1020) \rightarrow K^+K^-)]$ assuming $B(\phi(1020) \rightarrow K^+K^-) = (48.9 \pm 0.5) \times 10^{-2}$, which we rescale to our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Measured in Dalitz plot like analysis of $B_s \rightarrow J/\psi\pi^+\pi^-$ decays.

⁴ ABAZOV 12C reports $[\Gamma(B_s^0 \rightarrow J/\psi(1S)f_0(980), f_0 \rightarrow \pi^+\pi^-)/\Gamma(B_s^0 \rightarrow J/\psi(1S)\phi)] / [B(\phi(1020) \rightarrow K^+K^-)] = 0.275 \pm 0.041 \pm 0.061$ which we multiply by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ AALTONEN 11AB reports $[\Gamma(B_s^0 \rightarrow J/\psi(1S) f_0(980), f_0 \rightarrow \pi^+ \pi^-)] / \Gamma(B_s^0 \rightarrow J/\psi(1S) \phi)] / [B(\phi(1020) \rightarrow K^+ K^-)] = 0.257 \pm 0.020 \pm 0.014$ which we multiply by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁶ AAIJ 11 reports $[\Gamma(B_s^0 \rightarrow J/\psi(1S) f_0(980), f_0 \rightarrow \pi^+ \pi^-)] / \Gamma(B_s^0 \rightarrow J/\psi(1S) \phi)] / [B(\phi(1020) \rightarrow K^+ K^-)] = 0.252^{+0.046+0.027}_{-0.032-0.033}$ which we multiply by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.



$\Gamma(J/\psi(1S) f_0(1370), f_0 \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}$

Γ_{64}/Γ

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.34^{+0.11+0.085}_{-0.14-0.054}$ ¹ LI 11 BELL $e^+ e^- \rightarrow \gamma(5S)$

¹ The second error includes both the detector systematic and the uncertainty in the number of produced $\gamma(5S) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}$ pairs.

$\Gamma(J/\psi(1S) f_0(1370), f_0 \rightarrow \pi^+ \pi^-) / \Gamma(J/\psi(1S) \phi)$

Γ_{64}/Γ_{49}

VALUE (units 10^{-2}) DOCUMENT ID TECN COMMENT

$4.22^{+0.55}_{-3.76} \pm 0.05$ ^{1,2} AAIJ 12AO LHCb $p p$ at 7 TeV

¹ AAIJ 12AO reports $(4.19 \pm 0.53^{+0.12}_{-3.7}) \times 10^{-2}$ from a measurement of $[\Gamma(B_s^0 \rightarrow J/\psi(1S)f_0(1370), f_0 \rightarrow \pi^+\pi^-)/\Gamma(B_s^0 \rightarrow J/\psi(1S)\phi)] / [B(\phi(1020) \rightarrow K^+K^-)]$ assuming $B(\phi(1020) \rightarrow K^+K^-) = (48.9 \pm 0.5) \times 10^{-2}$, which we rescale to our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Measured in Dalitz plot like analysis of $B_s \rightarrow J/\psi\pi^+\pi^-$ decays.

$\Gamma(J/\psi(1S)f_2(1270), f_2 \rightarrow \pi^+\pi^-)/\Gamma(J/\psi(1S)\phi)$ Γ_{60}/Γ_{49}

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$9.9^{+3.4}_{-3.6} \pm 0.1$	1,2 AAIJ	12AO LHCb	$p p$ at 7 TeV

¹ AAIJ 12AO reports $(0.098 \pm 0.033^{+0.006}_{-0.015}) \times 10^{-2}$ from a measurement of $[\Gamma(B_s^0 \rightarrow J/\psi(1S)f_2(1270), f_2 \rightarrow \pi^+\pi^-)/\Gamma(B_s^0 \rightarrow J/\psi(1S)\phi)] / [B(\phi(1020) \rightarrow K^+K^-)]$ assuming $B(\phi(1020) \rightarrow K^+K^-) = (48.9 \pm 0.5) \times 10^{-2}$, which we rescale to our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Measured in Dalitz plot like analysis of $B_s \rightarrow J/\psi\pi^+\pi^-$ decays for the f_2 helicity state $\lambda = 0$.

$\Gamma(J/\psi(1S)\pi^+\pi^- \text{ (nonresonant)})/\Gamma(J/\psi(1S)\phi)$ Γ_{70}/Γ_{49}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$1.67^{+1.02}_{-0.32} \pm 0.02$	1,2 AAIJ	12AO LHCb	$p p$ at 7 TeV

¹ AAIJ 12AO reports $(1.66 \pm 0.31^{+0.96}_{-0.08}) \times 10^{-2}$ from a measurement of $[\Gamma(B_s^0 \rightarrow J/\psi(1S)\pi^+\pi^- \text{ (nonresonant)})/\Gamma(B_s^0 \rightarrow J/\psi(1S)\phi)] / [B(\phi(1020) \rightarrow K^+K^-)]$ assuming $B(\phi(1020) \rightarrow K^+K^-) = (48.9 \pm 0.5) \times 10^{-2}$, which we rescale to our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Measured in Dalitz plot like analysis of $B_s \rightarrow J/\psi\pi^+\pi^-$ decays.

$\Gamma(J/\psi(1S)\bar{K}^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{71}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.4 \times 10^{-5}$	90	1 AAIJ	14L LHCb	$p p$ at 7 TeV

¹ Measured with $B(B_s^0 \rightarrow J/\psi K_S^0\pi^+\pi^-) / B(B^0 \rightarrow J/\psi K_S^0\pi^+\pi^-)$ using PDG 12 values for the involved branching fractions.

$\Gamma(J/\psi(1S)K^+K^-)/\Gamma_{\text{total}}$ Γ_{72}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$7.9 \pm 0.7 \text{ OUR AVERAGE}$			
$7.70 \pm 0.08 \pm 0.72$	1 AAIJ	13AN LHCb	$p p$ at 7 TeV

10.1 $\pm 0.9 \pm 2.1$ ² THORNE 13 BELL $e^+e^- \rightarrow \gamma(5S)$

¹ Uses $f_s/f_d = 0.256 \pm 0.020$ and $B(B^+ \rightarrow J/\psi K^+) = (10.18 \pm 0.42) \times 10^{-4}$.

² Uses $f_s = (17.2 \pm 3.0)\%$ as the fraction of $\gamma(5S)$ decaying to $B_s^{(*)}\bar{B}_s^{(*)}$.

$\Gamma(J/\psi(1S)K^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{73}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$9.3 \pm 1.0 \pm 0.9$	¹ AAIJ	14L LHCb	$p p$ at 7 TeV

¹ AAIJ 14L reports $[\Gamma(B_s^0 \rightarrow J/\psi(1S)K^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)K^0 \pi^+ \pi^-)] = 2.12 \pm 0.15 \pm 0.18$ which we multiply by our best value $B(B^0 \rightarrow J/\psi(1S)K^0 \pi^+ \pi^-) = (4.4 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. This is an observation of $B_s^0 \rightarrow J/\psi K_S^0 K^\pm \pi^\mp$ with more than 10 standard deviations.

 $\Gamma(J/\psi(1S)\bar{K}^0 K^+ K^-)/\Gamma_{\text{total}}$ Γ_{74}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<12 \times 10^{-6}$	90	¹ AAIJ	14L LHCb	$p p$ at 7 TeV

¹ Measured with $B(B_s^0 \rightarrow J/\psi K_S^0 K^+ K^-)/B(B^0 \rightarrow J/\psi K_S^0 \pi^+ \pi^-)$ using PDG 12 values for the involved branching fractions.

 $\Gamma(J/\psi(1S)f'_2(1525))/\Gamma(J/\psi(1S)\phi)$ Γ_{75}/Γ_{49}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
21 ± 4 OUR AVERAGE			
21.5 $\pm 4.9 \pm 2.6$	¹ THORNE	13 BELL	$e^+ e^- \rightarrow \gamma(5S)$
21 $\pm 7 \pm 1$	2,3 ABAZOV	12AF D0	$p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

27 $\pm 4 \pm 1$ ⁴ AAIJ 12S LHCb Repl. by AAIJ 13AN

¹ Uses $B(f'_2(1525) \rightarrow K^+ K^-) = (44.4 \pm 1.1)\%$.

² ABAZOV 12AF reports $[\Gamma(B_s^0 \rightarrow J/\psi(1S)f'_2(1525))/\Gamma(B_s^0 \rightarrow J/\psi(1S)\phi)] \times B(f'_2(1525) \rightarrow K^+ K^-) / B(\phi(1020) \rightarrow K^+ K^-) = 0.19 \pm 0.05 \pm 0.04$ which we divide and multiply by our best values $B(f'_2(1525) \rightarrow K^+ K^-) = \frac{1}{2} (88.7 \pm 2.2) \times 10^{-2}$, $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

³ ABAZOV 12AF fits the invariant masses of the $K^+ K^-$ pair in the range $1.35 < M(K^+ K^-) < 2$ GeV.

⁴ AAIJ 12S reports $[(26.4 \pm 2.7 \pm 2.4) \times 10^{-2}$ from a measurement of $\Gamma(B_s^0 \rightarrow J/\psi(1S)f'_2(1525))/\Gamma(B_s^0 \rightarrow J/\psi(1S)\phi)] \times B(f'_2(1525) \rightarrow K^+ K^-) / B(\phi(1020) \rightarrow K^+ K^-)$ assuming $B(f'_2(1525) \rightarrow K^+ K^-) = (44.4 \pm 1.1) \times 10^{-2}$, $B(\phi(1020) \rightarrow K^+ K^-) = (48.9 \pm 0.5) \times 10^{-2}$, which we rescale to our best values $B(f'_2(1525) \rightarrow K^+ K^-) = \frac{1}{2} (88.7 \pm 2.2) \times 10^{-2}$, $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

 $\Gamma(J/\psi(1S)f'_2(1525))/\Gamma_{\text{total}}$ Γ_{75}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$2.61 \pm 0.20^{+0.56}_{-0.50}$	¹ AAIJ	13AN LHCb	$p p$ at 7 TeV

¹ Uses $f_s/f_d = 0.256 \pm 0.020$ and $B(B^+ \rightarrow J/\psi K^+) = (10.18 \pm 0.42) \times 10^{-4}$.

$\Gamma(\psi(2S)\eta)/\Gamma(J/\psi(1S)\eta)$ Γ_{80}/Γ_{52}

VALUE	DOCUMENT ID	TECN	COMMENT
0.83±0.14±0.12	¹ AAIJ	13AA LHCb	$p p$ at 7 TeV

¹ Assuming lepton universality for dimuon decay modes of J/ψ and $\psi(2S)$ mesons, the ratio $B(J/\psi \rightarrow \mu^+ \mu^-)/B(\psi(2S) \rightarrow \mu^+ \mu^-) = B(J/\psi \rightarrow e^+ e^-)/B(\psi(2S) \rightarrow e^+ e^-) = 7.69 \pm 0.19$ was used.

 $\Gamma(\psi(2S)\eta')/\Gamma(J/\psi(1S)\eta')$ Γ_{81}/Γ_{55}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
38.7±9.0±1.6	¹ AAIJ	15D LHCb	$p p$ at 7, 8 TeV

¹ Uses $J/\psi \rightarrow \mu^+ \mu^-$, $\eta' \rightarrow \rho^0 \gamma$, and $\eta' \rightarrow \eta \pi^+ \pi^-$ decays.

 $\Gamma(J/\psi(1S)p\bar{p})/\Gamma_{\text{total}}$ Γ_{76}/Γ

VALUE	CL %	DOCUMENT ID	TECN	COMMENT
<4.8 × 10⁻⁶	90	¹ AAIJ	13z LHCb	$p p$ at 7 TeV

¹ Uses $B(B_s^0 \rightarrow J/\psi(1S)\pi^+\pi^-) = (1.98 \pm 0.20) \times 10^{-4}$.

 $\Gamma(J/\psi(1S)\gamma)/\Gamma_{\text{total}}$ Γ_{77}/Γ

VALUE	CL %	DOCUMENT ID	TECN	COMMENT
<7.3 × 10⁻⁶	90	¹ AAIJ	15BB LHCb	$p p$ at 7, 8 TeV

¹ Branching fractions of normalization modes $B_s^0 \rightarrow J/\psi \gamma X$ taken from PDG 14. Uses $f_s/f_d = 0.259 \pm 0.015$.

 $\Gamma(J/\psi(1S)\pi^+\pi^-\pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_{78}/Γ_{56}

VALUE	DOCUMENT ID	TECN	COMMENT
0.371±0.015±0.022	¹ AAIJ	14Y LHCb	$p p$ at 7, 8 TeV

¹ Excludes contributions from $\psi(2S)$ and $\chi_{c1}(3872)$ decaying to $J/\psi(1S)\pi^+\pi^-$.

 $\Gamma(J/\psi(1S)f_1(1285))/\Gamma_{\text{total}}$ Γ_{79}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
7.0±1.3±0.4	¹ AAIJ	14Y LHCb	$p p$ at 7, 8 TeV

¹ AAIJ 14Y reports $(7.14 \pm 0.99)^{+0.83}_{-0.91} \pm 0.41) \times 10^{-5}$ from a measurement of $[\Gamma(B_s^0 \rightarrow J/\psi(1S)f_1(1285))/\Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow 2\pi^+ 2\pi^-)]$ assuming $B(f_1(1285) \rightarrow 2\pi^+ 2\pi^-) = 0.11^{+0.007}_{-0.006}$, which we rescale to our best value $B(f_1(1285) \rightarrow 2\pi^+ 2\pi^-) = (11.2^{+0.7}_{-0.6}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\psi(2S)\phi)/\Gamma_{\text{total}}$ Γ_{83}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

seen 1 BUSKULIC 93G ALEP $e^+ e^- \rightarrow Z$

$\Gamma(\psi(2S)\phi)/\Gamma(J/\psi(1S)\phi)$ Γ_{83}/Γ_{49}

VALUE	DOCUMENT ID	TECN	COMMENT
0.503±0.034 OUR AVERAGE			
0.500±0.034±0.011	1,2 AAIJ	12L	LHCb $p\bar{p}$ at 7 TeV
0.53 ± 0.10 ± 0.09	ABAZOV	09Y	D0 $p\bar{p}$ at 1.96 TeV
0.52 ± 0.13 ± 0.07	ABULENCIA	06N	CDF $p\bar{p}$ at 1.96 TeV
¹ AAIJ 12L reports $0.489 \pm 0.026 \pm 0.021 \pm 0.012$ from a measurement of $[\Gamma(B_s^0 \rightarrow \psi(2S)\phi)/\Gamma(B_s^0 \rightarrow J/\psi(1S)\phi)] \times [B(J/\psi(1S) \rightarrow e^+e^-)] / [B(\psi(2S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = (5.94 \pm 0.06) \times 10^{-2}$, $B(\psi(2S) \rightarrow e^+e^-) = (7.72 \pm 0.17) \times 10^{-3}$, which we rescale to our best values $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032) \times 10^{-2}$, $B(\psi(2S) \rightarrow e^+e^-) = (7.93 \pm 0.17) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best values. ² Assumes $B(J/\psi \rightarrow \mu^+\mu^-) / B(\psi(2S) \rightarrow \mu^+\mu^-) = B(J/\psi \rightarrow e^+e^-) / B(\psi(2S) \rightarrow e^+e^-) = 7.69 \pm 0.19$.			

 $\Gamma(\psi(2S)K^-\pi^+)/\Gamma_{\text{total}}$ Γ_{84}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
3.12±0.30±0.21			
1 AAIJ	15U	LHCb	$p\bar{p}$ at 7, 8 TeV

¹ AAIJ 15U reports $[\Gamma(B_s^0 \rightarrow \psi(2S)K^-\pi^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow \psi(2S)K^+\pi^-)] = (5.38 \pm 0.36 \pm 0.22 \pm 0.31) \times 10^{-2}$ which we multiply by our best value $B(B^0 \rightarrow \psi(2S)K^+\pi^-) = (5.8 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\psi(2S)\bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{85}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
3.3±0.5^{+0.2}_{-0.3}			
1 AAIJ	15U	LHCb	$p\bar{p}$ at 7, 8 TeV

¹ AAIJ 15U reports $[\Gamma(B_s^0 \rightarrow \psi(2S)\bar{K}^*(892)^0)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow \psi(2S)\bar{K}^*(892)^0)] = (5.58 \pm 0.57 \pm 0.40 \pm 0.32) \times 10^{-2}$ which we multiply by our best value $B(B^0 \rightarrow \psi(2S)\bar{K}^*(892)^0) = (5.9 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\chi_{c1}\phi)/\Gamma(J/\psi(1S)\phi)$ Γ_{86}/Γ_{49}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
18.9±1.8±1.5			
1 AAIJ	13AC	LHCb	$p\bar{p}$ at 7 TeV

¹ Uses $B(\chi_{c1} \rightarrow J/\psi\gamma) = (34.4 \pm 1.5)\%$.

 $\Gamma(\chi_{c2}K^+K^-)/\Gamma(\chi_{c1}K^+K^-)$ Γ_{88}/Γ_{87}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
17.1±3.1±1.0			
1 AAIJ	18AC	LHCb	$p\bar{p}$ at 7, 8, 13 TeV

¹ Measures the ratio for ± 15 MeV window around ϕ mass.

$\Gamma(\psi(2S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$	Γ_{82}/Γ_{56}
---	---------------------------

VALUE	DOCUMENT ID	TECN	COMMENT
0.34±0.04±0.03	¹ AAIJ	13AA LHCb	$p p$ at 7 TeV

¹ Assuming lepton universality for dimuon decay modes of J/ψ and $\psi(2S)$ mesons, the ratio $B(J/\psi \rightarrow \mu^+ \mu^-)/B(\psi(2S) \rightarrow \mu^+ \mu^-) = B(J/\psi \rightarrow e^+ e^-)/B(\psi(2S) \rightarrow e^+ e^-) = 7.69 \pm 0.19$ was used.

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_{89}/Γ
--	----------------------

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
6.8±0.8 OUR AVERAGE				
7.0±0.8±0.4		¹ AAIJ	17G LHCb	$p p$ at 7 and 8 TeV
6.1±1.7±0.3		² AALTONEN	12L CDF	$p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$10.0^{+2.4}_{-2.0} \pm 0.6$		³ AAIJ	12AR LHCb	Repl. by AAIJ 17G
< 120	90	⁴ PENG	10 BELL	$e^+ e^- \rightarrow \Upsilon(5S)$
< 12	90	⁵ AALTONEN	09C CDF	Repl. by AALTONEN 12L
< 17	90	⁶ ABULENCIA,A	06D CDF	Repl. by AALTONEN 09C
< 2320	90	⁷ ABE	00C SLD	$e^+ e^- \rightarrow Z$
< 1700	90	⁸ BUSKULIC	96V ALEP	$e^+ e^- \rightarrow Z$

¹ AAIJ 17G reports $[\Gamma(B_s^0 \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^+\pi^-)] \times [\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0)] = (9.15 \pm 0.71 \pm 0.83) \times 10^{-3}$ which we multiply or divide by our best values $B(B^0 \rightarrow K^+\pi^-) = (1.96 \pm 0.05) \times 10^{-5}$, $\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0) = 0.255 \pm 0.013$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² AALTONEN 12L reports $[\Gamma(B_s^0 \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^+\pi^-)] \times [\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0)] = 0.008 \pm 0.002 \pm 0.001$ which we multiply or divide by our best values $B(B^0 \rightarrow K^+\pi^-) = (1.96 \pm 0.05) \times 10^{-5}$, $\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0) = 0.255 \pm 0.013$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

³ AAIJ 12AR reports $[\Gamma(B_s^0 \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow \pi^+\pi^-)] \times [\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0)] = 0.050^{+0.011}_{-0.009} \pm 0.004$ which we multiply or divide by our best values $B(B^0 \rightarrow \pi^+\pi^-) = (5.12 \pm 0.19) \times 10^{-6}$, $\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0) = 0.255 \pm 0.013$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

⁴ Uses $\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*$ and assumes $B(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.3 \pm 2.9)\%$ and $\Gamma(\Upsilon(10860) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\Upsilon(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (90.1^{+3.8}_{-4.0})\%$.

⁵ Obtains this result from $(f_s/f_d) \cdot B(B_s \rightarrow \pi^+\pi^-)/B(B^0 \rightarrow K^+\pi^-) = 0.007 \pm 0.004 \pm 0.005$, assuming $f_s/f_d = 0.276 \pm 0.034$ and $B(B^0 \rightarrow K^+\pi^-) = (19.4 \pm 0.6) \times 10^{-6}$.

⁶ ABULENCIA,A 06D obtains this from $B(B_s \rightarrow \pi^+\pi^-) / B(B_s \rightarrow K^+K^-) < 0.05$ at 90% CL, assuming $B(B_s \rightarrow K^+K^-) = (33 \pm 6 \pm 7) \times 10^{-6}$.

⁷ ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

⁸ BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

$\Gamma(\pi^0\pi^0)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{90}/Γ
$<2.1 \times 10^{-4}$	90	¹ ACCIARRI	95H L3	$e^+e^- \rightarrow Z$	

¹ ACCIARRI 95H assumes $f_{B^0} = 39.5 \pm 4.0$ and $f_{B_s} = 12.0 \pm 3.0\%$.

 $\Gamma(\eta\pi^0)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{91}/Γ
$<1.0 \times 10^{-3}$	90	¹ ACCIARRI	95H L3	$e^+e^- \rightarrow Z$	

¹ ACCIARRI 95H assumes $f_{B^0} = 39.5 \pm 4.0$ and $f_{B_s} = 12.0 \pm 3.0\%$.

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{92}/Γ
$<1.5 \times 10^{-3}$	90	¹ ACCIARRI	95H L3	$e^+e^- \rightarrow Z$	

¹ ACCIARRI 95H assumes $f_{B^0} = 39.5 \pm 4.0$ and $f_{B_s} = 12.0 \pm 3.0\%$.

 $\Gamma(\rho^0\rho^0)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{93}/Γ
$<3.20 \times 10^{-4}$	90	¹ ABE	00C SLD	$e^+e^- \rightarrow Z$	

¹ ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

 $\Gamma(\eta'\eta')/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT	Γ_{94}/Γ
$3.3 \pm 0.7 \pm 0.1$	¹ AAIJ	150	LHCb $p\bar{p}$ at 7, 8 TeV	

¹ AAIJ 150 reports $[\Gamma(B_s^0 \rightarrow \eta'\eta')/\Gamma_{\text{total}}] / [B(B^+ \rightarrow \eta' K^+)] = 0.47 \pm 0.09 \pm 0.04$ which we multiply by our best value $B(B^+ \rightarrow \eta' K^+) = (7.06 \pm 0.25) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\eta'\phi)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{95}/Γ
$<0.82 \times 10^{-6}$	90	¹ AAIJ	17BA LHCb	$p\bar{p}$ at 7, 8 TeV	

¹ Corresponds to the 95% CL upper limit 1.01×10^{-6} . Uses the normalization mode $B^+ \rightarrow \eta' K^+$ with branching fraction $(70.6 \pm 2.5) \times 10^{-6}$ and the ratio of hadronisation fractions $f_s/f_d = 0.259 \pm 0.015$, which is assumed equal to f_s/f_u .

 $\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT	Γ_{99}/Γ
$3.48 \pm 0.23 \pm 0.39$	¹ AAIJ	17A LHCb	$p\bar{p}$ at 7, 8 TeV	

¹ Inclusive decays in mass range $400 < m(\pi^+\pi^-) < 1600$ MeV/c².

$\Gamma(\phi\rho^0)/\Gamma_{\text{total}}$				Γ_{98}/Γ
<u>VALUE (units 10^{-7})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.7 \pm 0.7 \pm 0.3$		1 AAIJ	17A LHCb	$p p$ at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6170 90 2 ABE 00C SLD $e^+ e^- \rightarrow Z$

¹ Signal evidence is 4 standard deviations.

² ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

$\Gamma(\phi f_0(980), f_0(980) \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}$				Γ_{96}/Γ
<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$1.12 \pm 0.16 \pm 0.14$	1 AAIJ	17A LHCb	$p p$ at 7, 8 TeV	

¹ Signal is observed with 8 standard deviations significance.

$\Gamma(\phi f_2(1270), f_2(1270) \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}$				Γ_{97}/Γ
<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$0.61 \pm 0.13^{+0.13}_{-0.08}$	1 AAIJ	17A LHCb	$p p$ at 7, 8 TeV	

¹ Signal is observed with 5 standard deviations significance.

$\Gamma(\phi\phi)/\Gamma_{\text{total}}$				Γ_{100}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
18.7 ± 1.5 OUR FIT				
$18.5 \pm 1.4 \pm 1.0$		1 AAIJ	15AS LHCb	$p p$ at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$14^{+6}_{-5} \pm 6$ 2 ACOSTA 05J CDF Repl. by AALTONEN 11AN
<1183 90 3 ABE 00C SLD $e^+ e^- \rightarrow Z$

¹ AAIJ 15AS reports $[\Gamma(B_s^0 \rightarrow \phi\phi)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^*(892)^0 \phi)] = 1.84 \pm 0.05 \pm 0.13$ which we multiply by our best value $B(B^0 \rightarrow K^*(892)^0 \phi) = (1.00 \pm 0.05) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Uses $B(B^0 \rightarrow J/\psi\phi) = (1.38 \pm 0.49) \times 10^{-3}$ and production cross-section ratio of $\sigma(B_s)/\sigma(B^0) = 0.26 \pm 0.04$.

³ ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

$\Gamma(\phi\phi)/\Gamma(J/\psi(1S)\phi)$				Γ_{100}/Γ_{49}
<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.73 ± 0.16 OUR FIT				
$1.78 \pm 0.14 \pm 0.20$	AALTONEN	11AN CDF	$p\bar{p}$ at 1.96 TeV	

$\Gamma(\phi\phi\phi)/\Gamma(\phi\phi)$				$\Gamma_{101}/\Gamma_{100}$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$0.117 \pm 0.030 \pm 0.015$	AAIJ	17BB LHCb	$p p$ at 7, 8 TeV	

$\Gamma(\pi^+ K^-)/\Gamma_{\text{total}}$ Γ_{102}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.6±0.6 OUR AVERAGE				
5.7±0.7±0.3		¹ AAIJ	12AR LHCb	$p p$ at 7 TeV
5.5±0.9±0.3		² AALTONEN	09C CDF	$p \bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 26	90	³ PENG	10 BELL	$e^+ e^- \rightarrow \gamma(5S)$
< 5.6	90	⁴ ABULENCIA,A	06D CDF	Repl. by AALTONEN 09C
<261	90	⁵ ABE	00C SLD	$e^+ e^- \rightarrow Z$
<210	90	⁶ BUSKULIC	96V ALEP	$e^+ e^- \rightarrow Z$
<260	90	⁷ AKERS	94L OPAL	$e^+ e^- \rightarrow Z$

¹ AAIJ 12AR reports $[\Gamma(B_s^0 \rightarrow \pi^+ K^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^+ \pi^-)] \times [\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0)] = 0.074 \pm 0.006 \pm 0.006$ which we multiply or divide by our best values $B(B^0 \rightarrow K^+ \pi^-) = (1.96 \pm 0.05) \times 10^{-5}$, $\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0) = 0.255 \pm 0.013$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² AALTONEN 09C reports $[\Gamma(B_s^0 \rightarrow \pi^+ K^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^+ \pi^-)] \times [B(\bar{b} \rightarrow B_s^0)/B(\bar{b} \rightarrow B^0)] / [B(\bar{b} \rightarrow B^0)] = 0.071 \pm 0.010 \pm 0.007$ which we multiply or divide by our best values $B(B^0 \rightarrow K^+ \pi^-) = (1.96 \pm 0.05) \times 10^{-5}$, $B(\bar{b} \rightarrow B_s^0) = (10.3 \pm 0.5) \times 10^{-2}$, $B(\bar{b} \rightarrow B^0) = (40.5 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

³ Uses $\gamma(10860) \rightarrow B_s^* \bar{B}_s^*$ and assumes $B(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.3 \pm 2.9)\%$ and $\Gamma(\gamma(10860) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (90.1^{+3.8}_{-4.0})\%$.

⁴ ABULENCIA,A 06D obtains this from $(f_s/f_d)(B(B_s \rightarrow \pi^+ K^-) / B(B^0 \rightarrow K^+ \pi^-)) < 0.08$ at 90% CL, assuming $f_s/f_d = 0.260 \pm 0.039$ and $B(B^0 \rightarrow K^+ \pi^-) = (18.9 \pm 0.7) \times 10^{-6}$.

⁵ ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

⁶ BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

⁷ Assumes $B(Z \rightarrow b\bar{b}) = 0.217$ and B_d^0 (B_s^0) fraction 39.5% (12%).

 $\Gamma(K^+ K^-)/\Gamma_{\text{total}}$ Γ_{103}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
25.4±1.7 OUR AVERAGE				
24.3±1.6±1.4		¹ AAIJ	12AR LHCb	$p p$ at 7 TeV
26.6±2.2±1.5		² AALTONEN	11N CDF	$p \bar{p}$ at 1.96 TeV
38 ± ¹⁰ ₉ ±7		³ PENG	10 BELL	$e^+ e^- \rightarrow \gamma(5S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<310	90	DRUTSKOY	07A BELL	$e^+ e^- \rightarrow \gamma(5S)$
33 ± 6 ± 7		⁴ ABULENCIA,A	06D CDF	Repl. by AALTONEN 11N
<283	90	⁵ ABE	00C SLD	$e^+ e^- \rightarrow Z$
< 59	90	⁶ BUSKULIC	96V ALEP	$e^+ e^- \rightarrow Z$
<140	90	⁷ AKERS	94L OPAL	$e^+ e^- \rightarrow Z$

¹ AAIJ 12AR reports $[\Gamma(B_s^0 \rightarrow K^+ K^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^+ \pi^-)] \times [\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0)] = 0.316 \pm 0.009 \pm 0.019$ which we multiply or divide by our best values

$B(B^0 \rightarrow K^+ \pi^-) = (1.96 \pm 0.05) \times 10^{-5}$, $\Gamma(\bar{B} \rightarrow B_s^0)/\Gamma(\bar{B} \rightarrow B^0) = 0.255 \pm 0.013$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

²AALTONEN 11N reports (f_s/f_d) $(B(B_s^0 \rightarrow K^+ K^-) / B(B^0 \rightarrow K^+ \pi^-)) = 0.347 \pm 0.020 \pm 0.021$. We multiply this result by our best value of $B(B^0 \rightarrow K^+ \pi^-) = (1.96 \pm 0.05) \times 10^{-5}$ and divide by our best value of f_s/f_d , where $1/2 f_s/f_d = 0.1275 \pm 0.0065$. Our first quoted uncertainty is the combined experiment's uncertainty and our second is the systematic uncertainty from using out best values.

³Uses $\gamma(10860) \rightarrow B_s^* \bar{B}_s^*$ and assumes $B(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.3 \pm 2.9)\%$ and $\Gamma(\gamma(10860) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (90.1 \pm 3.8) \pm 4.0\%$.

⁴ABULENCIA,A 06D obtains this from (f_s/f_d) $(B(B_s \rightarrow K^+ K^-) / B(B^0 \rightarrow K^+ \pi^-)) = 0.46 \pm 0.08 \pm 0.07$, assuming $f_s/f_d = 0.260 \pm 0.039$ and $B(B^0 \rightarrow K^+ \pi^-) = (18.9 \pm 0.7) \times 10^{-6}$.

⁵ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7 \pm 1.8) \pm 2.2\%$ and $f_{B_s} = (10.5 \pm 1.8) \pm 2.2\%$.

⁶BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

⁷Assumes $B(Z \rightarrow b\bar{b}) = 0.217$ and B_d^0 (B_s^0) fraction 39.5% (12%).

$\Gamma(K^0 \bar{K}^0)/\Gamma_{\text{total}}$

Γ_{104}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
$1.96 \pm 0.58 \pm 0.10 \pm 0.20$		¹ PAL	16	BELL $e^+ e^- \rightarrow \gamma(5S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.6 90 ²PENG 10 BELL Repl. by PAL 16

¹Observed in $B_s^0 \rightarrow K_S^0 K_S^0$ with significance of 5.1σ . The last uncertainty is due to the uncertainty of the total number of $B_s^0 \bar{B}_s^0$ pairs.

²Uses $\gamma(10860) \rightarrow B_s^* \bar{B}_s^*$ and assumes $B(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (19.3 \pm 2.9)\%$ and $\Gamma(\gamma(10860) \rightarrow B_s^* \bar{B}_s^*) / \Gamma(\gamma(10860) \rightarrow B_s^{(*)} \bar{B}_s^{(*)}) = (90.1 \pm 3.8) \pm 4.0\%$.

$\Gamma(K^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$

Γ_{105}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$9.4 \pm 2.1 \pm 0.3$	^{1,2} AAIJ	17BP LHCb	$p p$ at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

14 ± 4 ± 1 ³AAIJ 13BP LHCb Repl. by AAIJ 17BP

¹AAIJ 17BP reports $[\Gamma(B_s^0 \rightarrow K^0 \pi^+ \pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^0 \pi^+ \pi^-)] = 0.191 \pm 0.027 \pm 0.033$ which we multiply by our best value $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = (4.94 \pm 0.18) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²Used $f_s/f_d = 0.259 \pm 0.015$.

³AAIJ 13BP reports $[\Gamma(B_s^0 \rightarrow K^0 \pi^+ \pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^0 \pi^+ \pi^-)] = 0.29 \pm 0.06 \pm 0.04$ which we multiply by our best value $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = (4.94 \pm 0.18) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^-\pi^+)/\Gamma_{\text{total}}$	Γ_{107}/Γ		
<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.9 \pm 1.0 \pm 0.2$	1,2 AAIJ	14BM LHCb	$p p$ at 7 TeV

¹ AAIJ 14BM reports $[\Gamma(B_s^0 \rightarrow K^*(892)^-\pi^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^*(892)^+\pi^-)] = 0.39 \pm 0.13 \pm 0.05$ which we multiply by our best value $B(B^0 \rightarrow K^*(892)^+\pi^-) = (7.5 \pm 0.4) \times 10^{-6}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Uses $f_s/f_d = 0.259 \pm 0.015$.

$\Gamma(K^0 K^\pm \pi^\mp)/\Gamma_{\text{total}}$	Γ_{106}/Γ		
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$8.4 \pm 0.8 \pm 0.3$	1,2 AAIJ	17BP LHCb	$p p$ at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$7.3 \pm 0.9 \pm 0.3$	3 AAIJ	13BP LHCb	Repl. by AAIJ 17BP
-----------------------	--------	-----------	--------------------

¹ AAIJ 17BP reports $[\Gamma(B_s^0 \rightarrow K^0 K^\pm \pi^\mp)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^0 \pi^+ \pi^-)] = 1.70 \pm 0.07 \pm 0.15$ which we multiply by our best value $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = (4.94 \pm 0.18) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Used $f_s/f_d = 0.259 \pm 0.015$.

³ AAIJ 13BP reports $[\Gamma(B_s^0 \rightarrow K^0 K^\pm \pi^\mp)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^0 \pi^+ \pi^-)] = 1.48 \pm 0.12 \pm 0.14$ which we multiply by our best value $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = (4.94 \pm 0.18) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^\pm K^\mp)/\Gamma_{\text{total}}$	Γ_{108}/Γ		
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.12 \pm 0.21 \begin{array}{l} +0.07 \\[-4pt] -0.06 \end{array}$	1,2 AAIJ	14BM LHCb	$p p$ at 7 TeV

¹ AAIJ 14BM reports $[\Gamma(B_s^0 \rightarrow K^*(892)^\pm K^\mp)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^*(892)^+\pi^-)] = 1.49 \pm 0.22 \pm 0.18$ which we multiply by our best value $B(B^0 \rightarrow K^*(892)^+\pi^-) = (7.5 \pm 0.4) \times 10^{-6}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Uses $f_s/f_d = 0.259 \pm 0.015$.

$\Gamma(K_s^0 \bar{K}^*(892)^0 + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{109}/Γ		
<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$16.4 \pm 3.4 \pm 2.3$	1 AAIJ	16 LHCb	$p p$ at 7 TeV

¹ Measured relative to $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ using the value of $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = (4.96 \pm 0.2) \times 10^{-5}$.

$\Gamma(K^0 K^+ K^-)/\Gamma_{\text{total}}$	Γ_{110}/Γ			
<u>VALUE (units 10^{-7})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$12.9 \pm 6.4 \pm 0.5$	1,2,3 AAIJ	17BP LHCb	$p p$ at 7, 8 TeV	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<34	90	4 AAIJ	13BP LHCb	Repl. by AAIJ 17BP
-----	----	--------	-----------	--------------------

¹ AAIJ 17BP reports $[\Gamma(B_s^0 \rightarrow K^0 K^+ K^-)/\Gamma_{\text{total}}] / [\Gamma(B^0 \rightarrow K^0 \pi^+ \pi^-)] = 0.026 \pm 0.011 \pm 0.007$ which we multiply by our best value $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = (4.94 \pm 0.18) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² AAIJ 17BP also set the limit range $4-25 \times 10^{-7}$ at 90% CL using the world average value $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = (4.96 \pm 0.20) \times 10^{-5}$.

³ Used $f_s/f_d = 0.259 \pm 0.015$.

⁴ AAIJ 13BP reports $[\Gamma(B_s^0 \rightarrow K^0 K^+ K^-)/\Gamma_{\text{total}}] / [\Gamma(B^0 \rightarrow K^0 \pi^+ \pi^-)] < 0.068$ which we multiply by our best value $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = 4.94 \times 10^{-5}$.

$\Gamma(\bar{K}^*(892)^0 \rho^0)/\Gamma_{\text{total}}$ Γ_{111}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.67 \times 10^{-4}$	90	1 ABE	00C SLD	$e^+ e^- \rightarrow Z$

¹ ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

$\Gamma(\bar{K}^*(892)^0 K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{112}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
$1.11 \pm 0.26 \pm 0.06$		1 AAIJ	15AF LHCb	$p p$ at 7 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.81 \pm 0.46 \pm 0.56$		2 AAIJ	12F LHCb	Repl. by AAIJ 15AF
<168.1	90	3 ABE	00C SLD	$e^+ e^- \rightarrow Z$

¹ AAIJ 15AF reports $[\Gamma(B_s^0 \rightarrow \bar{K}^*(892)^0 K^*(892)^0)/\Gamma_{\text{total}}] / [\Gamma(B^0 \rightarrow K^*(892)^0 \phi)] = 1.11 \pm 0.22 \pm 0.12 \pm 0.06$ which we multiply by our best value $B(B^0 \rightarrow K^*(892)^0 \phi) = (1.00 \pm 0.05) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Uses $B^0 \rightarrow J/\psi K^{*0}$ for normalization and assumes $B(B^0 \rightarrow J/\psi K^{*0}) B(J/\psi \rightarrow \mu^+ \mu^-) B(K^{*0} \rightarrow K^+ \pi^-) = (1.33 \pm 0.06) \times 10^{-3}$ and $f_s/f_d = 0.253 \pm 0.031$. The second quoted error is total uncertainty including the error of 0.34 on f_s/f_d .

³ ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

$\Gamma(\phi K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{116}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$1.14 \pm 0.29 \pm 0.06$		1 AAIJ	13BW LHCb	$p p$ at 7 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1013	90	2 ABE	00C SLD	$e^+ e^- \rightarrow Z$
---------	----	-------	---------	-------------------------

¹ AAIJ 13BW reports $[\Gamma(B_s^0 \rightarrow \phi K^*(892)^0)/\Gamma_{\text{total}}] / [\Gamma(B^0 \rightarrow K^*(892)^0 \phi)] = 0.113 \pm 0.024 \pm 0.016$ which we multiply by our best value $B(B^0 \rightarrow K^*(892)^0 \phi) = (1.00 \pm 0.05) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{117}/Γ Test for $\Delta B=1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-8})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.5	90	1 AAIJ	17BJ LHCb	$p p$ at 7 and 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.84 $^{+2.03}_{-1.68}$ $^{+0.85}_{-0.18}$	2 AAIJ	13BQ LHCb	Repl. by AAIJ 17BJ	
<5900	90	3 BUSKULIC	96V ALEP	$e^+ e^- \rightarrow Z$

¹ Uses normalization mode $B(B^0 \rightarrow K^+ \pi^-) = (19.6 \pm 0.5) \times 10^{-6}$ and B production ratio $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.259 \pm 0.015$.

² Uses normalization mode $B(B^0 \rightarrow K^+ \pi^-) = (19.55 \pm 0.54) \times 10^{-6}$ and B production ratio $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.256 \pm 0.020$.

³ BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

 $\Gamma(p\bar{p}K^+K^-)/\Gamma_{\text{total}}$ Γ_{118}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
4.5$\pm 0.4 \pm 0.2$	1,2 AAIJ	17BD LHCb	$p p$ at 7, 8 TeV

¹ AAIJ 17BD reports $[\Gamma(B_s^0 \rightarrow p\bar{p}K^+K^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)K^*(892)^0)] / [B(J/\psi(1S) \rightarrow p\bar{p})] / [B(K^*(892) \rightarrow (K\pi)^\pm)] = 1.67 \pm 0.12 \pm 0.11$ which we multiply by our best values $B(B^0 \rightarrow J/\psi(1S)K^*(892)^0) = (1.27 \pm 0.05) \times 10^{-3}$, $B(J/\psi(1S) \rightarrow p\bar{p}) = (2.121 \pm 0.029) \times 10^{-3}$, $B(K^*(892) \rightarrow (K\pi)^\pm) = (99.901 \pm 0.009) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values. Reported value assumes $f_s/f_d = 0.259 \pm 0.015$.

² The branching ratio is given for $m_{p\bar{p}} < 2.85$ GeV.

 $\Gamma(p\bar{p}K^+\pi^-)/\Gamma_{\text{total}}$ Γ_{119}/Γ

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
13.9$\pm 2.5 \pm 0.5$	1,2 AAIJ	17BD LHCb	$p p$ at 7, 8 TeV

¹ AAIJ 17BD reports $[\Gamma(B_s^0 \rightarrow p\bar{p}K^+\pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)K^*(892)^0)] / [B(J/\psi(1S) \rightarrow p\bar{p})] / [B(K^*(892) \rightarrow (K\pi)^\pm)] = 0.52 \pm 0.08 \pm 0.05$ which we multiply by our best values $B(B^0 \rightarrow J/\psi(1S)K^*(892)^0) = (1.27 \pm 0.05) \times 10^{-3}$, $B(J/\psi(1S) \rightarrow p\bar{p}) = (2.121 \pm 0.029) \times 10^{-3}$, $B(K^*(892) \rightarrow (K\pi)^\pm) = (99.901 \pm 0.009) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values. Reported value assumes $f_s/f_d = 0.259 \pm 0.015$.

² The branching ratio is given for $m_{p\bar{p}} < 2.85$ GeV.

 $\Gamma(p\bar{p}K^+\pi^-)/\Gamma(p\bar{p}K^+K^-)$ $\Gamma_{119}/\Gamma_{118}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.31$\pm 0.05 \pm 0.02$	1,2 AAIJ	17BD LHCb	$p p$ at 7, 8 TeV

¹ Reports $B(B_s^0 \rightarrow p\bar{p}K^+\pi^-) / B(B^0 \rightarrow p\bar{p}K^+\pi^-) = 0.22 \pm 0.04 \pm 0.02 \pm 0.01$, where the third error is due to f_s/f_d .

² The ratio is given for $m_{p\bar{p}} < 2.85$ GeV and assuming $f_s/f_d = 0.259 \pm 0.015$.

$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_{120}/Γ
$\text{VALUE (units } 10^{-7})$	DOCUMENT ID TECN COMMENT
$4.3 \pm 2.0 \pm 0.2$	1,2 AAIJ 17BD LHCb $p\bar{p}$ at 7, 8 TeV

¹ AAIJ 17BD reports $[\Gamma(B_s^0 \rightarrow p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}] / [\text{B}(B^0 \rightarrow J/\psi(1S)K^*(892)^0)] / [\text{B}(J/\psi(1S) \rightarrow p\bar{p})] / [\text{B}(K^*(892) \rightarrow (K\pi)^\pm)] = 0.16 \pm 0.07 \pm 0.02$ which we multiply by our best values $\text{B}(B^0 \rightarrow J/\psi(1S)K^*(892)^0) = (1.27 \pm 0.05) \times 10^{-3}$, $\text{B}(J/\psi(1S) \rightarrow p\bar{p}) = (2.121 \pm 0.029) \times 10^{-3}$, $\text{B}(K^*(892) \rightarrow (K\pi)^\pm) = (99.901 \pm 0.009) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values. Reported value assumes $f_s/f_d = 0.259 \pm 0.015$.

² The branching ratio is given for $m_{p\bar{p}} < 2.85$ GeV.

$\Gamma(p\bar{\Lambda}K^- + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{121}/Γ
$\text{VALUE (units } 10^{-6})$	DOCUMENT ID TECN COMMENT
$5.5 \pm 0.6 \pm 0.8$	1,2 AAIJ 17AL LHCb $p\bar{p}$ at 7, 8 TeV
	¹ AAIJ 17AL reports $(5.46 \pm 0.61 \pm 0.82) \times 10^{-6}$ from a measurement of $[\Gamma(B_s^0 \rightarrow p\bar{\Lambda}K^- + \text{c.c.})/\Gamma_{\text{total}}] / [\text{B}(B^0 \rightarrow p\bar{\Lambda}\pi^-)]$ assuming $\text{B}(B^0 \rightarrow p\bar{\Lambda}\pi^-) = (3.14 \pm 0.29) \times 10^{-6}$.
	² AAIJ 17AL value represents the sum of $B_s^0 \rightarrow p\bar{\Lambda}K^-$ and $B_s^0 \rightarrow \bar{p}\Lambda K^+$ and assumes the fraction $f_s/f_d = 0.259 \pm 0.015$.

$\Gamma(\Lambda_c^- \Lambda\pi^+)/\Gamma_{\text{total}}$	Γ_{122}/Γ
$\text{VALUE (units } 10^{-4})$	DOCUMENT ID TECN COMMENT
$3.6 \pm 1.1 \pm 1.2$	1 SOLOVIEVA 13 BELL $e^+e^- \rightarrow \gamma(4S)$

¹ The second error is the total systematic uncertainty including the Λ_c absolute branching fractions and the normalization number of B_s events.

$\Gamma(\Lambda_c^- \Lambda_c^+)/\Gamma_{\text{total}}$	Γ_{123}/Γ
VALUE	CL\%
$< 8.0 \times 10^{-5}$	95

¹ Uses $\text{B}(\bar{B}^0 \rightarrow D^+ D_s^-) = (7.2 \pm 0.8) \times 10^{-3}$.

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	Γ_{124}/Γ
Test for $\Delta B=1$ weak neutral current.	
$\text{VALUE (units } 10^{-6})$	CL\%
< 3.1	90

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 8.7	90	² WICHT	08A	BELL	Repl. by DUTTA 15
< 53	90	DRUTSKOY	07A	BELL	Repl. by WICHT 08A
< 148	90	³ ACCIARRI	95I	L3	$e^+e^- \rightarrow Z$

¹ Assumes the fraction of $B_s^{(*)}\bar{B}_s^{(*)}$ in $b\bar{b}$ events is $f_s = (17.2 \pm 3.0)\%$.

² Assumes $\gamma(5S) \rightarrow B_s^*\bar{B}_s^* = (19.5^{+3.0}_{-2.3})\%$.

³ ACCIARRI 95I assumes $f_{B_s^0} = 39.5 \pm 4.0$ and $f_{B_s} = (12.0 \pm 3.0)\%$.

$\Gamma(\phi\gamma)/\Gamma_{\text{total}}$		Γ_{125}/Γ		
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
34 ± 4 OUR AVERAGE				
36 ± 5 ± 7		¹ DUTTA	15	BELL $e^+ e^- \rightarrow \gamma(5S)$
33.8 ± 3.4 ± 2.0		² AAIJ	13	LHCb $p\bar{p}$ at 7 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
39 ± 5		³ AAIJ	12AE	LHCb Repl. by AAIJ 13
57 $^{+18}_{-15}$ $^{+12}_{-11}$		⁴ WICHT	08A	BELL Repl. by DUTTA 15
<390	90	DRUTSKOY	07A	BELL $e^+ e^- \rightarrow \gamma(5S)$
<120	90	ACOSTA	02G	CDF $p\bar{p}$ at 1.8 TeV
<700	90	⁵ ADAM	96D	DLPH $e^+ e^- \rightarrow Z$
¹ Assumes the fraction of $B_s^{(*)}\overline{B}_s^{(*)}$ in $b\bar{b}$ events is $f_s = (17.2 \pm 3.0)\%$. The systematic uncertainty from f_s is 0.6×10^{-5} . ² AAIJ 13 reports $[\Gamma(B_s^0 \rightarrow \phi\gamma)/\Gamma_{\text{total}}] / [B(B_s^0 \rightarrow K^*(892)^0\gamma)] = 0.81 \pm 0.04 \pm 0.07$ which we multiply by our best value $B(B_s^0 \rightarrow K^*(892)^0\gamma) = (4.18 \pm 0.25) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. ³ Measures $B(B_s^0 \rightarrow K^{*0}\gamma)/B(B_s \rightarrow \phi\gamma) = 1.12 \pm 0.08(\text{stat})^{+0.06}_{-0.04}(\text{sys})^{+0.09}_{-0.08}(f_s/f_d)$ and uses current world-average value of $B(B_s^0 \rightarrow K^{*0}\gamma) = (4.33 \pm 0.15) \times 10^{-5}$. ⁴ Assumes $\gamma(5S) \rightarrow B_s^*\overline{B}_s^* = (19.5^{+3.0}_{-2.3})\%$. ⁵ ADAM 96D assumes $f_{B_s^0} = f_{B_s^-} = 0.39$ and $f_{B_s} = 0.12$.				

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$		Γ_{126}/Γ		
VALUE (units 10^{-9})	CL%	DOCUMENT ID	TECN	COMMENT
3.0±0.4 OUR AVERAGE				
2.8 $^{+0.8}_{-0.7}$		AABOUD	19L	ATLS $p\bar{p}$ at 7, 8, 13 TeV
3.0 ± 0.6 $^{+0.3}_{-0.2}$		AAIJ	17AI	LHCb $p\bar{p}$ at 7, 8, 13 TeV
13 $^{+9}_{-7}$		¹ AALTONEN	13F	CDF $p\bar{p}$ at 1.96 TeV
3.0 $^{+1.0}_{-0.9}$		² CHATRCHYAN	13AW	CMS $p\bar{p}$ at 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.9 $^{+1.1}_{-0.8}$		³ AABOUD	16L	ATLS $p\bar{p}$ at 7, 8 TeV, Repl. by AABOUD 16L
2.8 $^{+0.7}_{-0.6}$		⁴ KHACHATRYAN	15BE	LHCb $p\bar{p}$ at 7, 8 TeV
3.2 $^{+1.4}_{-1.2}$ $^{+0.5}_{-0.3}$		⁵ AAIJ	13B	LHCb Repl. by AAIJ 13BA
2.9 $^{+1.1}_{-1.0}$ $^{+0.3}_{-0.1}$		⁶ AAIJ	13BA	LHCb Repl. by KHACHATRYAN 15BE
<12	90	⁷ ABAZOV	13C	D0 $p\bar{p}$ at 1.96 TeV
<19	90	⁸ AAD	12AE	ATLS $p\bar{p}$ at 7 TeV
<12	90	⁹ AAIJ	12A	LHCb Repl. by AAIJ 12W
< 3.8	90	¹⁰ AAIJ	12W	LHCb Repl. by AAIJ 13B
< 6.4	90	¹¹ CHATRCHYAN	12A	CMS $p\bar{p}$ at 7 TeV
<43	90	¹² AAIJ	11B	LHCb Repl. by AAIJ 12A
<35	90	¹³ AALTONEN	11AG	CDF $p\bar{p}$ at 1.96 TeV

<16	90	¹⁴ CHATRCHYAN 11T CMS	Repl. by CHATRCHYAN 12A
<42	90	¹⁵ ABAZOV 10S D0	$p\bar{p}$ at 1.96 TeV
¹ Uses normalization mode $B(B^+ \rightarrow J/\psi K^+) = (10.22 \pm 0.35) \times 10^{-4}$ and B production ratio $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.28 \pm 0.04$.			
² Uses B production ratio $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.256 \pm 0.020$ and $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (6.0 \pm 0.2) \times 10^{-5}$ for normalization.			
³ This value corresponds to an upper limit of $< 3.0 \times 10^{-9}$ at 95% C.L. It uses $f_s/f_d = 0.24 \pm 0.02$.			
⁴ Determined from the joint fit to CMS and LHCb data. Uncertainty includes both statistical and systematic component.			
⁵ Uses B production ratio $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.256 \pm 0.020$ and two normalization modes: $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (6.01 \pm 0.21) \times 10^{-5}$ and $B(B^0 \rightarrow K^+ \pi^-) = (1.94 \pm 0.06) \times 10^{-5}$.			
⁶ Uses B production ratio $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.259 \pm 0.015$ and normalization modes $B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+$ and $B^0 \rightarrow K^+ \pi^-$.			
⁷ Uses normalization mode $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (6.01 \pm 0.21) \times 10^{-5}$ and B production ratio $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.263 \pm 0.017$.			
⁸ Uses B production ratio $f(\bar{b} \rightarrow B^+)/f(\bar{b} \rightarrow B_s^0) = 3.75 \pm 0.29$ and $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (6.0 \pm 0.2) \times 10^{-5}$.			
⁹ Uses B production ratio $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.267^{+0.021}_{-0.020}$ and three normalization modes $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (6.01 \pm 0.21) \times 10^{-5}$, $B(B^0 \rightarrow K^+ \pi^-) = (1.94 \pm 0.06) \times 10^{-5}$, and $B(B_s^0 \rightarrow J/\psi \phi \rightarrow \mu^+ \mu^- K^+ K^-) = (3.4 \pm 0.9) \times 10^{-5}$.			
¹⁰ Uses B production ratio $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.267^{+0.021}_{-0.020}$ and three normalization modes of $B^+ \rightarrow J/\psi K^+$, $B^0 \rightarrow K^+ \pi^-$, and $B_s^0 \rightarrow J/\psi \phi$.			
¹¹ Uses $f_s/f_u = 0.267 \pm 0.021$ and $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (6.0 \pm 0.2) \times 10^{-5}$.			
¹² Uses B production ratio $f(\bar{b} \rightarrow B^+)/f(\bar{b} \rightarrow B_s^0) = 3.71 \pm 0.47$ and three normalization modes.			
¹³ Uses B production ratio $f(\bar{b} \rightarrow B^+)/f(\bar{b} \rightarrow B_s^0) = 3.55 \pm 0.47$ and $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (6.01 \pm 0.21) \times 10^{-5}$.			
¹⁴ Uses B production ratio $f(\bar{b} \rightarrow B^+)/f(\bar{b} \rightarrow B_s^0) = 3.55 \pm 0.42$ and $B(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+) = (6.0 \pm 0.2) \times 10^{-5}$.			
¹⁵ Uses B production ratio $f(\bar{b} \rightarrow B^+)/f(\bar{b} \rightarrow B_s^0) = 3.86 \pm 0.59$, and the number of $B^+ \rightarrow J/\psi K^+$ decays.			

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_{127}/Γ Test for $\Delta B = 1$ weak neutral current.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.8 \times 10^{-7}$	90	AALTONEN 09P	CDF	$p\bar{p}$ at 1.96 TeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<5.4 \times 10^{-5}$	90	¹ ACCIARRI 97B	L3	$e^+ e^- \rightarrow Z$

¹ ACCIARRI 97B assume PDG 96 production fractions for B^+ , B^0 , B_s^0 , and Λ_b . $\Gamma(\tau^+ \tau^-)/\Gamma_{\text{total}}$ Γ_{128}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.8 \times 10^{-3}$	95	¹ AAIJ	17AJ LHCb	$p\bar{p}$ at 7, 8 TeV

¹ Assuming no contribution from $B^0 \rightarrow \tau^+ \tau^-$.

$\Gamma(\mu^+\mu^-\mu^+\mu^-)/\Gamma_{\text{total}}$		Γ_{129}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.5 \times 10^{-9}$	95	AAIJ	17N LHCb	$p\bar{p}$ at 7, 8 TeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<1.2 \times 10^{-8}$	90	¹ AAIJ	13AW LHCb	Repl. by AAIJ 17N

¹ Also reports a limit of $< 1.6 \times 10^{-8}$ at 95% CL.

$\Gamma(SP, S \rightarrow \mu^+\mu^-, P \rightarrow \mu^+\mu^-)/\Gamma_{\text{total}}$		Γ_{130}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.2 \times 10^{-9}$	95	AAIJ	17N LHCb	$p\bar{p}$ at 7, 8 TeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<1.2 \times 10^{-8}$	90	¹ AAIJ	13AW LHCb	Repl. by AAIJ 17N

¹ Also reports a limit of $< 1.6 \times 10^{-8}$ at 95% CL.

$\Gamma(\phi(1020)\mu^+\mu^-)/\Gamma_{\text{total}}$		Γ_{131}/Γ		
VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<32	90	¹ ABAZOV	06G D0	$p\bar{p}$ at 1.96 TeV
$< 4.7 \times 10^2$	90	ACOSTA	02D CDF	$p\bar{p}$ at 1.8 TeV

¹ Uses $B(B_s^0 \rightarrow J/\psi \phi) = 9.3 \times 10^{-4}$.

$\Gamma(\phi(1020)\mu^+\mu^-)/\Gamma(J/\psi(1S)\phi)$		Γ_{131}/Γ_{49}		
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
0.76 ± 0.09 OUR AVERAGE Error includes scale factor of 1.9.				
0.741 ^{+0.042} _{-0.040} ± 0.029	AAIJ	15AQ LHCb	$p\bar{p}$ at 7, 8 TeV	
1.13 ± 0.19 ± 0.07	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.674 ^{+0.061} _{-0.056} ± 0.016	¹ AAIJ	13X LHCb	Repl. by AAIJ 15AQ	
1.11 ± 0.25 ± 0.09	AALTONEN	11L CDF	Repl. by AALTONEN 11AI	
< 2.3	90	AALTONEN	09B CDF	Repl. by AALTONEN 11L

¹ Replaced by AAIJ 15AQ.

$\Gamma(\overline{K}^*(892)^0\mu^+\mu^-)/\Gamma_{\text{total}}$		Γ_{132}/Γ		
VALUE (units 10^{-8})	CL%	DOCUMENT ID	TECN	COMMENT
2.9 ± 1.0 ± 0.4	¹ AAIJ	18AB LHCb	$p\bar{p}$ at 7, 8, 13 TeV	
$\bullet \bullet \bullet$ Normalizes to $B(B^0 \rightarrow J/\psi K^{*0}) = 1.19 \pm 0.01 \pm 0.08\%$ and $B(J/\psi \rightarrow \mu^+\mu^-) = 5.96 \pm 0.03\%$, and uses $f_s/f_d = 0.259 \pm 0.015$.				

$\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{133}/Γ

VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT
$8.4 \pm 1.6 \pm 0.3$	¹ AAIJ	15S	LHCb $p\bar{p}$ at 7, 8 TeV

¹ AAIJ 15S reports $(8.6 \pm 1.5 \pm 0.7 \pm 0.7) \times 10^{-8}$ from a measurement of $[\Gamma(B_s^0 \rightarrow \pi^+\pi^-\mu^+\mu^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)K^*(892)^0)]$ assuming $B(B^0 \rightarrow J/\psi(1S)K^*(892)^0) = (1.3 \pm 0.1) \times 10^{-3}$, which we rescale to our best value $B(B^0 \rightarrow J/\psi(1S)K^*(892)^0) = (1.27 \pm 0.05) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\phi\nu\bar{\nu})/\Gamma_{\text{total}}$ Γ_{134}/Γ Test for $\Delta B = 1$ weak neutral current.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.4 \times 10^{-3}$	90	¹ ADAM	96D	DLPH $e^+e^- \rightarrow Z$

¹ ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

 $\Gamma(e^\pm\mu^\mp)/\Gamma_{\text{total}}$ Γ_{135}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.4 \times 10^{-9}$	90	¹ AAIJ	18T	LHCb $p\bar{p}$ at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.1 \times 10^{-8}$	90	² AAIJ	13BMLHCB	Repl. by AAIJ 18T
$<2.0 \times 10^{-7}$	90	AALTONEN	09P	CDF $p\bar{p}$ at 1.96 TeV
$<6.1 \times 10^{-6}$	90	ABE	98V	CDF Repl. by AALTONEN 09P
$<4.1 \times 10^{-5}$	90	³ ACCIARRI	97B	$L3 e^+e^- \rightarrow Z$

¹ AAIJ 18T uses normalization modes $B(B^0 \rightarrow K^+\pi^-) = (19.6 \pm 0.5) \times 10^{-6}$ and $B(B^+ \rightarrow J/\psi K^+) = (1.026 \pm 0.031) \times 10^{-3}$ with B production ratio $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.259 \pm 0.015$. The upper limit increases to 6×10^{-9} with the assumption of B_L -dominated decay amplitude.

² Uses normalization mode $B(B^0 \rightarrow K^+\pi^-) = (19.4 \pm 0.6) \times 10^{-6}$ and B production ratio $f(\bar{b} \rightarrow B_s^0)/f(\bar{b} \rightarrow B_d^0) = 0.256 \pm 0.020$.

³ ACCIARRI 97B assume PDG 96 production fractions for B^+ , B^0 , B_s , and Λ_b .

POLARIZATION IN B_s^0 DECAY

In decays involving two vector mesons, one can distinguish among the states in which meson polarizations are both longitudinal (L), or both are transverse and parallel (\parallel), or perpendicular (\perp) to each other with the parameters Γ_L/Γ , Γ_\perp/Γ , and the relative phases ϕ_\parallel and ϕ_\perp . In decays involving two tensor mesons, the transverse polarization states are described by parameters $\Gamma_{\parallel 1}$, $\Gamma_{\parallel 2}$, $\Gamma_{\perp 1}$, $\Gamma_{\perp 2}$ and their relative phases $\phi_{\parallel 1}$, $\phi_{\parallel 2}$, $\phi_{\perp 1}$, $\phi_{\perp 2}$. See also the review on “Polarization in B Decays.”

 Γ_L/Γ in $B_s^0 \rightarrow D_s^*\rho^+$ Γ_{136}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$1.05^{+0.08+0.03}_{-0.10-0.04}$	LOUVOT	10	BELL $e^+e^- \rightarrow \gamma(5S)$

Γ_L/Γ in $B_s^0 \rightarrow J/\psi(1S)\phi$

VALUE		DOCUMENT ID	TECN	COMMENT
0.523 ± 0.005 OUR AVERAGE				Error includes scale factor of 1.1.
0.522 ± 0.003 ± 0.007	¹ AAD	16AP ATLS	$p\bar{p}$ at 7, 8 TeV	
0.510 ± 0.005 ± 0.011	KHACHATRY...16S	CMS	$p\bar{p}$ at 8 TeV	
0.5241 ± 0.0034 ± 0.0067	AAIJ	15I LHCb	$p\bar{p}$ at 7, 8 TeV	
0.524 ± 0.013 ± 0.015	² AALTONEN	12D CDF	$p\bar{p}$ at 1.96 TeV	
0.558 +0.017 -0.019	^{2,3} ABAZOV	12D D0	$p\bar{p}$ at 1.96 TeV	
0.61 ± 0.14 ± 0.02	⁴ AFFOLDER	00N CDF	$p\bar{p}$ at 1.8 TeV	
0.56 ± 0.21 ± 0.02 -0.04	ABE	95Z CDF	$p\bar{p}$ at 1.8 TeV	

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE		DOCUMENT ID	TECN	COMMENT
0.529 ± 0.006 ± 0.012	¹ AAD	14U ATLS	Repl. by AAD 16AP	
0.539 ± 0.014 ± 0.016	² AAD	12CV ATLS	Repl. by AAD 14U	
0.555 ± 0.027 ± 0.006	⁵ ABAZOV	09E D0	Repl. by ABAZOV 12D	
0.531 ± 0.020 ± 0.007	² AALTONEN	08J CDF	Repl. by AALTONEN 12D	
0.62 ± 0.06 ± 0.01	ACOSTA	05 CDF	Repl. by AALTONEN 08J	

¹ Measured using the flavor tagged, time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays.

² Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays.

³ The error includes both statistical and systematic uncertainties.

⁴ AFFOLDER 00N measurements are based on 40 B_s^0 candidates obtained from a data sample of 89 pb^{-1} . The P -wave fraction is found to be $0.23 \pm 0.19 \pm 0.04$.

⁵ Measured the angular and lifetime parameters for the time-dependent angular untagged decays $B_d^0 \rightarrow J/\psi K^{*0}$ and $B_s^0 \rightarrow J/\psi\phi$.

 Γ_L/Γ in $B_s^0 \rightarrow D_s^{*+} D_s^{*-}$

VALUE		DOCUMENT ID	TECN	COMMENT
0.06 +0.18 -0.17 ± 0.03		ESEN	13 BELL	$e^+ e^- \rightarrow \gamma(5S)$

 $\Gamma_{||}/\Gamma$ in $B_s^0 \rightarrow J/\psi(1S)\phi$

VALUE		DOCUMENT ID	TECN	COMMENT
0.228 ± 0.007 OUR AVERAGE				
0.227 ± 0.004 ± 0.006	¹ AAD	16AP ATLS	$p\bar{p}$ at 7, 8 TeV	
0.231 ± 0.014 ± 0.015	² AALTONEN	12D CDF	$p\bar{p}$ at 1.96 TeV	
0.231 +0.024 -0.030	^{2,3} ABAZOV	12D D0	$p\bar{p}$ at 1.96 TeV	

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE		DOCUMENT ID	TECN	COMMENT
0.220 ± 0.008 ± 0.009	¹ AAD	14U ATLS	Repl. by AAD 16AP	
0.224 ± 0.010 ± 0.009	² AAD	12CV ATLS	Repl. by AAD 14U	
0.244 ± 0.032 ± 0.014	⁴ ABAZOV	09E D0	Repl. by ABAZOV 12D	
0.230 ± 0.029 ± 0.011	² AALTONEN	08J CDF	Repl. by AALTONEN 12D	
0.260 ± 0.084 ± 0.013	ACOSTA	05 CDF	Repl. by AALTONEN 08J	

¹ Measured using a tagged, time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays.

² Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays.

³ The error includes both statistical and systematic uncertainties.

⁴ Measured the angular and lifetime parameters for the time-dependent angular untagged decays $B_d^0 \rightarrow J/\psi K^{*0}$ and $B_s^0 \rightarrow J/\psi\phi$.

Γ_{\perp}/Γ in $B_s^0 \rightarrow J/\psi(1S)\phi$ VALUE **0.249 ± 0.006 OUR AVERAGE** $0.243 \pm 0.008 \pm 0.012$ $0.2504 \pm 0.0049 \pm 0.0036$ DOCUMENT ID TECN COMMENTKHACHATRY...16S CMS $p p$ at 8 TeV
AAIJ 15I LHCb $p p$ at 7, 8 TeV ϕ_{\parallel} in $B_s^0 \rightarrow J/\psi(1S)\phi$ VALUE (rad) **$3.19^{+0.08}_{-0.09}$ OUR AVERAGE** $3.15 \pm 0.10 \pm 0.05$ $3.48^{+0.07}_{-0.09} \pm 0.68$ $3.26^{+0.10}_{-0.17} {}^{+0.06}_{-0.07}$ 3.15 ± 0.22 ¹ ABAZOV 12D D0 $p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $2.72^{+1.12}_{-0.27} \pm 0.26$ DOCUMENT ID TECN COMMENTAAD 16AP ATLS $p p$ at 7, 8 TeV
KHACHATRY...16S CMS $p p$ at 8 TeV
AAIJ 15I LHCb $p p$ at 7, 8 TeV
ABAZOV 09E D0 Repl. by ABAZOV 12D¹ The error includes both statistical and systematic uncertainties. ϕ_{\perp} in $B_s^0 \rightarrow J/\psi(1S)\phi$ VALUE (rad) **3.2 ± 0.4 OUR AVERAGE**

Error includes scale factor of 2.8.

 $4.15 \pm 0.32 \pm 0.16$ ¹ AAD 16AP ATLS $p p$ at 7, 8 TeV $2.98 \pm 0.36 \pm 0.66$ KHACHATRY...16S CMS $p p$ at 8 TeV $3.08^{+0.14}_{-0.15} \pm 0.06$ AAIJ 15I LHCb $p p$ at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $3.89 \pm 0.47 \pm 0.11$ ¹ AAD 14U ATLS Repl. by AAD 16AP¹ Measured using a tagged, time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays. Γ_{\perp}/Γ in $B_s^0 \rightarrow \psi(2S)\phi$ VALUEDOCUMENT ID TECN COMMENT **$0.264^{+0.024}_{-0.023} \pm 0.002$** ¹ AAIJ 16AK LHCb $p p$ at 7, 8 TeV¹ Measured using time-dependent angular analysis of $B_s^0 \rightarrow \psi(2S)\phi$ decays. ϕ_{\parallel} in $B_s^0 \rightarrow \psi(2S)\phi$ VALUE (rad)DOCUMENT ID TECN COMMENT **$3.67^{+0.13}_{-0.18} \pm 0.03$** ¹ AAIJ 16AK LHCb $p p$ at 7, 8 TeV¹ Measured using time-dependent angular analysis of $B_s^0 \rightarrow \psi(2S)\phi$ decays. ϕ_{\perp} in $B_s^0 \rightarrow \psi(2S)\phi$ VALUE (rad)DOCUMENT ID TECN COMMENT **$3.29^{+0.43}_{-0.39} \pm 0.04$** ¹ AAIJ 16AK LHCb $p p$ at 7, 8 TeV¹ Measured using time-dependent angular analysis of $B_s^0 \rightarrow \psi(2S)\phi$ decays.

Γ_L/Γ for $B_s^0 \rightarrow J/\psi(1S)\bar{K}^*(892)^0$

Longitudinal polarization fraction, equals to f_L using notation of “Polarization in B decays” review.

VALUE	DOCUMENT ID	TECN	COMMENT
0.497±0.025±0.025	AAIJ	15AV LHCb	$p\bar{p}$ at 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			

0.50 ± 0.08 ± 0.02 ¹ AAIJ 12AP LHCb Repl. by AAIJ 15AV

¹ The non-resonant $K\pi$ background contributions are subtracted. Also reports an S -wave amplitude $|A_S|^2 = 0.07^{+0.15}_{-0.07}$.

 $\Gamma_{||}/\Gamma$ for $B_s^0 \rightarrow J/\psi(1S)\bar{K}^*(892)^0$

Parallel polarization fraction, equals to $1 - f_L - f_\perp$ using notation of “Polarization in B decays” review.

VALUE	DOCUMENT ID	TECN	COMMENT
0.179±0.027±0.013	AAIJ	15AV LHCb	$p\bar{p}$ at 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			

0.19 ^{+0.10} _{-0.08} ± 0.02 ¹ AAIJ 12AP LHCb Repl. by AAIJ 15AV

¹ The non-resonant $K\pi$ background contributions are subtracted. Also reports an S -wave amplitude $|A_S|^2 = 0.07^{+0.15}_{-0.07}$.

 $\Gamma_{||}/\Gamma$ of $K^*(892)^0$ in $B_s^0 \rightarrow \psi(2S)\bar{K}^*(892)^0$

VALUE	DOCUMENT ID	TECN	COMMENT
0.524±0.056±0.029	AAIJ	15U LHCb	$p\bar{p}$ at 7, 8 TeV

 Γ_L/Γ in $B_s^0 \rightarrow \phi\phi$

VALUE	DOCUMENT ID	TECN	COMMENT
0.362±0.014 OUR AVERAGE			
0.364±0.012±0.009	AAIJ	14AE LHCb	$p\bar{p}$ at 7, 8 TeV
0.348±0.041±0.021	AALTONEN	11AN CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.365±0.022±0.012	AAIJ	12P LHCb	Repl. by AAIJ 14AE

 Γ_\perp/Γ in $B_s^0 \rightarrow \phi\phi$

VALUE	DOCUMENT ID	TECN	COMMENT
0.309±0.015 OUR AVERAGE			Error includes scale factor of 1.1.
0.305±0.013±0.005	AAIJ	14AE LHCb	$p\bar{p}$ at 7, 8 TeV
0.365±0.044±0.027	AALTONEN	11AN CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.291±0.024±0.010	AAIJ	12P LHCb	Repl. by AAIJ 14AE

 $\phi_{||}$ in $B_s^0 \rightarrow \phi\phi$

VALUE (rad)	DOCUMENT ID	TECN	COMMENT
2.55±0.11 OUR AVERAGE			
2.54±0.07±0.09	¹ AAIJ	14AE LHCb	$p\bar{p}$ at 7, 8 TeV
2.71 ^{+0.31} _{-0.36} ± 0.22	² AALTONEN	11AN CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.57±0.15±0.06	³ AAIJ	12P LHCb	Repl. by AAIJ 14AE

- ¹ AAIJ 14AE reports measurement of ϕ_{\perp} and $\phi_{\perp} - \phi_{\parallel}$, which we convert into ϕ_{\parallel} . Statistical uncertainty includes correlation between measured parameters, while systematic uncertainties are assumed uncorrelated.
² AALTONEN 11AN quotes $\cos\phi_{\parallel} = -0.91^{+0.15}_{-0.13} \pm 0.09$ which we convert to ϕ_{\parallel} taking the smaller solution.
³ AAIJ 12P quotes $\cos\phi_{\parallel} = -0.844 \pm 0.068 \pm 0.029$ which we convert to ϕ_{\parallel} , taking the smaller solution.

 ϕ_{\perp} in $B_s^0 \rightarrow \phi\phi$

VALUE (rad)	DOCUMENT ID	TECN	COMMENT
2.67±0.23±0.07	AAIJ	14AE	LHCb $p p$ at 7, 8 TeV

 Γ_L/Γ in $B_s^0 \rightarrow K^{*0}\bar{K}^{*0}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.208±0.032±0.046	1 AAIJ	18S	LHCb $p p$ at 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.201±0.057±0.040	2 AAIJ	15AF	LHCb Repl. by AAIJ 18S
0.31 ± 0.12 ± 0.04	AAIJ	12F	LHCb Repl. by AAIJ 15AF

¹ Measured in angular analysis, which takes into account S-, P- and D-wave contributions.

² Measured in angular analysis, which takes into account S-wave contributions.

 Γ_{\perp}/Γ in $B_s^0 \rightarrow K^{*0}\bar{K}^{*0}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.38±0.11±0.04	AAIJ	12F	LHCb $p p$ at 7 TeV

 $\Gamma_{\parallel}/\Gamma$ in $B_s^0 \rightarrow K^*(892)^0\bar{K}^*(892)^0$

VALUE	DOCUMENT ID	TECN	COMMENT
0.297±0.029±0.042	1 AAIJ	18S	LHCb $p p$ at 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.215±0.046±0.015	AAIJ	15AF	LHCb Repl. by AAIJ 18S

¹ Measured in angular analysis, which takes into account S-, P- and D-wave contributions.

 Φ_{\parallel} in $B_s^0 \rightarrow K^*(892)^0\bar{K}^*(892)^0$

VALUE	DOCUMENT ID	TECN	COMMENT
2.40±0.11±0.33	1 AAIJ	18S	LHCb $p p$ at 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5.31±0.24±0.14	AAIJ	15AF	LHCb Repl. by AAIJ 18S

¹ Measured in angular analysis, which takes into account S-, P- and D-wave contributions.

 Φ_{\perp} in $B_s^0 \rightarrow K^*(892)^0\bar{K}^*(892)^0$

VALUE (rad)	DOCUMENT ID	TECN	COMMENT
2.62±0.26±0.64	1 AAIJ	18S	LHCb $p p$ at 7, 8 TeV

¹ Measured in angular analysis, which takes into account S-, P- and D-wave contributions.

 Γ_L/Γ in $B_s^0 \rightarrow \phi\bar{K}^{*0}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.51±0.15±0.07	AAIJ	13BW	LHCb $p p$ at 7 TeV

$\Gamma_{||}/\Gamma \text{ in } B_s^0 \rightarrow \phi \bar{K}^{*0}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.21±0.11±0.02	AAIJ	13BW	LHCb $p p$ at 7 TeV

 $\phi_{||} \text{ in } B_s^0 \rightarrow \phi \bar{K}^{*0}$

VALUE (rad)	DOCUMENT ID	TECN	COMMENT
1.75±0.53±0.29	¹ AAIJ	13BW	LHCb $p p$ at 7 TeV

¹ Measures $\cos(\phi_{||}) = -0.18 \pm 0.52 \pm 0.29$, which we convert to $\phi_{||}$ by taking the smaller solution.

 $\Gamma_L/\Gamma \text{ in } B_s^0 \rightarrow \bar{D}^{*0} \phi$

VALUE	DOCUMENT ID	TECN	COMMENT
0.73±0.15±0.04	AAIJ	18AY	LHCb $p p$ at 7 and 8 TeV

 $\Gamma_L/\Gamma \text{ in } B_s^0 \rightarrow K^*(892)^0 \bar{K}_2^*(1430)^0$

VALUE	DOCUMENT ID	TECN	COMMENT
0.911±0.020±0.165	¹ AAIJ	18S	LHCb $p p$ at 7, 8 TeV

¹ Measured in angular analysis, which takes into account S-, P- and D-wave. contributions.

 $\Gamma_{||}/\Gamma \text{ in } B_s^0 \rightarrow K^*(892)^0 \bar{K}_2^*(1430)^0$

VALUE	DOCUMENT ID	TECN	COMMENT
0.012±0.008±0.053	¹ AAIJ	18S	LHCb $p p$ at 7, 8 TeV

¹ Measured in angular analysis, which takes into account S-, P- and D-wave. contributions.

 $\Gamma_L/\Gamma \text{ in } B_s^0 \rightarrow K_2^*(1430)^0 \bar{K}^*(892)^0$

VALUE	DOCUMENT ID	TECN	COMMENT
0.62±0.16±0.25	¹ AAIJ	18S	LHCb $p p$ at 7, 8 TeV

¹ Measured in angular analysis, which takes into account S-, P- and D-wave. contributions.

 $\Gamma_{||}/\Gamma \text{ in } B_s^0 \rightarrow K_2^*(1430)^0 \bar{K}^*(892)^0$

VALUE	DOCUMENT ID	TECN	COMMENT
0.24±0.10±0.14	¹ AAIJ	18S	LHCb $p p$ at 7, 8 TeV

¹ Measured in angular analysis, which takes into account S-, P- and D-wave. contributions.

 $\Gamma_L/\Gamma \text{ in } B_s^0 \rightarrow K_2^*(1430)^0 \bar{K}_2^*(1430)^0$

VALUE	DOCUMENT ID	TECN	COMMENT
0.25±0.14±0.18	¹ AAIJ	18S	LHCb $p p$ at 7, 8 TeV

¹ Measured in angular analysis, which takes into account S-, P- and D-wave. contributions.

 $\Gamma_{||1}/\Gamma \text{ in } B_s^0 \rightarrow K_2^*(1430)^0 \bar{K}_2^*(1430)^0$

VALUE	DOCUMENT ID	TECN	COMMENT
0.17±0.11±0.14	¹ AAIJ	18S	LHCb $p p$ at 7, 8 TeV

¹ Measured in angular analysis, which takes into account S-, P- and D-wave. contributions.

$\Gamma_{\perp 1}/\Gamma$ in $B_s^0 \rightarrow K_2^*(1430)^0 \bar{K}_2^*(1430)^0$

VALUE	DOCUMENT ID	TECN	COMMENT
0.30±0.18±0.21	¹ AAIJ	18S LHCb	$p p$ at 7, 8 TeV

¹ Measured in angular analysis, which takes into account S -, P - and D -wave. contributions.

 $\Gamma_{|| 2}/\Gamma$ in $B_s^0 \rightarrow K_2^*(1430)^0 \bar{K}_2^*(1430)^0$

VALUE	DOCUMENT ID	TECN	COMMENT
0.015±0.033±0.107	¹ AAIJ	18S LHCb	$p p$ at 7, 8 TeV

¹ Measured in angular analysis, which takes into account S -, P - and D -wave . contributions.

 $F_L(B_s^0 \rightarrow \phi \mu^+ \mu^-)$ ($0.10 < q^2 < 2.00 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
0.20^{+0.08}_{-0.09}±0.02	AAIJ	15AQ LHCb	$p p$ at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.37^{+0.19}_{-0.17} \pm 0.07$	AAIJ	13X LHCb	Repl. by AAIJ 15AQ
---------------------------------	------	----------	--------------------

 $F_L(B_s^0 \rightarrow \phi \mu^+ \mu^-)$ ($2.00 < q^2 < 5.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
0.68^{+0.16}_{-0.13}±0.03	AAIJ	15AQ LHCb	$p p$ at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.53^{+0.25}_{-0.23} \pm 0.10$	¹ AAIJ	13X LHCb	Repl. by AAIJ 15AQ
---------------------------------	-------------------	----------	--------------------

¹ Measured in $2.0 < q^2 < 4.3 \text{ GeV}^2/c^4$.

 $F_L(B_s^0 \rightarrow \phi \mu^+ \mu^-)$ ($5.0 < q^2 < 8.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
0.54^{+0.10}_{-0.09}±0.02	AAIJ	15AQ LHCb	$p p$ at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.81^{+0.11}_{-0.13} \pm 0.05$	¹ AAIJ	13X LHCb	Repl. by AAIJ 15AQ
---------------------------------	-------------------	----------	--------------------

¹ Measured in $4.3 < q^2 < 8.68 \text{ GeV}^2/c^4$.

 $F_L(B_s^0 \rightarrow \phi \mu^+ \mu^-)$ ($11.0 < q^2 < 12.5 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
0.29±0.11±0.04	AAIJ	15AQ LHCb	$p p$ at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.33^{+0.14}_{-0.12} \pm 0.06$	¹ AAIJ	13X LHCb	Repl. by AAIJ 15AQ
---------------------------------	-------------------	----------	--------------------

¹ Measured in $10.09 < q^2 < 12.90 \text{ GeV}^2/c^4$.

$F_L(B_s^0 \rightarrow \phi\mu^+\mu^-)$ ($15.0 < q^2 < 17.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.23^{+0.09}_{-0.08} \pm 0.02$	AAIJ	15AQ LHCb	$p p$ at 7, 8 TeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$0.34^{+0.18}_{-0.17} \pm 0.07$	¹ AAIJ	13X LHCb	Repl. by AAIJ 15AQ

¹ Measured in $14.18 < q^2 < 16 \text{ GeV}^2/c^4$.

$F_L(B_s^0 \rightarrow \phi\mu^+\mu^-)$ ($17.0 < q^2 < 19.0 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.40^{+0.13}_{-0.15} \pm 0.02$	AAIJ	15AQ LHCb	$p p$ at 7, 8 TeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$0.16^{+0.17}_{-0.10} \pm 0.07$	¹ AAIJ	13X LHCb	Repl. by AAIJ 15AQ

¹ Measured in $16.0 < q^2 < 19.0 \text{ GeV}^2/c^4$.

$F_L(B_s^0 \rightarrow \phi\mu^+\mu^-)$ ($1.00 < q^2 < 6.00 \text{ GeV}^2/c^4$)

VALUE	DOCUMENT ID	TECN	COMMENT
$0.63^{+0.09}_{-0.09} \pm 0.03$	AAIJ	15AQ LHCb	$p p$ at 7, 8 TeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$0.56^{+0.17}_{-0.16} \pm 0.09$	AAIJ	13X LHCb	Repl. by AAIJ 15AQ

$B_s^0-\overline{B}_s^0$ MIXING

For a discussion of $B_s^0-\overline{B}_s^0$ mixing see the note on “ $B^0-\overline{B}^0$ Mixing” in the B^0 Particle Listings above.

χ_s is a measure of the time-integrated $B_s^0-\overline{B}_s^0$ mixing probability that produced $B_s^0(\overline{B}_s^0)$ decays as a $\overline{B}_s^0(B_s^0)$. Mixing violates $\Delta B \neq 2$ rule.

$$\chi_s = \frac{x_s^2}{2(1+x_s^2)}$$

$$x_s = \frac{\Delta m_{B_s^0}}{\Gamma_{B_s^0}} = (m_{B_{sH}^0} - m_{B_{sL}^0}) \tau_{B_s^0},$$

where H, L stand for heavy and light states of two B_s^0 CP eigenstates and

$$\tau_{B_s^0} = \frac{1}{0.5(\Gamma_{B_{sH}^0} + \Gamma_{B_{sL}^0})}.$$

$$\Delta m_{B_s^0} = m_{B_{sH}^0} - m_{B_{sL}^0}$$

$\Delta m_{B_s^0}$ is a measure of 2π times the B_s^0 - \bar{B}_s^0 oscillation frequency in time-dependent mixing experiments.

"OUR EVALUATION" is provided by the Heavy Flavor Averaging Group (HFLAV) by taking into account correlations between measurements.

VALUE ($10^{12} \text{ } \text{Hz} \text{ s}^{-1}$)	CL%	DOCUMENT ID	TECN	COMMENT
17.757 \pm 0.021 OUR EVALUATION				
17.756 \pm 0.021 OUR AVERAGE				
17.711 $^{+0.055}_{-0.057}$ \pm 0.011	1 AAIJ	15I LHCb	$p\bar{p}$ at 7, 8 TeV	
17.768 \pm 0.023 \pm 0.006	2 AAIJ	13BI LHCb	$p\bar{p}$ at 7 TeV	
17.93 \pm 0.22 \pm 0.15	3 AAIJ	13CF LHCb	$p\bar{p}$ at 7 TeV	
17.63 \pm 0.11 \pm 0.02	4 AAIJ	12I LHCb	$p\bar{p}$ at 7 TeV	
17.77 \pm 0.10 \pm 0.07	5 ABULENCIA,A 06G CDF	$p\bar{p}$ at 1.96 TeV		
• • • We do not use the following data for averages, fits, limits, etc. • • •				
17-21	90	6 ABAZOV	06B D0	$p\bar{p}$ at 1.96 TeV
17.31 $^{+0.33}_{-0.18}$ \pm 0.07	7 ABULENCIA	06Q CDF	Repl. by ABULENCIA,A 06G	
> 8.0	95	8 ABDALLAH	04J DLPH	$e^+e^- \rightarrow Z^0$
> 4.9	95	9 ABDALLAH	04J DLPH	$e^+e^- \rightarrow Z^0$
> 8.5	95	10 ABDALLAH	04J DLPH	$e^+e^- \rightarrow Z^0$
> 5.0	95	11 ABDALLAH	03B DLPH	$e^+e^- \rightarrow Z$
> 10.3	95	12 ABE	03 SLD	$e^+e^- \rightarrow Z$
> 10.9	95	13 HEISTER	03E ALEP	$e^+e^- \rightarrow Z$
> 5.3	95	14 ABE	02V SLD	$e^+e^- \rightarrow Z$
> 1.0	95	15 ABBIENDI	01D OPAL	$e^+e^- \rightarrow Z$
> 7.4	95	16 ABREU	00Y DLPH	Repl. by ABDALLAH 04J
> 4.0	95	17 ABREU,P	00G DLPH	$e^+e^- \rightarrow Z$
> 5.2	95	18 ABBIENDI	99S OPAL	$e^+e^- \rightarrow Z$
< 96	95	19 ABE	99D CDF	$p\bar{p}$ at 1.8 TeV
> 5.8	95	20 ABE	99J CDF	$p\bar{p}$ at 1.8 TeV
> 9.6	95	21 BARATE	99J ALEP	$e^+e^- \rightarrow Z$
> 7.9	95	22 BARATE	98C ALEP	Repl. by BARATE 99J
> 3.1	95	23 ACKERSTAFF	97U OPAL	Repl. by ABBIENDI 99S
> 2.2	95	24 ACKERSTAFF	97V OPAL	Repl. by ABBIENDI 99S
> 6.5	95	25 ADAM	97 DLPH	Repl. by ABREU 00Y
> 6.6	95	26 BUSKULIC	96M ALEP	Repl. by BARATE 98C
> 2.2	95	24 AKERS	95J OPAL	Sup. by ACKERSTAFF 97V
> 5.7	95	27 BUSKULIC	95J ALEP	$e^+e^- \rightarrow Z$
> 1.8	95	24 BUSKULIC	94B ALEP	$e^+e^- \rightarrow Z$

¹ Measured using time-dependent angular analysis of $B_s^0 \rightarrow J/\psi K^+ K^-$ decays.

² Measured using $B_s^0 \rightarrow D_s^- \pi^+$ decays.

³ Measured using $B_s^0 \rightarrow D_s^- \mu^+ \nu_\mu X$ decays.

⁴ Measured using $B_s^0 \rightarrow D_s^- \pi^+$ and $D_s^- \pi^+ \pi^- \pi^+$ decays.

⁵ Significance of oscillation signal is 5.4σ . Also reports $|V_{td} / V_{ts}| = 0.2060 \pm 0.0007^{+0.0081}_{-0.0060}$.

- ⁶ A likelihood scan over the oscillation frequency, Δm_s , gives a most probable value of 19 ps^{-1} and a range of $17 < \Delta m_s < 21 \text{ (ps}^{-1})$ at 90% C.L. assuming Gaussian uncertainties. Also excludes $\Delta m_s < 14.8 \text{ ps}^{-1}$ at 95% C.L.
- ⁷ Significance of oscillation signal is 0.2%. Also reported the value $|V_{td} / V_{ts}| = 0.208^{+0.001}_{-0.002} {}^{+0.008}_{-0.006}$.
- ⁸ Uses leptons emitted with large momentum transverse to a jet and improved techniques for vertexing and flavor-tagging.
- ⁹ Updates of D_s -lepton analysis.
- ¹⁰ Combined results from all Delphi analyses.
- ¹¹ Events with a high transverse momentum lepton were removed and an inclusively reconstructed vertex was required.
- ¹² ABE 03 uses the novel “charge dipole” technique to reconstruct separate secondary and tertiary vertices originating from the $B \rightarrow D$ decay chain. The analysis excludes $\Delta m_s < 4.9 \text{ ps}^{-1}$ and $7.9 < \Delta m_s < 10.3 \text{ ps}^{-1}$.
- ¹³ Three analyses based on complementary event selections: (1) fully-reconstructed hadronic decays; (2) semileptonic decays with D_s exclusively reconstructed; (3) inclusive semileptonic decays.
- ¹⁴ ABE 02V uses exclusively reconstructed D_s^- mesons and excludes $\Delta m_s < 1.4 \text{ ps}^{-1}$ and $2.4 < \Delta m_s < 5.3 \text{ ps}^{-1}$ at 95%CL.
- ¹⁵ Uses fully or partially reconstructed $D_s \ell$ vertices and a mixing tag as a flavor tagging.
- ¹⁶ Replaced by ABDALLAH 04A. Uses $D_s^- \ell^+$, and $\phi \ell^+$ vertices, and a multi-variable discriminant as a flavor tagging.
- ¹⁷ Uses inclusive D_s vertices and fully reconstructed B_s decays and a multi-variable discriminant as a flavor tagging.
- ¹⁸ Uses ℓ - Q_{hem} and ℓ - ℓ .
- ¹⁹ ABE 99D assumes $\tau_{B_s^0} = 1.55 \pm 0.05 \text{ ps}$ and $\Delta\Gamma/\Delta m = (5.6 \pm 2.6) \times 10^{-3}$.
- ²⁰ ABE 99J uses ϕ ℓ - ℓ correlation.
- ²¹ BARATE 99J uses combination of an inclusive lepton and D_s^- -based analyses.
- ²² BARATE 98C combines results from $D_s h$ - ℓ / Q_{hem} , $D_s h$ - K in the same side, $D_s \ell$ - ℓ / Q_{hem} and $D_s \ell$ - K in the same side.
- ²³ Uses ℓ - Q_{hem} .
- ²⁴ Uses ℓ - ℓ .
- ²⁵ ADAM 97 combines results from $D_s \ell$ - Q_{hem} , ℓ - Q_{hem} , and ℓ - ℓ .
- ²⁶ BUSKULIC 96M uses D_s lepton correlations and lepton, kaon, and jet charge tags.
- ²⁷ BUSKULIC 95J uses ℓ - Q_{hem} . They find $\Delta m_s > 5.6$ [> 6.1] for $f_s = 10\%$ [12%]. We interpolate to our central value $f_s = 10.5\%$.

$$x_s = \Delta m_{B_s^0} / \Gamma_{B_s^0}$$

This is derived by the Heavy Flavor Averaging Group (HFLAV) from the results on $\Delta m_{B_s^0}$ and “OUR EVALUATION” of the B_s^0 mean lifetime.

<u>VALUE</u>	<u>DOCUMENT ID</u>
26.81±0.08 OUR EVALUATION	

$$\chi_s$$

This is a B_s^0 - \bar{B}_s^0 integrated mixing parameter derived from x_s above and OUR EVALUATION of $\Delta\Gamma_{B_s^0}/\Gamma_{B_s^0}$.

<u>VALUE</u>	<u>DOCUMENT ID</u>
0.499308±0.000004 OUR EVALUATION	

CP VIOLATION PARAMETERS in B_s^0

$$\text{Re}(\epsilon_{B_s^0}) / (1 + |\epsilon_{B_s^0}|^2)$$

CP impurity in B_s^0 system.

“OUR EVALUATION” is an average obtained by the Heavy Flavor Averaging Group (HFLAV) and described at <http://www.slac.stanford.edu/xorg/hflav/>. It is the result of a fit to B_d and B_s *CP* asymmetries, which includes the B_s measurements listed below and the B_d measurements listed in the B_d section, and takes into account correlations between those measurements.

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

-0.15±0.70 OUR EVALUATION

0.0 ±1.1 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.

$0.98 \pm 0.65 \pm 0.5$	¹ AAIJ	16G	LHCb	$p\bar{p}$ at 7, 8 TeV
-2.15 ± 1.85	² ABAZOV	14	D0	$p\bar{p}$ at 1.96 TeV
$-2.8 \pm 1.9 \pm 0.4$	³ ABAZOV	13	D0	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$-0.15 \pm 1.25 \pm 0.90$	⁴ AAIJ	14D	LHCb	Repl. by AAIJ 16G
-4.5 ± 2.7	⁵ ABAZOV	11U	D0	Repl. by ABAZOV 14
$-0.4 \pm 2.3 \pm 0.4$	⁶ ABAZOV	10E	D0	Repl. by ABAZOV 13
-3.6 ± 1.9	⁷ ABAZOV	10H	D0	Repl. by ABAZOV 11U
$6.1 \pm 4.8 \pm 0.9$	⁸ ABAZOV	07A	D0	Repl. by ABAZOV 10E

¹ AAIJ 16G reports a measurement of time-integrated flavor-specific asymmetry in $B_s^0 \rightarrow \mu^+ D_s^- X$ decays, $A_{SL}^s = (0.39 \pm 0.26 \pm 0.20)\%$, which is approximately equal to $4 \times \text{Re}(\epsilon_{B_s^0}) / (1 + |\epsilon_{B_s^0}|^2)$.

² ABAZOV 14 uses the dimuon charge asymmetry with different impact parameters from which it reports $A_{SL}^s = (-0.86 \pm 0.74) \times 10^{-2}$.

³ ABAZOV 13 reports a measurement of time-integrated flavor-specific asymmetry in mixed semileptonic $B_s^0 \rightarrow \mu^+ D_s^- X$ decays $A_{SL}^s = (-1.12 \pm 0.74 \pm 0.17)\%$ which is approximately equal to $4 \times \text{Re}(\epsilon_{B_s^0}) / (1 + |\epsilon_{B_s^0}|^2)$.


⁴ AAIJ 14D reports a measurement of time-integrated flavor-specific asymmetry in $B_s^0 \rightarrow \mu^+ D_s^- X$ decays, $A_{SL}^s = (-0.06 \pm 0.50 \pm 0.36)\%$, which is approximately equal to $4 \times \text{Re}(\epsilon_{B_s^0}) / (1 + |\epsilon_{B_s^0}|^2)$.

⁵ ABAZOV 11U uses the dimuon charge asymmetry with different impact parameters from which it reports $A_{SL}^s = (-18.1 \pm 10.6) \times 10^{-3}$.

⁶ ABAZOV 10E reports a measurement of flavor-specific asymmetry in $B_{(s)}^0 \rightarrow \mu^+ D_{(s)}^{*-} X$ decays with a decay-time analysis including initial-state flavor tagging, $A_{SL}^s = (-1.7 \pm 9.1^{+1.4}_{-1.5}) \times 10^{-3}$ which is approximately equal to $4 \times \text{Re}(\epsilon_{B_s^0}) / (1 + |\epsilon_{B_s^0}|^2)$.

⁷ ABAZOV 10H reports a measurement of like-sign dimuon charge asymmetry of $A_{SL}^s = (-9.57 \pm 2.51 \pm 1.46) \times 10^{-3}$ in semileptonic b -hadron decays. Using the measured production ratio of B_d^0 and B_s^0 , and the asymmetry of $B_d^0 A_{SL}^s = (-4.7 \pm 4.6) \times 10^{-3}$ measured from B -factories, they obtain the asymmetry for B_s^0 .

⁸ The first direct measurement of the time integrated flavor untagged charge asymmetry in semileptonic B_s^0 decays is reported as $2 \times A_{SL}^s$ (untagged) = $A_{SL}^s = (2.45 \pm 1.93 \pm 0.35) \times 10^{-2}$.



$C_{KK}(B_s^0 \rightarrow K^+ K^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.14±0.11±0.03	AAIJ	13BO LHCb	$p p$ at 7 TeV

$S_{KK}(B_s^0 \rightarrow K^+ K^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.30±0.12±0.04	AAIJ	13BO LHCb	$p p$ at 7 TeV

$r_B(B_s^0 \rightarrow D_s^\mp K^\pm)$

r_B and δ_B are the amplitude ratio and relative strong phase between the amplitudes of $A(B_s^0 \rightarrow D_s^+ K^-)$ and $A(B_s^0 \rightarrow D_s^- K^+)$,

VALUE	DOCUMENT ID	TECN	COMMENT
0.37±0.10 -0.09	¹ AAIJ	18U LHCb	$p p$ at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.53^{+0.17}_{-0.16}$	² AAIJ	14BF LHCb	Repl. by AAIJ 18U
------------------------	-------------------	-----------	-------------------

¹ Measured in $B_s^0 \rightarrow D_s^\mp K^\pm$ decays, constraining $-2\beta_s$ by the measurement of $\phi_s = -0.030 \pm 0.033$ from HFLAV.

² Measured in $B_s^0 \rightarrow D_s^\mp K^\pm$ decays, constraining $-2\beta_s$ by the measurement of $\phi_s = 0.01 \pm 0.07 \pm 0.0$ from AAIJ 13AR. At 68% CL.

$\delta_B(B_s^0 \rightarrow D_s^\pm K^\mp)$

VALUE (°)	DOCUMENT ID	TECN	COMMENT
358^{+13}_{-14}	¹ AAIJ	18U LHCb	$p\bar{p}$ at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

3^{+19}_{-20}	² AAIJ	14BF LHCb	Repl. by AAIJ 18U
-----------------	-------------------	-----------	-------------------

¹ Measured in $B_s^0 \rightarrow D_s^\mp K^\pm$ decays, constraining $-2\beta_s$ by the measurement of $\phi_s = 0.030 \pm 0.033$ from HFLAV. The value is modulo 180° .

² Measured in $B_s^0 \rightarrow D_s^\mp K^\pm$ decays, constraining $-2\beta_s$ by the measurement of $\phi_s = 0.01 \pm 0.07 \pm 0.0$ from AAIJ 13AR. The value is modulo 180° at 68% CL.

 CP Violation phase β_s

$-2\beta_s$ is the weak phase difference between B_s^0 mixing amplitude and the $B_s^0 \rightarrow J/\psi\phi$ decay amplitude driven by the $b \rightarrow c\bar{c}s$ transition (such as $B_s \rightarrow J/\psi\phi$, $J/\psi K^+ K^-$, $J/\psi\pi^+\pi^-$, and $D_s^+ D_s^-$). The Standard Model value of β_s is $\arg(-\frac{V_{ts}V_{tb}^*}{V_{cs}V_{cb}^*})$ if penguin contributions are neglected.

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFLAV) and are described at <http://www.slac.stanford.edu/xorg/hflav/>. The averaging/scaling procedure takes into account correlation between the measurements.

VALUE (10^{-2} rad)	DOCUMENT ID	TECN	COMMENT
1.1 ± 1.6 OUR EVALUATION			
1.8 ± 1.7 OUR AVERAGE			
$11.9 \pm 10.7 \pm 3.4$	¹ AAIJ	17V LHCb	$p\bar{p}$ at 7, 8 TeV
$4.5 \pm 3.9 \pm 2.1$	² AAD	16AP ATLAS	$p\bar{p}$ at 7, 8 TeV
$-11.5^{+14}_{-14.5} \pm 1$	³ AAIJ	16AK LHCb	$p\bar{p}$ at 7, 8 TeV
$3.75 \pm 4.85 \pm 1.55$	⁴ KHACHATRYAN...16S CMS	CMS	$p\bar{p}$ at 8 TeV
$2.9 \pm 2.5 \pm 0.3$	⁵ AAIJ	15I LHCb	$p\bar{p}$ at 7, 8 TeV
$-1 \pm 9 \pm 1$	⁶ AAIJ	14AY LHCb	$p\bar{p}$ at 7, 8 TeV
$-3.5 \pm 3.4 \pm 0.4$	⁷ AAIJ	14S LHCb	$p\bar{p}$ at 7, 8 TeV
	⁸ AALTONEN	12AJ CDF	$p\bar{p}$ at 1.96 TeV
28^{+18}_{-19}	⁹ ABAZOV	12D D0	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$5.0 \pm 6.5 \pm 7.0$	¹⁰ AAIJ	18S LHCb	$p\bar{p}$ at 7, 8 TeV
6^{+8}_{-7}	^{11,12} AAIJ	15K LHCb	$p\bar{p}$ at 7, 8 TeV
$-6 \pm 13 \pm 3$	¹³ AAD	14U ATLAS	Repl. by AAD 16AP
$-17 \pm 15 \pm 3$	¹⁴ AAIJ	14AE LHCb	$p\bar{p}$ at 7, 8 TeV
$-0.5 \pm 3.5 \pm 0.5$	¹⁵ AAIJ	13AR LHCb	Repl. by AAIJ 15I
	¹⁶ AAIJ	13AY LHCb	$p\bar{p}$ at 7 TeV
$-11.0 \pm 20.5 \pm 5.0$	¹⁷ AAD	12CV ATLAS	Repl. by AAD 14U
$22 \pm 22 \pm 1$	¹⁸ AAIJ	12B LHCb	Repl. by AAIJ 12Q
$-8 \pm 9 \pm 3$	¹⁹ AAIJ	12D LHCb	Repl. by AAIJ 13AR
$0.95^{+8.70+0.15}_{-8.65-0.20}$	²⁰ AAIJ	12Q LHCb	Repl. by AAIJ 13AR
	²¹ AALTONEN	12D CDF	Repl. by AALTONEN 12AJ

		22 AALTONEN	08G CDF	Repl. by AALTONEN 12D
28	$+12$ -15	$+4$ -1	9,23 ABAZOV	08AM D0
39.5	± 28.0	$+0.5$ -7.0	24,25 ABAZOV	07 D0
35	$+20$ -24	25,26 ABAZOV	07N D0	Repl. by ABAZOV 08AM

¹ Measured using time-dependent angular analysis of $B_s^0 \rightarrow J/\psi K^+ K^-$ in the region $m(KK) > 1.05$ GeV.

² AAD 16AP reports $\phi_s = -2 \beta_s = -0.090 \pm 0.078 \pm 0.041$ rad. that was measured using a time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays.

³ AAIJ 16AK reports $\phi_s = -2 \beta_s = 0.23^{+0.29}_{-0.28} \pm 0.02$ rad. that was measured using a time-dependent angular analysis of $B_s^0 \rightarrow \psi(2S) \phi$ decays.

⁴ KHACHATRYAN 16S reports $\phi_s = -2 \beta_s = -0.075 \pm 0.097 \pm 0.031$ rad. that was measured using a time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays.

⁵ AAIJ 15I reports $\phi_s = -2 \beta_s = -0.058 \pm 0.049 \pm 0.006$ rad. that was measured using a time-dependent angular analysis of $B_s^0 \rightarrow J/\psi K^+ K^-$ decays. It also combines this result with that of AAIJ 14S and quotes $\phi_s = -2 \beta_s = -0.010 \pm 0.039$ rad.

⁶ AAIJ 14AY reports $\phi_s = -2 \beta_s = 0.02 \pm 0.17 \pm 0.02$ rad. in a time-dependent fit to $B_s^0 \rightarrow D_s^+ D_s^-$, while allowing CP violation in decay.

⁷ AAIJ 14S reports $\phi_s = -2 \beta_s = 0.070 \pm 0.068 \pm 0.008$ rad. and $|\lambda| = 0.89 \pm 0.05 \pm 0.01$, when direct CP violation is allowed. Measured using a time-dependent fit to $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$ decays.

⁸ AALTONEN 12AJ reports $-\pi/2 < \beta_s < -1.51$ or $-0.06 < \beta_s < 0.30$, or $1.26 < \beta_s < \pi/2$ rad. at 68% CL. Measured using the time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays.

⁹ ABAZOV 12D reports $\phi_s = -2 \beta_s = -0.55^{+0.38}_{-0.36}$ rad. that was measured using a time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays. A single error includes both statistical and systematic uncertainties.

¹⁰ AAIJ 18S reports $\phi_s = -2 \beta_s = -0.10 \pm 0.13 \pm 0.14$ rad measured in $B_s^0 \rightarrow (K^+ \pi^-)(K^- \pi^+)$ in the region $0.75 < m(K^\pm \pi^\mp) < 1.6$ GeV. This is a $b \rightarrow d \bar{d}s$ transition with a decay amplitude phase different from that of $b \rightarrow c \bar{c}s$ transition.

¹¹ AAIJ 15K reports $-2\beta_s = -0.12^{+0.14}_{-0.16}$ rad. The value was obtained by measuring time-dependent CP asymmetry in $B_s^0 \rightarrow K^+ K^-$ and using a U-spin relation between $B_s^0 \rightarrow K^+ K^-$ and $B^0 \rightarrow \pi^+ \pi^-$.

¹² Results are also presented using additional inputs on $B^0 \rightarrow \pi^0 \pi^0$ and $B^+ \rightarrow \pi^+ \pi^0$ decays from other experiments and isospin symmetry assumptions. The dependence of the results on the maximum allowed amount of U-spin breaking up to 50% is also included.

¹³ AAD 14U reports $\phi_s = -2 \beta_s = 0.12 \pm 0.25 \pm 0.05$ rad. that was measured using a time-dependent angular analysis of $B_s^0 \rightarrow J/\psi \phi$ decays.

¹⁴ Measured in $B_s^0 \rightarrow \phi \phi$ decays. This is a $b \rightarrow s \bar{s}s$ transition with a decay amplitude phase different from that of $b \rightarrow c \bar{c}s$ transition.

¹⁵ AAIJ 13AR reports $\phi_s = -2\beta_s = 0.01 \pm 0.07 \pm 0.01$ rad. obtained from combined fit to $B_s^0 \rightarrow J/\psi K^+ K^-$ and $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$ data sets. Also reports separate results of $\phi_s = 0.07 \pm 0.09 \pm 0.01$ rad. from $B_s^0 \rightarrow J/\psi K^+ K^-$ decays and $\phi_s = -0.14^{+0.17}_{-0.16} \pm 0.01$ rad. from $B_s^0 \rightarrow J/\psi \pi^+ \pi^-$ decays.

- 16 AAIJ 13AY uses $B_s^0 \rightarrow \phi\phi$ mode, and reports the 68% CL interval of $\phi_s = -2\beta_s$ as $[-2.46, -0.76]$ rad.
- 17 AAD 12CV reports $\phi_s = -2\beta_s = 0.22 \pm 0.41 \pm 0.10$ rad. that was measured using a time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays.
- 18 Reports $\phi_s = -2\beta_s = -0.44 \pm 0.44 \pm 0.02$ rad. that was measured using a time-dependent fit to $B_s^0 \rightarrow J/\psi f_0(980)$ decays.
- 19 Reports $\phi_s = -2\beta_s = 0.15 \pm 0.18 \pm 0.06$ rad. that was measured using a time-dependent angular analysis of $B_s^0 \rightarrow J/\psi\phi$ decays.
- 20 Reports $\phi_s = -2\beta_s = -0.019^{+0.173+0.004}_{-0.174-0.003}$ rad. which was measured using a time-dependent fit to $B_s^0 \rightarrow J/\psi\pi^+\pi^-$ decays, with the $\pi^+\pi^-$ mass within 775–1550 MeV. Searches for, but finds no evidence, for direct CP violation in $B_s^0 \rightarrow J/\psi\pi\pi$ decays.
- 21 Reports $0.02 < \phi_s < 0.52$ or $1.08 < \phi_s < 1.55$ rad. at 68% C.L. confidence regions in the two-dimensional space of ϕ_s and $\Delta\Gamma_{B_s^0}$ from $B_s^0 \rightarrow J/\psi\phi$ decays.
- 22 Reports $0.32 < 2\beta_s < 2.82$ rad. at 68% C.L. and confidence regions in the two-dimensional space of $2\beta_s$ and $\Delta\Gamma$ from the first measurement of $B_s^0 \rightarrow J/\psi\phi$ decays using flavor tagging. The probability of a deviation from SM prediction as large as the level of observed data is 15%.
- 23 Reports $\phi_s = -2\beta_s$ and obtains 90% CL interval $-0.03 < \beta_s < 0.60$ rad.
- 24 The first direct measurement of the CP -violating mixing phase is reported from the time-dependent analysis of flavor untagged $B_s^0 \rightarrow J/\psi\phi$ decays.
- 25 Reports ϕ_s which equals to $-2\beta_s$.
- 26 Combines D0 collaboration measurements of time-dependent angular distributions in $B_s^0 \rightarrow J/\psi\phi$ and charge asymmetry in semileptonic decays. There is a 4-fold ambiguity in the solution.

$|\lambda| (B_s^0 \rightarrow J/\psi(1S)\phi)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.964 \pm 0.019 \pm 0.007$	AAIJ	15I	LHCb $p p$ at 7, 8 TeV

$|\lambda|$

VALUE	DOCUMENT ID	TECN	COMMENT
1.002 ± 0.017 OUR AVERAGE			
$1.035 \pm 0.034 \pm 0.089$	1 AAIJ	18S	LHCb $p p$ at 7, 8 TeV
$0.994 \pm 0.018 \pm 0.006$	2 AAIJ	17V	LHCb $p p$ at 7, 8 TeV
$1.045^{+0.069}_{-0.050} \pm 0.007$	3 AAIJ	16AK	LHCb $p p$ at 7, 8 TeV
$1.04 \pm 0.07 \pm 0.03$	4 AAIJ	14AE	LHCb $p p$ at 7, 8 TeV
$0.91^{+0.18}_{-0.15} \pm 0.02$	5 AAIJ	14AY	LHCb $p p$ at 7, 8 TeV

¹ Measured in $B_s^0 \rightarrow (K^+\pi^-)(K^-\pi^+)$ in the region $0.75 < m(K^\pm\pi^\mp) < 1.6$ GeV.

² Measured using time-dependent angular analysis of $B_s^0 \rightarrow J/\psi K^+ K^-$ in the region $m(KK) > 1.05$ GeV.

³ Measured using time-dependent angular analysis of $B_s^0 \rightarrow \psi(2S)\phi$ decays.

⁴ Measured in $B_s^0 \rightarrow \phi\phi$ decays.

⁵ Measured in $B_s^0 \rightarrow D_s^+ D_s^-$ decays.

A, CP violation parameter

$$A = -2 \operatorname{Re}(\lambda) / (1 + |\lambda|^2)$$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.75±0.12 OUR AVERAGE			
-0.79±0.07±0.10	1 AAIJ	180 LHCb	$p p$ at 7, 8 TeV
0.49 ^{+0.77} _{-0.65} ±0.06	2 AAIJ	15AL LHCb	$p p$ at 7, 8 TeV

¹ Measured in $B_s^0 \rightarrow K^+ K^-$ decays.

² Measured in $B_s^0 \rightarrow J/\psi K_S^0$ decays.

C, CP violation parameter

$$C = (1 - |\lambda|^2) / (1 + |\lambda|^2)$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.19±0.06 OUR AVERAGE			
0.20±0.06±0.02	1 AAIJ	180 LHCb	$p p$ at 7, 8 TeV
-0.28±0.41±0.08	2 AAIJ	15AL LHCb	$p p$ at 7, 8 TeV

¹ Measured in $B_s^0 \rightarrow K^+ K^-$ decays.

² Measured in $B_s^0 \rightarrow J/\psi K_S^0$ decays.

S, CP violation parameter

$$S = -2 \operatorname{Im}(\lambda) / (1 + |\lambda|^2)$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.17±0.06 OUR AVERAGE			
0.18±0.06±0.02	1 AAIJ	180 LHCb	$p p$ at 7, 8 TeV
-0.08±0.40±0.08	2 AAIJ	15AL LHCb	$p p$ at 7, 8 TeV

¹ Measured in $B_s^0 \rightarrow K^+ K^-$ decays.

² Measured in $B_s^0 \rightarrow J/\psi K_S^0$ decays.

 $A_{CP}^L(B_s \rightarrow J/\psi \bar{K}^*(892)^0)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.048±0.057±0.020			

 $A_{CP}^{\parallel}(B_s \rightarrow J/\psi \bar{K}^*(892)^0)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.171±0.152±0.028			

 $A_{CP}^{\perp}(B_s \rightarrow J/\psi \bar{K}^*(892)^0)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.049±0.096±0.025			

 $A_{CP}(B_s \rightarrow \pi^+ K^-)$

A_{CP} is defined as

$$\frac{B(\bar{B}_s^0 \rightarrow f) - B(B_s^0 \rightarrow \bar{f})}{B(\bar{B}_s^0 \rightarrow f) + B(B_s^0 \rightarrow \bar{f})},$$

the CP -violation asymmetry of exclusive B_s^0 and \bar{B}_s^0 decay.

VALUE	DOCUMENT ID	TECN	COMMENT
0.221±0.015 OUR AVERAGE			
0.213±0.015±0.007	AAIJ	180	LHCb $p\bar{p}$ at 7, 8 TeV
0.22 ±0.07 ±0.02	AALTONEN	14P	CDF $p\bar{p}$ at 1.96 TeV
0.27 ±0.04 ±0.01	AAIJ	13AX	LHCb $p\bar{p}$ at 7 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.27 ±0.08 ±0.02	AAIJ	12V	LHCb Repl. by AAIJ 13AX
0.39 ±0.15 ±0.08	AALTONEN	11N	CDF Repl. by AALTONEN 14P

$A_{CP}(B_s^0 \rightarrow [K^+ K^-]_D \bar{K}^*(892)^0)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.04±0.07±0.02	AAIJ	14BN	LHCb $p\bar{p}$ at 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.04±0.16±0.01	AAIJ	13L	LHCb Repl. by AAIJ 14BN

$A_{CP}(B_s^0 \rightarrow [\pi^+ \pi^-]_D K^*(892)^0)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.01±0.03±0.02	AAIJ	14BN	LHCb $p\bar{p}$ at 7, 8 TeV

$A_{CP}(B_s^0 \rightarrow [\pi^+ \pi^-]_D K^*(892)^0)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.06±0.13±0.02	AAIJ	14BN	LHCb $p\bar{p}$ at 7, 8 TeV

$A^\Delta(B_s \rightarrow \phi\gamma)$

$A^\Delta(B_s \rightarrow \phi\gamma)$ is the multiplicative coefficient of the $\sinh(\Delta\Gamma t/2)$ term in the $B_s \rightarrow \phi\gamma$ decay rate time dependence.

VALUE	DOCUMENT ID	TECN	COMMENT
-0.98±0.46±0.23 -0.52±0.20	1 AAIJ	17B	LHCb $p\bar{p}$ at 7, 8 TeV

¹ Measured in time dependent analysis without initial flavor tagging.

CPT VIOLATION PARAMETERS

In the B_s^0 mixing, propagating mass eigenstates can be written as

$$\begin{aligned} |B_{sL}\rangle &\propto p \sqrt{1-\xi} |B_s^0\rangle + q \sqrt{1+\xi} |\bar{B}_s^0\rangle \\ |B_{sH}\rangle &\propto p \sqrt{1+\xi} |B_s^0\rangle - q \sqrt{1-\xi} |\bar{B}_s^0\rangle \end{aligned}$$

where parameter ξ controls CPT violation. If ξ is zero, then CPT is conserved. The parameter ξ can be written as

$$\xi = \frac{2(M_{11}-M_{22})-i(\Gamma_{11}-\Gamma_{22})}{-2\Delta m_s+i\Delta\Gamma_s} \approx \frac{-2\beta^\mu \Delta a_\mu}{2\Delta m_s-i\Delta\Gamma_s},$$

where M_{ii} , Γ_{ii} , Δm_s , and $\Delta\Gamma_s$ are parameters of Hamiltonian governing B_s oscillations, β^μ is the B_s^0 meson velocity and Δa_μ characterizes Lorentz-invariance violation.

Δa_\perp

VALUE (10^{-12} GeV)	CL%	DOCUMENT ID	TECN	COMMENT
-0.47±0.39±0.08		1 AAIJ	16E	LHCb $p\bar{p}$ at 7, 8 TeV
< 1.2	95	2 ABAZOV	15L	D0 $p\bar{p}$ at 1.96 TeV

¹ Uses $B_s^0 \rightarrow J/\psi K^+ K^-$ decays.

² Measured in semileptonic $B_s^0 \rightarrow D_s^- \mu^+ X$ decays. Also extracts limit on time and longitudinal components ($-0.8 < \Delta a_T - 0.396 \Delta a_Z < 3.9$) 10^{-13} GeV.

$\Delta a_{||}$

VALUE (10^{-14} GeV)	DOCUMENT ID	TECN	COMMENT
-0.89 ± 1.41 ± 0.36	¹ AAIJ	16E LHCb	$p\bar{p}$ at 7, 8 TeV

¹ Uses $B_s^0 \rightarrow J/\psi K^+ K^-$ decays.

Δa_X

VALUE (10^{-14} GeV)	DOCUMENT ID	TECN	COMMENT
+1.01 ± 2.08 ± 0.71	¹ AAIJ	16E LHCb	$p\bar{p}$ at 7, 8 TeV

¹ Uses $B_s^0 \rightarrow J/\psi K^+ K^-$ decays.

Δa_Y

VALUE (10^{-14} GeV)	DOCUMENT ID	TECN	COMMENT
-3.83 ± 2.09 ± 0.71	¹ AAIJ	16E LHCb	$p\bar{p}$ at 7, 8 TeV

¹ Uses $B_s^0 \rightarrow J/\psi K^+ K^-$ decays.

$\text{Re}(\xi)$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.022 ± 0.033 ± 0.003	¹ AAIJ	16E LHCb	$p\bar{p}$ at 7, 8 TeV

¹ Uses $B_s^0 \rightarrow J/\psi K^+ K^-$ decays.

$\text{Im}(\xi)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.004 ± 0.011 ± 0.002	¹ AAIJ	16E LHCb	$p\bar{p}$ at 7, 8 TeV

¹ Uses $B_s^0 \rightarrow J/\psi K^+ K^-$ decays.

PARTIAL BRANCHING FRACTIONS IN $B_s \rightarrow \phi \ell^+ \ell^-$

$\mathbf{B}(B_s \rightarrow \phi \ell^+ \ell^-) (0.1 < q^2 < 2.0 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
1.14 ± 0.16 OUR AVERAGE			
1.11 ± 0.14 ± 0.09	¹ AAIJ	15AQ LHCb	$p\bar{p}$ at 7, 8 TeV
2.78 ± 0.95 ± 0.89	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.897 ± 0.207 ± 0.097	¹ AAIJ	13X LHCb	Repl. by AAIJ 15AQ
¹ Measured in $B_s^0 \rightarrow \phi \mu^+ \mu^-$ decays.			

$\mathbf{B}(B_s \rightarrow \phi \ell^+ \ell^-) (2.0 < q^2 < 5.0 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
0.77 ± 0.12 ± 0.06	¹ AAIJ	15AQ LHCb	$p\bar{p}$ at 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.529 ± 0.182 ± 0.057	^{1,2} AAIJ	13X LHCb	Repl. by AAIJ 15AQ
0.58 ± 0.55 ± 0.19	² AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV

¹ Measured in $B_s^0 \rightarrow \phi \mu^+ \mu^-$ decays.

² Measured in $2 < q^2 < 4.3 \text{ GeV}^2/c^4$.

$B(B_s \rightarrow \phi \ell^+ \ell^-) (5.0 < q^2 < 8.0 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
0.96±0.13±0.08	¹ AAIJ	15AQ LHCb	$p\bar{p}$ at 7, 8 TeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$1.38^{+0.25}_{-0.23} \pm 0.14$	^{1,2} AAIJ	13X LHCb	Repl. by AAIJ 15AQ
$1.34 \pm 0.83 \pm 0.43$	² AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
1 Measured in $B_s^0 \rightarrow \phi \mu^+ \mu^-$ decays. 2 Measured in $4.3 < q^2 < 8.68 \text{ GeV}^2/c^4$.			

$B(B_s \rightarrow \phi \ell^+ \ell^-) (11.0 < q^2 < 12.5 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
0.71±0.10±0.06	¹ AAIJ	15AQ LHCb	$p\bar{p}$ at 7, 8 TeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$1.18^{+0.22}_{-0.21} \pm 0.14$	^{1,2} AAIJ	13X LHCb	Repl. by AAIJ 15AQ
$2.98 \pm 0.95 \pm 0.95$	² AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
1 Measured in $B_s^0 \rightarrow \phi \mu^+ \mu^-$ decays. 2 Measured in $10.9 < q^2 < 12.86 \text{ GeV}^2/c^4$.			

$B(B_s \rightarrow \phi \ell^+ \ell^-) (15.0 < q^2 < 17.0 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
0.90 ± 0.11 ± 0.07	¹ AAIJ	15AQ LHCb	$p\bar{p}$ at 7, 8 TeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$0.760^{+0.189}_{-0.169} \pm 0.087$	^{1,2} AAIJ	13X LHCb	Repl. by AAIJ 15AQ
$1.86 \pm 0.66 \pm 0.59$	² AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
1 Measured in $B_s^0 \rightarrow \phi \mu^+ \mu^-$ decays. 2 Measured in $14.18 < q^2 < 16 \text{ GeV}^2/c^4$.			

$B(B_s \rightarrow \phi \ell^+ \ell^-) (17.0 < q^2 < 19.0 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
0.79±0.11±0.07	¹ AAIJ	15AQ LHCb	$p\bar{p}$ at 7, 8 TeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$1.06^{+0.23}_{-0.21} \pm 0.12$	^{1,2} AAIJ	13X LHCb	Repl. by AAIJ 15AQ
$2.32 \pm 0.76 \pm 0.74$	² AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
1 Measured in $B_s^0 \rightarrow \phi \mu^+ \mu^-$ decays. 2 Measured in $16 < q^2 < 19 \text{ GeV}^2/c^4$.			

$B(B_s \rightarrow \phi \ell^+ \ell^-) (1.0 < q^2 < 6.0 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
1.28±0.18 OUR AVERAGE			
$1.29 \pm 0.16 \pm 0.10$	¹ AAIJ	15AQ LHCb	$p\bar{p}$ at 7, 8 TeV
$1.14 \pm 0.79 \pm 0.36$	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$1.14^{+0.25}_{-0.23} \pm 0.13$	¹ AAIJ	13X LHCb	Repl. by AAIJ 15AQ

¹ Measured in $B_s^0 \rightarrow \phi \mu^+ \mu^-$ decays.

$\mathbf{B}(B_s \rightarrow \phi \ell^+ \ell^-) (0.0 < q^2 < 4.3 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
$3.30 \pm 1.09 \pm 1.05$	AALTONEN	11AI	CDF $p\bar{p}$ at 1.96 TeV

PRODUCTION ASYMMETRIES

$A_P(B_s^0)$

$$A_P(B_s^0) = [\sigma(\overline{B}_s^0) - \sigma(B_s^0)] / [\sigma(\overline{B}_s^0) + \sigma(B_s^0)]$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
1.2 ± 1.6 OUR AVERAGE			
$-0.65 \pm 2.88 \pm 0.59$	¹ AAIJ	17BF LHCb	$p\bar{p}$ at 7 TeV
$1.98 \pm 1.90 \pm 0.59$	¹ AAIJ	17BF LHCb	$p\bar{p}$ at 8 TeV
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$1.09 \pm 2.61 \pm 0.66$	² AAIJ	14BP LHCb	Repl. by AAIJ 17BF, $p\bar{p}$ at 7 TeV

¹ Based on time-dependent analysis of $B_s^0 \rightarrow D_s^- \pi^+$ in kinematic range $2 < p_T < 30$ GeV/c and $2.1 < \eta < 4.5$.

² Based on time-dependent analysis of $B_s^0 \rightarrow D_s^- \pi^+$ in kinematic range $4 < p_T < 30$ GeV/c and $2.5 < \eta < 4.5$.

B_s^0 REFERENCES

AABOUD	19L	1812.03017	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AAIJ	18AB	JHEP 1807 020	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18AC	JHEP 1808 191	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18AY	PR D98 071103	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18AZ	PR D98 072006	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18O	PR D98 032004	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18S	JHEP 1803 140	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18T	JHEP 1803 078	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18U	JHEP 1803 059	R. Aaij <i>et al.</i>	(LHCb Collab.)
SIRUNYAN	18BY	EPJ C78 457	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AAIJ	17A	PR D95 012006	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17AI	PRL 118 191801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17AJ	PRL 118 251802	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17AL	PRL 119 041802	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17AN	PRL 119 101801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17B	PRL 118 021801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BA	JHEP 1705 158	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BD	PR D96 051103	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BF	PL B774 139	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BJ	PRL 119 232001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BP	JHEP 1711 027	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17G	PRL 118 081801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17N	JHEP 1703 001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17U	JHEP 1707 021	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17V	JHEP 1708 037	R. Aaij <i>et al.</i>	(LHCb Collab.)
AABOUD	16L	EPJ C76 513	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AAD	16AP	JHEP 1608 147	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAIJ	16	JHEP 1601 012	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	16AK	PL B762 253	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	16AL	PL B762 484	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	16C	PRL 116 161802	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	16E	PRL 116 241601	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	16G	PRL 117 061803	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	16P	PR D93 092008	R. Aaij <i>et al.</i>	(LHCb Collab.)

AAIJ	16U	JHEP 1603 040	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABAZOV	16C	PR D94 012001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
KHACHATRY...	16Q	PL B756 84	V. Khachatryan <i>et al.</i>	(CMS Collab.)
KHACHATRY...	16S	PL B757 97	V. Khachatryan <i>et al.</i>	(CMS Collab.)
PAL	16	PRL 116 161801	B. Pal <i>et al.</i>	(BELLE Collab.)
AAIJ	15AC	JHEP 1505 019	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15AD	JHEP 1506 130	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15AF	JHEP 1507 166	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15AG	JHEP 1508 005	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15AL	JHEP 1506 131	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15AQ	JHEP 1509 179	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15AS	JHEP 1510 053	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15AV	JHEP 1511 082	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15BB	PR D92 112002	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15D	JHEP 1501 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15I	PRL 114 041801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15K	PL B741 1	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15O	PRL 115 051801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15S	PL B743 46	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15U	PL B747 484	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABAZOV	15A	PRL 114 062001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	15L	PRL 115 161601	V.M. Abazov <i>et al.</i>	(D0 Collab.)
DUTTA	15	PR D91 011101	D. Dutta <i>et al.</i>	(BELLE Collab.)
KHACHATRY...	15BE	NAT 522 68	V. Khachatryan <i>et al.</i>	(CMS and LHCb Collab.)
OSWALD	15	PR D92 072013	C. Oswald <i>et al.</i>	(BELLE Collab.)
AAD	14U	PR D90 052007	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAIJ	14AA	PRL 112 202001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14AE	PR D90 052011	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14AX	PRL 113 172001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14AY	PRL 113 211801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14BF	JHEP 1411 060	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14BH	PR D90 072003	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14BM	NJP 16 123001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14BN	PR D90 112002	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14BP	PL B739 218	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14BR	PR D89 092006	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14D	PL B728 607	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14E	JHEP 1404 114	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14F	PRL 112 111802	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14L	JHEP 1407 140	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14R	PL B736 446	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14S	PL B736 186	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14Y	PRL 112 091802	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTENON	14P	PRL 113 242001	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	14	PR D89 012002	V.M. Abazov <i>et al.</i>	(D0 Collab.)
PDG	14	CP C38 070001	K. Olive <i>et al.</i>	(PDG Collab.)
AAIJ	13	NP B867 1	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13A	NP B867 547	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13AA	NP B871 403	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13AB	NP B873 275	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13AC	NP B874 663	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13AL	PR D87 071101	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13AN	PR D87 072004	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13AP	PR D87 092007	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13AQ	PR D87 112009	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13AR	PR D87 112010	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13AW	PRL 110 211801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13AX	PRL 110 221601	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13AY	PRL 110 241802	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13B	PRL 110 021801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13BA	PRL 111 101805	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13BI	NJP 15 053021	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13BM	PRL 111 141801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13BO	JHEP 1310 183	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13BP	JHEP 1310 143	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13BQ	JHEP 1310 005	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13BW	JHEP 1311 092	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13BX	PL B727 403	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13CF	EPJ C73 2655	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13L	JHEP 1303 067	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13X	JHEP 1307 084	R. Aaij <i>et al.</i>	(LHCb Collab.)

AAIJ	13Z	JHEP 1309 006	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTENON	13F	PR D87 072003	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	13	PRL 110 011801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	13C	PR D87 072006	V.M. Abazov <i>et al.</i>	(D0 Collab.)
CHATRCHYAN	13AW	PRL 111 101804	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
ESEN	13	PR D87 031101	S. Esen <i>et al.</i>	(BELLE Collab.)
OSWALD	13	PR D87 072008	C. Oswald <i>et al.</i>	(BELLE Collab.)
Also		PR D90 119901 (errat.)	C. Oswald <i>et al.</i>	(BELLE Collab.)
SOLOVIEVA	13	PL B726 206	E. Solovieva <i>et al.</i>	(BELLE Collab.)
THORNE	13	PR D88 114006	F. Thorne <i>et al.</i>	(BELLE Collab.)
AAD	12AE	PL B713 387	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAD	12CV	JHEP 1212 072	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAIJ	12	PL B707 349	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12A	PL B708 55	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12AE	PR D85 112013	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12AG	JHEP 1206 115	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12AM	PRL 109 131801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12AN	PRL 109 152002	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12AO	PR D86 052006	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12AP	PR D86 071102	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12AR	JHEP 1210 037	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12AX	PR D86 112005	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12B	PL B707 497	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12D	PRL 108 101803	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12E	PL B708 241	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12F	PL B709 50	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12I	PL B709 177	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12L	EPJ C72 2118	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12O	PL B713 172	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12P	PL B713 369	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12Q	PL B713 378	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12R	PL B716 393	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12S	PRL 108 151801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12V	PRL 108 201601	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12W	PRL 108 231801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTENON	12AJ	PRL 109 171802	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTENON	12C	PRL 108 201801	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTENON	12D	PR D85 072002	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTENON	12L	PRL 108 211803	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	12AF	PR D86 092011	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	12C	PR D85 011103	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	12D	PR D85 032006	V.M. Abazov <i>et al.</i>	(D0 Collab.)
CHATRCHYAN	12A	JHEP 1204 033	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
LEES	12A	PR D85 011101	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LI	12	PRL 108 181808	J. Li <i>et al.</i>	(BELLE Collab.)
PDG	12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)
AAIJ	11	PL B698 115	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	11A	PL B698 14	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	11B	PL B699 330	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	11D	PL B706 32	R. Aaij	(LHCb Collab.)
AAIJ	11E	PR D84 092001	R. Aaij <i>et al.</i>	(LHCb Collab.)
Also		PR D85 039904 (errat.)	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTENON	11A	PR D83 052012	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTENON	11AB	PR D84 052012	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTENON	11AG	PRL 107 191801	T. Aaltonen <i>et al.</i>	(CDF Collab.)
Also		PRL 107 239903 (errat.)	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTENON	11AI	PRL 107 201802	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTENON	11AN	PRL 107 261802	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTENON	11AP	PRL 107 272001	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTENON	11L	PRL 106 161801	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTENON	11N	PRL 106 181802	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	11U	PR D84 052007	V.M. Abazov <i>et al.</i>	(D0 Collab.)
CHATRCHYAN	11T	PRL 107 191802	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
LI	11	PRL 106 121802	J. Li <i>et al.</i>	(BELLE Collab.)
ABAZOV	10E	PR D82 012003	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	10H	PRL 105 081801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
Also		PR D82 032001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	10S	PL B693 539	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ESEN	10	PRL 105 201802	S. Esen <i>et al.</i>	(BELLE Collab.)
LOUVOT	10	PRL 104 231801	R. LOUVOT <i>et al.</i>	(BELLE Collab.)
PENG	10	PR D82 072007	C.-C. Peng <i>et al.</i>	(BELLE Collab.)

AALTONEN	09AQ	PRL 103 191802	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	09B	PR D79 011104	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	09C	PRL 103 031801	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	09P	PRL 102 201801	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	09E	PRL 102 032001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	09G	PRL 102 051801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	09I	PRL 102 091801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	09Y	PR D79 111102	V.M. Abazov <i>et al.</i>	(D0 Collab.)
LOUVOT	09	PRL 102 021801	R. Louvot <i>et al.</i>	(BELLE Collab.)
AALTONEN	08F	PRL 100 021803	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	08G	PRL 100 161802	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	08J	PRL 100 121803	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	08AM	PRL 101 241801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
WICHT	08A	PRL 100 121801	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABAZOV	07	PRL 98 121801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	07A	PRL 98 151801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	07N	PR D76 057101	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	07Y	PRL 99 241801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABULENCIA	07C	PRL 98 061802	A. Abulencia <i>et al.</i>	(CDF Collab.)
DRUTSKOY	07	PRL 98 052001	A. Drutskoy <i>et al.</i>	(BELLE Collab.)
DRUTSKOY	07A	PR D76 012002	A. Drutskoy <i>et al.</i>	(BELLE Collab.)
ABAZOV	06B	PRL 97 021802	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	06G	PR D74 031107	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	06V	PRL 97 241801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABULENCIA	06J	PRL 96 191801	A. Abulencia <i>et al.</i>	(CDF Collab.)
ABULENCIA	06N	PRL 96 231801	A. Abulencia <i>et al.</i>	(CDF Collab.)
ABULENCIA	06Q	PRL 97 062003	A. Abulencia <i>et al.</i>	(CDF Collab.)
ABULENCIA,A	06D	PRL 97 211802	A. Abulencia <i>et al.</i>	(CDF Collab.)
ABULENCIA,A	06G	PRL 97 242003	A. Abulencia <i>et al.</i>	(CDF Collab.)
ACOSTA	06	PRL 96 202001	D. Acosta <i>et al.</i>	(CDF Collab.)
ABAZOV	05B	PRL 94 042001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	05W	PRL 95 171801	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ACOSTA	05	PRL 94 101803	D. Acosta <i>et al.</i>	(CDF Collab.)
ACOSTA	05J	PRL 95 031801	D. Acosta <i>et al.</i>	(CDF Collab.)
ABDALLAH	04A	PL B585 63	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABDALLAH	04J	EPJ C35 35	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABDALLAH	03B	EPJ C28 155	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ABE	03	PR D67 012006	K. Abe <i>et al.</i>	(SLD Collab.)
HEISTER	03E	EPJ C29 143	A. Heister <i>et al.</i>	(ALEPH Collab.)
ABE	02V	PR D66 032009	K. Abe <i>et al.</i>	(SLD Collab.)
ACOSTA	02D	PR D65 111101	D. Acosta <i>et al.</i>	(CDF Collab.)
ACOSTA	02G	PR D66 112002	D. Acosta <i>et al.</i>	(CDF Collab.)
ABBIENDI	01D	EPJ C19 241	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABE	00C	PR D62 071101	K. Abe <i>et al.</i>	(SLD Collab.)
ABREU	00Y	EPJ C16 555	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU,P	00G	EPJ C18 229	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AFFOLDER	00N	PRL 85 4668	T. Affolder <i>et al.</i>	(CDF Collab.)
BARATE	00K	PL B486 286	R. Barate <i>et al.</i>	(ALEPH Collab.)
ABBIENDI	99S	EPJ C11 587	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABE	99D	PR D59 032004	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	99J	PRL 82 3576	F. Abe <i>et al.</i>	(CDF Collab.)
BARATE	99J	EPJ C7 553	R. Barate <i>et al.</i>	(ALEPH Collab.)
Also		EPJ C12 181 (errat.)	R. Barate <i>et al.</i>	(ALEPH Collab.)
ABE	98B	PR D57 5382	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	98V	PRL 81 5742	F. Abe <i>et al.</i>	(CDF Collab.)
ACCIARRI	98S	PL B438 417	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACKERSTAFF	98F	EPJ C2 407	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	98G	PL B426 161	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
BARATE	98C	EPJ C4 367	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARATE	98Q	EPJ C4 387	R. Barate <i>et al.</i>	(ALEPH Collab.)
PDG	98	EPJ C3 1	C. Caso <i>et al.</i>	(PDG Collab.)
ACCIARRI	97B	PL B391 474	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	97C	PL B391 481	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACKERSTAFF	97U	ZPHY C76 401	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ACKERSTAFF	97V	ZPHY C76 417	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ADAM	97	PL B414 382	W. Adam <i>et al.</i>	(DELPHI Collab.)
ABE	96B	PR D53 3496	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96N	PRL 77 1945	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96Q	PR D54 6596	F. Abe <i>et al.</i>	(CDF Collab.)
ABREU	96F	ZPHY C71 11	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ADAM	96D	ZPHY C72 207	W. Adam <i>et al.</i>	(DELPHI Collab.)

BUSKULIC	96E	ZPHY C69 585	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	96M	PL B377 205	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	96V	PL B384 471	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
PDG	96	PR D54 1	R. M. Barnett <i>et al.</i>	(PDG Collab.)
ABE	95R	PRL 74 4988	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	95Z	PRL 75 3068	F. Abe <i>et al.</i>	(CDF Collab.)
ACCIARRI	95H	PL B363 127	M. Acciarri <i>et al.</i>	(L3 Collab.)
ACCIARRI	95I	PL B363 137	M. Acciarri <i>et al.</i>	(L3 Collab.)
AKERS	95G	PL B350 273	R. Akers <i>et al.</i>	(OPAL Collab.)
AKERS	95J	ZPHY C66 555	R. Akers <i>et al.</i>	(OPAL Collab.)
BUSKULIC	95J	PL B356 409	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	95O	PL B361 221	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABREU	94D	PL B324 500	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	94E	ZPHY C61 407	P. Abreu <i>et al.</i>	(DELPHI Collab.)
Also		PL B289 199	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AKERS	94J	PL B337 196	R. Akers <i>et al.</i>	(OPAL Collab.)
AKERS	94L	PL B337 393	R. Akers <i>et al.</i>	(OPAL Collab.)
BUSKULIC	94B	PL B322 441	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	94C	PL B322 275	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABE	93F	PRL 71 1685	F. Abe <i>et al.</i>	(CDF Collab.)
ACTON	93H	PL B312 501	P.D. Acton <i>et al.</i>	(OPAL Collab.)
BUSKULIC	93G	PL B311 425	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABREU	92M	PL B289 199	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACTON	92N	PL B295 357	P.D. Acton <i>et al.</i>	(OPAL Collab.)
BUSKULIC	92E	PL B294 145	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
LEE-FRANZINI	90	PRL 65 2947	J. Lee-Franzini <i>et al.</i>	(CUSB II Collab.)